

Mothers should be educated to stop the night breastfeeding after one year of age and shift to use of cup at 10 months of age, and start cleaning the child's mouth and teeth regularly once the first primary tooth has erupted (Bourne, 2007). Furthermore, the American Dental Association (2009) recommended weaning from the breast soon after the child's first birthday. Breastfeeding was recommended up to 2 years of age, with the introduction of appropriate solid food after the age of 6 months to complement the nutrients supplied by breast milk (WHO, 2003; WHO, 1998).

Bottle feeding

Bottle feeding, especially nocturnal feeding is a risk factor for ECC since most contain some form of sugar (Azevedo *et al.* 2005; Twetman, Garcia-Godoy and Goepferd, 2000; Picton and Wiltshear, 1970). In developed countries the primary risk factor for ECC considered to be the use of a "nap time" bottle that contains fermentable carbohydrates such as milk, milk with sugar, sweetened milk with formula, fruit juice or other sweetened solutions (Ribeiro and Ribeiro, 2004). However, under normal conditions milk is not cariogenic and may provide some protection against caries and it is the added sweeteners to the bottle and use of pacifiers dipped in sweetened liquids that leads to the development of ECC (Holt *et al.* 1982).

Bottle feeding all day and night, duration of feeding and sweetened contents are significant risk factors for ECC (Hallet and O'Rourke, 2003; Karjalainen *et al.* 2001; Milgrom *et al.* 2000). Among Syrian children age 3-5 year-olds diagnosed with ECC, bottle feeding occurred in 63%, and breast feeding in 37% (Qadri, Nourallah and Splieth, 2012).

In Lithuania, among 3 year-old children 95% who had S-ECC were sleeping with a bottle containing carbohydrates during the day and night (Slabšinskienė *et al.* 2010). This has been shown to be the most significant risk factor of S-ECC together with a high frequency of sugar intake (Johnsen and Nowjack-Raymer, 1989; Dilley *et al.* 1980). The reduction of the infant salivary flow rate during sleep decreases the oral clearance and increases the contact time of plaque and substrates making the teeth more susceptible to caries (Schafer and Adair, 2000; Seow, 1998; Firestone, 1982).

Parent's education

The level of mother's education has been found to be an important indicator for the development of dental caries in their children. Low mothers' educational levels were associated with severe dental caries in preschool children (Traebert *et al.* 2009).

Dini *et al.* (2000) and Al-Hosani and Rugg-Gunn (1998) showed that the level of parent's education was correlated with the occurrence and severity of ECC, in which low prevalence of dental caries and low mean dmft scores were associated with higher levels of parental education. In another study carried by Subramaniam and Prashanth (2012) thirty eight percent of the children whose mothers were uneducated had high caries prevalence (37.9%) compared to the children whose mothers had higher education.

A higher prevalence of ECC and higher sugar consumption rate was found among Ugandan children who had mothers with lower level of education (Kiwauka *et al.* 2004). A recent study carried out in Sudan showed that there was no significant effect of the mother's education on the prevalence of dental caries: 65% of children whose mothers were uneducated had caries while the prevalence of dental caries was 64.6% in those who had educated mothers (Awooda *et al.* 2013). Sufia *et al.* (2009) showed that the higher the level of maternal education the better the dental health practices of their children. These practices are tooth brushing at correct the time using a tooth brush and toothpaste and sugar consumption control.

Mother's age

In a study carried out in Pakistan to investigate the influence of mother's age, education, occupation and income on the dental health behaviour and caries experience of the preschool children found that there was no association between mother's age and caries experience of the child (Sufia *et al.* 2009). However, children with younger mothers experienced more dental caries than those with older mothers, probably due to the fact that younger mothers paid less attention to the dental health of children (Matilla *et al.* 2000; Paunio *et al.* 1993).

In contrast, Sufia and co-workers (2009) reported that a higher percentage of younger mothers cleaned their children's teeth after meals using tooth brushes and toothpaste and had taken their children to the dentist.

2.7 Risk assessment

The goal of caries-risk assessment in young children is to prevent caries initiation before the first signs of disease (Tinanoff, Kanellis and Vargas, 2002). “Caries risk assessment is the determination of the incidence of caries during a certain time period” (AAPD, 2005). The American Academy of Paediatric Dentistry in (2003b) identified several groups at risk for ECC: (1) children with special healthcare needs, (2) from low socioeconomic and ethno-cultural groups, (3) with suboptimal exposure to topical or systemic fluoride, (4) with poor dietary and feeding habits, (5) whose caregivers and/or siblings have caries, (6) and those with visible caries, white spots, plaque, or decay.

Risk assessment strategies that are most applicable for use in clinical practice, can be easily performed, are inexpensive, require no special equipment or supplies, and provide reliable results. Indicators of caries risk that meet these criteria includes: previous caries experience, presence of white spot lesions or enamel defects, visible plaque, perceived risk by dental professionals, screening tests for mutans streptococci and socioeconomic level of family (Tinanoff *et al.* 2002).

Previous caries experience

One of the best predictors of future caries is previous caries experience (Birkeland *et al.* 1997). Children under the age of 5 years with a history of dental caries should be classified as being at high risk for future decay (Tinanoff and Reisine, 2009).

White spot lesions

White spot lesions are the precursors to cavitated lesions. There are only a few studies that have examined the staining of pits and fissures (Steiner *et al.* 1992), and hypoplastic lesions as caries- risk variables (Li *et al.* 1996).

Visible plaque

The presence of visible plaque on primary teeth can be used as indicator of caries risk. Several studies have shown that there is a correlation between visible plaque on primary teeth and caries risk (Alaluusua and Malmivirta, 1994; Roeters *et al.* 1995).

Perceived risk by dental professional

Dentists without using specific criteria are reasonably able to predict caries risk in children without time or money consuming methods (Alanen *et al.* 1994).

Microbiological testing

The quantity of MS in a child's oral cavity is a risk indicator for dental caries. The count of salivary MS can be determine by simple microbiological testing in the dental clinic and have been reported to have a good specificity (ability to correctly identify those who will not get the disease) but less sensitivity (ability to correctly identify those who will get the disease) (Edelstein and Tinanoff, 1989). In general children who are highly infected with MS have higher caries prevalence and higher caries rate than children with low MS levels (Litt *et al.* 1995).

Socioeconomic status

Socioeconomic status can influence health literacy which in turn can affect general health (Zafar *et al.* 2009).

There is strong evidence to support the association between socioeconomic status and dental caries prevalence. Preschool children from low income families with lower levels of maternal education, especially of illiterate mothers are more likely to have caries (Ann *et al.* 2011; Hashim *et al.* 2009; Maciel *et al.* 2001; Vargas *et al.* 1998; Beck, 1998).

Social status has an indirect influence on caries risk due to its association with certain behaviors such as inappropriate bottle feeding habits, poor oral hygiene (lack of tooth brushing) and high sugar consumption (Litt *et al.* 1995). Eckersley and Blinkhorn (2001) in Northwest England found that children from low income families had visited the dentist less frequently, at an older age and only when there was a dental problem.

2.8 Consequences of ECC

Dental caries in preschool children remain a major problem because it affects high numbers preschool children in both developed and developing countries (Vadiakas, 2008). It occurs five times more frequently than asthma and seven times more commonly than hay fever (DHHS, 2000).

There is strong evidence that untreated dental disease is an important aetiological factor in the pathogenesis of infective endocarditis, which is a condition that carries high mortality rate (Child, 1996). Caries in general and ECC in particular represent a serious public health problem (Begzati, Birisha and Meqa, 2010). It affects normal growth, development and social adaptation of young children (Vadiakas, 2008). It's infectious, and results in impairment of nutrition and esthetics with accompanying psychological problems (Wyne *et al.* 1995).

More than 51 million school hours are lost each year to dental-related illness; poor children suffer nearly 12 times more than rich ones (Manson, 2005).

ECC has been related to failure to thrive, decreased capability to study, increased risk of caries development in the permanent dentition and an increase in the possibility of enamel defects on the successors of carious primary teeth (Broadbent *et al.* 2005; Acs *et al.* 1999; Gray *et al.* 1991).

Rapid progression of dental caries usually causes complications such as pulpitis and apical periodontitis leading to excessive dental treatment such as: pulp therapy and extractions that makes treatment more difficult (Vinckier *et al.* 2001). Moreover, early extraction of primary teeth may predispose to malocclusion in the permanent dentition (Greenwell *et al.* 1990). ECC was shown to adversely affect the child's development, especially body weight and height (Ayhan, Suskan and Yildirim, 1997). Severe forms of ECC may require treatment under general anaesthesia, and the treatment cost is high (Peressini *et al.* 2004).

2.9 Treatment of ECC

Most researchers agree that the best approach to the “treatment” of ECC is prevention, since treatment of ECC is expensive and mainly carried out under general anaesthesia or sedation and this is risky for children (Peretz and Eidelman, 2003). Treatment of ECC includes different types of interventions depending on the progression of the disease, the age of the child, and the social, behavioural and medical history of the child (Zafar *et al.* 2009).

2.10 Prevention of ECC

Dental Health education

Educational programmes have been used to decrease dental caries in children (Gallagher and Rowe, 2001; Gomes, Fonseca and Rodrigues, 2001). There are three key messages currently used in dental health education: reduce the frequency of sugar intake, brush teeth regularly with a fluoridated toothpaste and visited a dentist regularly (Health Education Authority, 1989). People should be motivated to use fluoridated toothpaste with correct method of tooth brushing for effective removal of dental plaque (Ramya *et al.* 2011).

Oral health education preventive programmes for preschool children often involve education of the parent or carer. These educational messages attempt to persuade parents/carers not to put children to bed with a bottle, to reduce the child’s high frequency consumption of sugars, and to brush children’s teeth daily.

Burt and Eklund. (1999) have suggested that while educational programmes may improve knowledge, they only have a temporary effect on plaque levels and no marked effect on caries experience. Despite these limitations, oral health education undoubtedly remains an important component of preventive dental programmes (Tinanoff *et al.* 2002).

Dental visit

The first visit to dentist should occur before the eruption of the first tooth, so that parents can be taught how to clean their infant’s teeth. During the first dental visit, a risk assessment is necessary to counsel the parent on how to prevent dental caries (Zafar *et al.* 2009).

In a study carried in Thailand, 88% of children had never visited a dentist and for those who visited dentist the main reason was pain (Senesombath *et al.* 2010). Lo, Loo and Lee. (2009) in Hong Kong found that only 26% of children had visited a dentist and in the United States, Kopycka- Kedzierawski and Billings (2011) found that about thirty percent of children had never had a dental check-up.

Fluoride and dental caries

The action of fluoride is in the process of remineralisation and inhibition of demineralisation of enamel (Featherstone, 2000). Topical application of fluoride is most effective in the prevention of ECC (Sanchez and Childers, 2000).

Community water fluoridation

The effectiveness of water fluoridation in reduction of dental caries is 50-70% in children (CDC, 1999). The optimum concentration of fluoride is in the range of 0.7 mg/L-1.2 mg/L (CDC, 2001). Water fluoridation has been shown to be one of the most cost-effective means of reducing tooth decay in children (Klein *et al.* 1985). In studies examining the before and after measurements of caries, starting or continuing fluoridation decreased dental caries experience among children aged 4 to 17 years by a median of 29.1% during 3 to 12 years of follow-up (Kumar and Moss, 2008).

Fluoride varnish

The application of fluoride varnish every 6 months is effective in preventing dental caries in both primary and permanent teeth of children and adolescents (ADA, 2006) and it reduces dental caries by a third (Marinho *et al.* 2002). The varnishes must be applied by a professional to maintain their cariostatic effect.

Salt fluoridation reduces the prevalence of dental caries to be about 50% (Estupinan *et al.* 2001).

Fluoride tablets

Fluoride tablets provide both systemic and topical effects and it reduces the new carious lesions for up to 30% (Allukian and Horowitz, 2003; Stephen, 1993).

Fluoridated toothpaste

Proper oral hygiene and use of fluoridated toothpaste are the most important factors in the prevention of dental caries (Slabsinskiene *et al.* 2010). To prevent ECC in preschool children by home-care approaches, brushing with a small amount of fluoridated toothpaste is effective when it starts soon after tooth eruption (Pine *et al.* 2000).

The efficacy of fluoridated toothpastes in the prevention of dental caries in children and adolescents has been reported (Marinho *et al.* 2003; Mohan *et al.* 1998). Fluoridated toothpastes have contributed considerably to recent declines in the incidence of dental caries (Naylor and Murray, 1983). The use of a “pea-size” of toothpaste, low in fluoride concentration (<500 ppm) has been recommended by the European Academy of Paediatric Dentistry (Oulis *et al.* 2000).

In Nigerian preschool children aged 6-71 months, more than half of children brushed their teeth with toothbrush and paste (Sowole *et al.* 2007).

Nearly all preschool children aged 3-5 years in Uganda brushed their teeth at least daily with fluoridated toothpaste (Kiwauka *et al.* 2004).

Results of a similar study in Lithuania showed that 62.5% of the parents who have caries-free children started to brush their teeth soon after the eruption of the first tooth, and 52.5% of the parents having children with S-ECC did not brush their children's teeth at all (Slabsinskiene *et al.* 2010).

Lo *et al.* (2009) found that children with low dmft scores are those who started tooth brushing at an earlier age and those who brushed their teeth more frequently (Lo *et al.* 2009). A high dmft score was reported for children who did not brush their teeth at all (Begzati *et al.* 2010). Senesombath *et al.* (2010) found that children aged 36-47 months with fewer dental caries are those with higher tooth brushing frequency and those whose parents or caregivers help and supervise them during brushing.

Other studies have shown that daily tooth brushing with fluoridated toothpaste in 3-6 year olds significantly reduces caries incidence (Schwarz *et al.* 1998; Sjogren *et al.* 1995).

Marinho *et al.* (2003) in a systematic review concluded that supervised tooth brushing with fluoridated toothpaste containing 1000-1100 ppm fluoride will reduce the prevalence of dental caries in children by 32%. In another systematic review of 70 controlled clinical trials Marinho *et al.* (2009) found that tooth brushing was associated with 24% reduction of dental caries. Lower percentages of dental caries were found in children who used fluoridated dentifrices (Subramaniam and Prashanth, 2012).

Children who have an adult assisting with brushing are more likely to be caries-free than children who brushed their own teeth (Peretz and Eidelman, 2003). Preschool children whose parents began caring earlier for their teeth within the first year of life, and those whose parents helped them during tooth brushing have better dental health status and lower dmft scores (1.56) (Pieper *et al.* 2012).

In India, preschool children who started brushing their teeth after 1 year of age showed a high severity of dental caries, and less dental caries was found among children who brushed their teeth twice daily (Retnakumari and Cyriac, 2012). Gibson and Williams (1999) concluded that the consumption of sugars and sugary foods in children who brushed their teeth twice a day or more did not appear to be associated with caries. Similar findings were reported by Burt and Pai (2001) that the consumption of sugars remains a moderate risk factor for caries in those who had adequate exposure to fluoride.

The results of a study in children aged 6-60 months showed that 9% of children at age 60 months were brushing their teeth less than once per day, 47% were brushing once a day, and 43% were brushing 2 or more times per day (Franzman *et al.* 2004). In the same study Franzman *et al.* (2004) suggested that parents with high educational levels were brushing their children's teeth more frequently than those with lower educational levels. Parents should supervise children aged 2-7 years during brushing and should ensure that only a small pea sized amount of fluoridated toothpaste is used and that swallowing of the paste is avoided (DHC, 2002).

An evaluation of a supervised tooth brushing program for 5 and 6 year olds children in England found a reduction of 11% in the caries increment (Jackson *et al.* 2005).

This review has shown that dental caries is a public health problem among preschool children all over the world. Although it is preventable disease, dental health education, use of fluoride especially a supervised tooth brushing with a fluoridated toothpaste and regular dental visit are the main preventive measures for dental caries.



CHAPTER 3: AIM and OBJECTIVES

3.1 Aim

To determine the prevalence of dental caries and tooth brushing habits among 3 to 5 year-old preschool children in Khartoum State, Sudan.

3.2 Objectives

To determine:

1. The dmft of the sample
2. The proportion of caries free teeth
3. Tooth brushing frequency

and

4. To compare dmft and tooth brushing habits
5. To make recommendations to Directorate of Oral Health- Ministry of Health- Khartoum State.



CHAPTER 4: METHODOLOGY

4.1 Study design

This study is a descriptive, cross sectional study of prevalence of dental caries and tooth brushing habits among preschool children 3 to 5 year-olds.

4.2 Study site and study population

This study was carried out among a systematically randomly selected kindergarten schools in the 7 localities of Khartoum state: Khartoum, Jabal Awlia, Om Dorman, Umbada, Karari, Khartoum North and Sharq Elnil. These 7 localities comprise 3,133 kindergarten schools. Basic information regarding the schools was obtained from the Ministry of Education, Khartoum state and more detailed information regarding the number and age of the children was obtained from the localities.

4.3 Sample size and sample

The sample size for pre-school children was calculated according to the following formula:

$$n = \frac{z^2 pq}{d^2} \cdot deff$$

Where:

n= sample size

z= (1.96) value of normal curve corresponding to level of confidence (95%)

p= (0.607) proportion of students with caries in pre-school children

q= 1-p (0.393) students with no caries in basic public schools.

d= desired margin of error (5%)

deff = design effect (1.5)

$$n = \frac{(1.96)^2 * 0.607 * 0.393}{(0.05)^2} = 367$$

$$(0.05)^2$$

$$n * deff = 367 * 1.5 = 551 \approx 555$$

Twenty eight kindergarten schools were randomly selected, and 20 children aged between 3-5 years were selected from each school.

4.4 Inclusion criteria

- Preschool children attending kindergartens in Khartoum- State aged 3-5 year-olds.
- Kindergarten schools had more than 20 children in the 3-5 year age group.

4.5 Data collection

Data was collected by using two instruments: a data capture sheet modified from WHO assessment form (1997) for clinical dental examination (Appendix 1) and a structured administered questionnaire for mothers/guardians (Appendix 2). The questionnaire was written in English but administered in Arabic to the mother by the researcher during the interview.

4.5.1 Cross infection control measures

A sterile set of instruments was used for each child. Gloves were changed before examination on every subject. Face mask, sterile cotton rolls also had been used.

Used instruments were collected and putted in a separate box and then washed and autoclaved at the end of the working day. Used gloves, masks, tissue papers and pouches were disposed in a waste bag.

4.5.2 Questionnaire

The dental health related habits were assessed using structured administered questionnaire that was completed by interviewing the mothers/guardians (n=553) (Appendix 2). The questionnaire obtained information on the knowledge about the causes of dental caries, tooth brushing habits of their children (age of starting brushing, frequency, tool of brushing, help of mother during brushing and fluoride

content in the tooth paste), breast feeding and bottle feeding habits, dietary habits and dental visits.

The questionnaire was piloted prior to the main study on 12 mothers/guardians to check the clarity and ease of understanding. Following the pilot, some of the questions were reformulated in the final draft.

4.5.3 Clinical dental examination

Clinical dental examinations were carried out at the kindergarten schools by a calibrated dentist with the child seated in an upright position. All children were examined with a dental mirror and a probe in the class room of the school using natural daylight.

4.5.3.1 Diagnostic criteria

The criteria for the diagnosis of dental caries were according to the criteria of the WHO (1997).


Dentition coding was as follows:

1) Sound crown =A

A crown was recorded as sound if it showed no evidence of untreated clinical caries.

The stages of caries that precede cavitation, as well as other conditions similar to the early stages of caries, are excluded because they cannot be reliably diagnosed. Thus, crown with the following defects, in the absence of other positive criteria, were coded as sound:

1. White or chalky spots.
2. Discoloured or rough spots.
3. Stained pits or fissures in the enamel that catch the explorer but do not have a detectably softened floor, undermined enamel, or softening of the walls.
4. Dark, shiny, hard, pitted areas of enamel in a tooth showing signs of moderate to severe fluorosis.
5. Lesions that, on the basis of distribution or history, appeared to be due to abrasion.



2) Decayed tooth = B

Caries is recorded as present when a lesion in a pit or fissure, or on a smooth tooth surface, has an unmistakable cavity, undermined enamel, or a detectably softened floor or wall. A crown with a temporary filling, or one which is sealed but also decayed should also be included in this category.

3) Filled with decay = C

A crown was recorded as filled with decay, when it has one or more permanent restorations and one or more areas that are decayed.

4) Filled no decay = D

A crown was considered filled with out decay, when one or more permanent restorations are present and there is no caries anywhere on the crown. A tooth that has been crowned because of previous decay is recorded in this category. A tooth that has been crowned for reasons other than decay (e.g. a bridge abutment).

5) Missing tooth, as a result of caries = E

This code was used for tooth that has been extracted because of caries and was recorded under coronal status.

6) Not recorded = F

This code was used for any tooth that could not be examined for any reason.

4.5.3.2 Calibration

Prior to the clinical dental examinations, the examiner was calibrated on a group of pre-selected children (12 children were examined) who possessed the same characteristics to be assessed in the main study in order to assess intra-examiner agreement. The kappa statistic was 0.925.

4.6 Data analysis

The collected data from the dental examination and from the questionnaire was categorized, coded and entered into a computer. The data was captured in Excel.

Basic descriptive analysis was done using the Excel environment. The database was imported into the statistical package for social sciences (SPSS), version 11.5 computer software programs to perform complex statistical analysis.

Descriptive statistics was used to describe the demographic factors. The tests of significance used were independent sample t-test, Chi-square test and one way ANOVA test.

4.7 Ethical considerations

The protocol was submitted to the Senate Research Ethics Committee of the University of The Western Cape for ethical approval and permission to carry out the study.

Participation was voluntary and they were informed that they were free to withdraw from the study at any time and that all information gathered from the study would be kept strictly confidential


Two informed consent forms (Appendix 3 and 4) were signed by the mother/ guardian prior to the interview and before the oral examination. The examination of the study population did not include any radiographic examination. After clinical dental examinations mother/ guardian were informed about any necessary dental treatments and those who needed treatment were referred for the necessary treatments to the dental clinic of their choice.

CHAPTER 5: RESULTS

5.1 Demographic information

Five hundred and fifty three preschool children aged between 3 and 5 years old participated in the study with their mothers or guardians (n=553). The demographic characteristics of the study population are presented in Figure 4. Girls (n= 287) slightly outnumbered boys (n= 266).

The calculated statistical sample size was 551 children and this was rounded to 555 children and the final sample size was 553 since two unusable questionnaires were excluded from the data analysis.



About half of the mothers were aged between 30 and 39 years old, 31.5% had a university level education and 8% were uneducated (Table 1).

Table 1: Distribution of mothers/guardians according to age and level of education

		n	Percent
Age	<20 years	6	1.1
	20-29 years	210	38.0
	30-39 years	277	50.1
	≥40 years	60	10.8
Level of education	No education	45	8.1
	Primary/basic	143	25.9
	Intermediate school	18	3.3
	High school	155	28.0
	University	174	31.5
	Postgraduate	18	3.3
Total		553	100

5.2 Caries Experience (Children)

The caries experience was 52.4% with a mean dmft of 2.27(11.09% decayed, 0.3% missing due to caries and 0.03% filled). The decayed component comprised the bulk of the mean dmft and the filled component the least (Table 2). The most commonly decayed tooth was the lower right second molar (9.6%).

Table 2: Percentage of decayed, missing and filled teeth

dmft Status	Percentage
Caries-free	88.58
Decayed	11.09
Filled	0.03
Missing as a result of caries	0.3
Total	100

There was an increase in the dmft scores with increasing age, with the mean dmft being 1.35, 2.18 and 2.79 in the 3, 4 and 5 year olds respectively. There was no significant difference in the mean dmft scores between males and females (Table 3).

Table 3: Mean dmft according to age and gender

	n	Mean dmft	Std. Deviation
Age			
3 years	103	1.53	2.46
4 years	254	2.18	3.24
5 years	196	2.79	4.04
Gender			
Male	266	2.28	3.32
Female	287	2.27	3.57
Total	553	2.27	3.45

5.3 Dental Caries and Mothers' characteristics


As shown in Table 4, children with mothers younger than 20 years old had lower dmft scores than those with mothers older than 20 years.

Table 4: Association between age of mothers and dmft

	Frequency	Mean dmft	Std. Deviation
<20 years	6	3.33	3.50
20-29 years	210	2.24	3.60
30-39 years	277	2.24	3.29
≥40 years	60	2.43	3.66
Total	553	2.27	3.45

Chi-square test performed, p-value = 0.865

The differences in dmft between of children of uneducated and educated mothers were very variable and not significant (Figure 5).



Chi-square test performed, p -value = 0.213

5.4 Tooth brushing habits

The number of children who brushed their teeth regularly (at least twice a day) was 490 (88.6%) (Table 5).




Table 5: Does your child brush his/her teeth?

	n	Percent
Regularly brush	490	88.6
Sometimes brush	53	9.58
Not brush	10	1.8
Total	553	100

According to Table 6, only 6.8% of mothers reported that they started brushing soon after the first tooth erupted, while 92.6% only after their child turned 3 years old. 83.3% of children were brushing their teeth once a day and 15.1% were brushing twice a day (Table 6). Table 6 also depicts information regarding tooth brushing tools, mother assistance during brushing, using fluoridated toothpaste and spitting out the toothpaste after brushing.

Table 6: Tooth brushing habits

	n	Percent
Age of starting tooth brushing		
Soon after the first tooth came out	37	6.81
When they turned 1 year old	101	18.60
When they turned 2 years old	233	42.90
When they turned 3 years old	132	24.30
Others	40	7.36
Tooth brushing frequency		
Once a day	453	83.4
Twice a day	82	15.1
Three times a day	8	1.5
Tooth brushing Tools		
Toothbrush	542	99.9
Miswak	1	0.001
Mother helps during brushing		
Yes	285	52.5
Sometimes	100	18.4
No	158	29.1
Knowledge of using fluoridated toothpaste		
No	8	1.47
Yes	226	41.62
Don't know	309	56.90
Spitting out the toothpaste		
Yes	488	89.87
No	45	8.28
Don't know	10	1.84
Total	543	100

By comparing the age of starting brushing with dmft, it was seen that children whose mothers started to brush their teeth at an earlier age, had lower dmft scores than those whose mothers started to brush their teeth after 2 years (Table 7).

Table 7: Association between dmft and age starting brushing

	N	Mean dmft	Std. Deviation
Soon after the first tooth came out?	37	1.97	2.53
When they turned 1 year old	101	1.86	2.97
When they turned 2 years old	233	2.21	3.14
When they turned 3 years old	132	2.96	4.53
Others	40	2.20	2.96
Total	543	2.31	3.47

One way ANOVA test performed, p-value = 0.139 (not significant)

The mean dmft decreased with the increased frequency of tooth brushing. Lower dmft scores were found among children who brushed their teeth three times a day (Table 8).

Table 8: Association between dmft and tooth brushing frequency

	n	Mean dmft	Std. Deviation
Once a day	453	2.35	3.49
Twice a day	82	2.21	3.44
Three times a day	8	1.25	2.12
Total	543	2.31	3.47

One way ANOVA test performed, p-value = 0.646 (not significant)

No statistically significant difference was found between the mother help during brushing and the mean dmft of their children (Table 9).

Table 9: Association between dmft and mother help during brushing

	n	Mean dmft	Std. Deviation
Yes helping children	285	2.45	3.63
Sometimes helping children	100	1.92	2.77
Not helping children	158	2.31	3.56
Total	543	2.31	3.47

One way ANOVA test performed, p-value = 0.423


There was no significant difference between the mean dmft scores of children whose mothers were sure that they were using a toothpaste containing fluoride and children whose mothers did not (Table 10).

Table 10: Association between dmft and use of fluoridated tooth paste

	N	Mean dmft	Std. Deviation
Using fluoridated toothpaste	226	2.27	3.15
Not using fluoridated toothpaste	8	1.38	2.07
Don't know	309	2.37	3.72
Total	543	2.31	3.47

One way ANOVA test performed, p-value=0.703

Figure 6 shows that the highest percentage (99.3%) of children brushed their teeth in the morning and 14.9% were brushed at night.



There was no statistical difference between the dmft and the time of brushing per day, however, children who brushed their teeth during the day had a lower dmft score 1.12 (Table 11).

Table 11: Association between dmft and time of tooth brushing during the day

	Frequency	Mean dmft	Std. Deviation	<i>p-value</i>
Morning	539	2.32	3.48	0.746
During the day	17	1.12	2.34	0.15
At night	81	2.17	3.39	0.697
After meals	4	3.25	2.22	0.587

One way ANOVA test performed, p-value>0.05(not significant)

5.5 Tooth brushing and Mother characteristics

The highest percentage of children (77.78%) who brushed their teeth regularly was found among the children whose mothers had a post-graduate degree and the highest proportion of children who did not brush their teeth regularly was from uneducated mothers and this was statistically significant (Table 12).

Table 12: Association between mother level of education and tooth brushing habit

Level of education	Does your child brush his /her teeth?			Total
	Yes	Sometimes	No	
None	17	5	21	43
%	39.53	11.63	48.84	100
Primary/basic education	71	22	48	141
%	50.35	15.60	34.04	100
Intermediate school	12	1	4	17
%	70.59	5.88	23.53	100
High school	74	32	47	153
%	48.37	20.92	30.72	100
University	97	38	36	171
%	56.73	22.22	21.05	100
Postgraduate	14	2	2	18
%	77.78	11.11	11.11	100
Total	285	100	158	543
%	52.49	18.42	29.10	100

Chi square test performed, p-value=0.007

There was no significant association between the mother level of education and the age at which the child started brushing their teeth (p -value= 0.058). More than 42% of mothers started brushing their child's teeth after they turned 2 years old. Around 22% of postgraduate mothers started brushing their children teeth soon after the first tooth erupted (Table 13).

Table 13: Association between the mother level of education and age of starting tooth brushing for children

Level of education	Age of starting tooth brushing					Total
	Soon after first tooth erupted	When they turned 1 year old	When they turned 2 years old	When they turned 3 years old	Others	
None	2	11	12	13	5	43
%	4.65	25.58	27.91	30.23	11.63	100
Primary/basic education	9	16	64	45	7	141
%	6.38	11.35	45.39	31.91	4.96	100
Intermediate school	2	2	6	6	1	17
%	11.76	11.76	35.29	35.29	5.88	100
High school	8	27	70	34	14	153
%	5.23	17.65	45.75	22.22	9.15	100
University	12	41	75	31	12	171
%	7.02	23.98	43.86	18.13	7.02	100
Postgraduate	4	4	6	3	1	18
%	22.22	22.22	33.33	16.67	5.56	100
Total	37	101	233	132	40	543
%	6.81	18.60	42.91	24.31	7.37	100

Chi square test performed, p-value=0.058

UNIVERSITY of the
WESTERN CAPE

Ninety percent of children with uneducated mothers brushed their teeth once a day. The highest proportion of children who brushed twice a day was found among the children whose mothers had a post- graduate education (Table 14).

Table 14: Association between mother level of education and brushing frequency

Level of education	Brushing frequency			Total
	Once a day	Twice a day	Three times a day	
None educated	39	2	2	43
%	90.70	4.65	4.65	100
Primary/basic education	110	28	3	141
%	78.01	19.86	2.13	100
Intermediate school	17	0	0	17
%	100	0	0	100
High school	130	20	3	153
%	84.97	13.07	1.96	100
University	143	28	0	171
%	83.63	16.37	0	100
Postgraduate	14	4	0	18
%	77.78	22.22	0	100
Total	453	82	8	543
%	83.43	15.10	1.47	100

Chi square test performed, p-value=0.073

The association between the mother's level of education and the use of fluoridated toothpaste was highly significant (p -value=0.001). The highest percentage of mothers that used fluoridated toothpaste was found among the postgraduate educated mothers (77.78%) and the lowest among the uneducated mothers (9.3%) (Table 15).

Table 15: Association between mother level of education and use of toothpaste that contains fluoride

Level of education	Does your child use toothpaste that contains fluoride?			Total
	Yes	No	Don't know	
None educated	4	2	37	43
%	9.30	4.65	86.05	100
Primary /basic education	30	1	110	141
%	21.28	0.71	78.01	100
Intermediate school	2	0	15	17
%	11.76	0.00	88.24	100
High school	69	3	81	153
%	45.10	1.96	52.94	100
University	107	2	62	171
%	62.57	1.17	36.26	100
Postgraduate	14	0	4	18
%	77.78	0.00	22.22	100
Total	226	8	309	543
%	41.62	1.47	56.91	100

Chi square test performed, p-value=0.001

There was no significant difference in the association between the mother's age and the assistance of mother during brushing. All mothers younger than 20 years old helped their children to brush their teeth (Table 16).

Table 16: Association between mother age and mother help during brushing

Mother age	Do you help your child to brush his/her teeth?			Total
	Yes help n (%)	Sometimes help n (%)	Not help n (%)	
<20	4 (66.67)	0	2 (33.33)	6 (100)
20-29	101 (48.79)	41 (19.81)	65 (31.40)	207 (100)
30-39	148 (54.81)	52 (19.26)	70 (25.93)	270 (100)
≥40	32 (53.33)	7 (11.67)	21(35)	60 (100)
Total	285 (52.49)	100 (18.42)	158 (29.1)	543 (100)

Chi square test performed, p-value =0.42

5.6 Causes of tooth decay

Table 17 depicts the knowledge of mothers regarding the causes of dental caries. 'Too much sugar' was reported to be the main cause of dental caries (82.8%).

Table 17: Causes of tooth decay

	n	Percent
Too much sugar	458	82.8
Bad oral hygiene	395	71.4
Germ	188	34
Others	21	3.8

5.7 Feeding habits

The vast majority of mothers (97.28%) reported breast feeding their children and about 25% of children were bottle-fed. The percentage of both breast-fed and bottle-fed children together was 22.9% (Table 18).

Table 18: Feeding habits

	n	Percent
Breast feeding		
Yes	538	97.28
No	15	2.71
Bottle feeding		
Yes	140	25.31
Sometimes	6	1.08
No	407	73.59
Breast and bottle feedings	127	22.96
Total	553	100

The association between the dmft and breast feeding showed that the mean dmft was slightly lower in children who were breast fed (Table 19).

Table 19: Association between breast-feeding and dmft

Did you breast-feed your child?	n	Mean dmft	Std. Deviation
No breast- fed	15	3.4	3.07
Breast- fed	538	2.24	3.46

Independent samples T-test performed, p-value=0.200

No significant difference was found in dmft scores between children who were bottle-fed and children who were not bottle-fed (Table 20).

Table 20: Association between bottle-feeding and dmft

	n	Mean dmft	Std. Deviation
Not bottle-fed	407	2.21	3.47
Bottle-fed	140	2.51	3.41

Independent samples T-test performed, p-value=0.376

Data on bottle content, frequency of bottle feeding per day, sugar addition to the bottle and nocturnal use of bottle are presented in Table 21.

Table 21: Bottle-feeding characteristics

	n	Percent
What do/did you put in the bottle (milk)?		
Yes	135	92.5
No	11	7.5
What do/did you put in the bottle (juice)?		
Yes	82	56.2
No	64	43.8
What do/did you put in the bottle (other)?		
Yes	4	2.7
No	142	97.3
Bottle-feeding frequency		
Once	16	11
Twice	37	25.3
Thrice	41	28.1
More than three times	52	35.6
Do/did you add sugar to the contents in the bottle?		
Yes	94	64.4
Sometimes	6	4.1
No	46	31.5
Does your child fall asleep with the bottle in his/her mouth?		
Yes	37	25.3
Sometimes	6	4.1
No	103	70.5
Total	146	100

The association between the dmft and the content of bottle was not statistically significant (Table 22).

Table 22: Association between the bottle-content and dmft

	n	Mean dmft	Std. Deviation	sig
Milk				
Yes	135	2.56	3.44	0.167
No	11	1.09	2.12	
Juice				
Yes	82	2.18	3.15	0.299
No	64	2.78	3.64	
Other				
Yes	4	0.25	0.50	0.188
No	142	2.51	3.40	

Independent samples T-test performed, P-value>0.05

When dmft was compared to the frequency of bottle feeding there was no statistically significant difference. The highest dmft scores were found among children who were bottle-fed three and more times per day (Table 23).

Table 23: Association between bottle-feeding frequency and dmft

	n	Mean dmft	Std. Deviation
Once a day	16	2.25	3.51
Twice a day	37	1.68	2.39
Thrice a day	41	2.88	3.21
More than three times a day	51	2.76	4.02
Total	145	2.46	3.38

One way ANOVA test performed, p-value=0.381

Table 24 shows that there was no association between dmft scores and the addition of sugar to the bottle content.

Table 24: Association between adding sugar to the bottle content and dmft

Adding sugar	n	Mean dmft	Std. Deviation
Yes	94	2.34	3.59
Sometimes	6	3.83	2.23
No	46	2.48	3.05
Total	146	2.45	3.38

One way ANOVA test performed, p-value=0.578

Higher dmft scores were found among children who sleep with a bottle (Table 25), but this was not significant.


Table 25: Association between falling sleep with the bottle in a mouth and dmft

	N	Mean dmft	Std. Deviation
Yes	37	3.00	3.37
Sometimes	6	2.00	1.90
No	103	2.27	3.45
Total	146	2.45	3.38

One way ANOVA test performed, p-value=0.506

5.8 Sugar consumption

Nearly 30% eat sugar-containing food more than three times per day (Figure 7 and 8). More than 95% of the children were eating sugars every day (Figure 9), Sweets and chocolates were most commonly eaten.




Eating sugary food was significantly associated with higher dmft scores (Table 26).

Table 26: Association between eating sugar containing food and dmft

Eating sugary food?	Frequency	Mean dmft	Std. Deviation
Yes	530	2.36	3.49
Sometimes	21	0.33	0.80
No	2	0.50	0.71
Total	553	2.27	3.45

One way ANOVA test performed, $p\text{-value}=0.023$




There was a statistically significant difference between foods containing sugar (sweets and chocolates) and the dmft scores (Table 27).

Table 27: Association between type of sugar containing food and dmft

	n	Mean dmft	Std. Deviation	p-value
Sweets				0.001
Yes	518	2.36	3.53	
No	33	1.06	1.48	
Chocolates				0.001
Yes	407	2.54	3.66	
No	144	1.56	2.67	
Biscuits and cookies				0.237
Yes	460	2.35	3.52	
No	91	1.92	3.07	

Independent samples T-test performed



A highly significant difference was found between the mean dmft and the frequency of eating food containing sugar per day. The dmft scores increased with the increasing number of times sugar was eaten per day ranging from 1.34 once per day to 2.8 for three and more times per day (Table 28).

Table 28: Association between frequency of eating sugary food and dmft

	n	Mean dmft	Std. Deviation
Once a day	94	1.43	2.40
Twice a day	157	1.80	3.01
Thrice a day	137	2.80	4.00
More than three times a day	163	2.80	3.72
Total	551	2.28	3.45

One way ANOVA test performed, p-value=0.001

5.9 Dental visits

Dental visits were not common as only about 9% of children had visited a dentist and the most common reason for the visit was toothache (Table 29 and 30). In 41.2% of children the dentist only did an examination of the mouth and teeth (Table 31).

Table 29: Dental visits

	n	Percent
Visited dentist	51	9.22
Not visited dentist	502	90.78
Total	553	100

Table 30: Reasons for taking child to the dentist

	n	Percent
Routine check-up	8	15.7
Toothache	25	49
Others	18	35.3
Total	51	100

Table 31: What did the dentist do?

	n	Percent
Only an examination	21	41.2
Pulled out the tooth/teeth	13	25.5
Did a filling	7	13.7
Other	16	31.4

Table 32: Association between mother's age and dental visit

Age	Dental visit		
	Visited	Not visited	Total
<20	0	6 (100)	6 (100)
20-29	18 (8.57)	192 (91.43)	210 (100)
30-39	27 (9.75)	250 (90.25)	277 (100)
≥40	6 (10)	54 (90)	60 (100)
Total	51 (9.22)	502 (90.78)	553 (100)

Chi square test performed, p-value =0.837

Table 32 depicts the relationship between the age of mothers and dental visit. All children with uneducated mothers never visited a dentist. The highest number of dental visits was found among children whose mothers were university graduated (27%) while the lowest was among children with uneducated mothers (2.2%) (Table33).

Table 33: Association between mother level of education and dental visit

Level of education	Have you ever taken your child to the dentist?		
	Yes n (%)	No n (%)	Total n (%)
Not educated	1 (2.22)	44 (97.78)	45 (100)
Primary/basic education	9 (6.29)	134 (93.71)	143 (100)
Intermediate school	2 (11.11)	16 (88.89)	18 (100)
High school	10 (6.45)	145 (93.55)	155 (100)
University	27 (15.52)	147 (84.48)	174 (100)
Postgraduate	2 (11.11)	16 (88.89)	18(100)
Total	51 (9.22)	502 (90.78)	553 (100)

Chi square test performed, p-value =0.016

CHAPTER 6: DISCUSSION

Dental caries in preschool children is a public health problem although it is a preventable disease and one in which prevention should begin early in life. The present study investigated the prevalence of dental caries and tooth brushing habits in a random sample of preschool children in Khartoum State, Sudan. The prevalence of dental caries was measured using the WHO criteria (1997) and information on the causes of dental caries, tooth brushing habits, feeding habits, dietary habits and dental visits was obtained by interviewing the mothers/guardians of the children using a structured administered questionnaire.

The prevalence of dental caries varies widely throughout the world and it is not easy to compare the prevalence of ECC between the different studies due to the differing measurement criteria that are used. Studies are conducted for a variety of reasons, each with different aims and objectives. The external validity of the findings is therefore restricted i.e. the extent to which inferences can be made is limited to specific populations and may not necessarily reflect local, regional or national trends. This is important as there may be considerable variations in the prevalence and manifestation of ECC within sub-populations in an area as diverse as sub-Saharan Africa.

The prevalence of dental caries among the preschool children in the present study was 52.4% in Khartoum State, Sudan. The mean dmft was 2.27 and according to the WHO criteria this mean dmft is low. A study in Khartoum State a decade ago, by Raadal *et al.* (1993) reported a lower dmft score of 1.68 and a more recent 2013 study in the same age group found a higher dmft of 3.53 (Awooda *et al.* 2013). This big variation in the dmft scores seems to be due to the different sample selection criteria.

Internationally, the mean dmft in studies carried out in different countries in Asia was found to range from a low of 1.5 in Hong Kong and 1.85 in Pakistan to 5.5 in Thailand (Lo *et al.* 2009; Sufia *et al.* 2011; Senesombath *et al.* 2010).

In the present study the mean dmft was 1.35, 2.18 and 2.79 among 3, 4 and 5 year-olds respectively, while the results of a study in Uganda showed higher dmft scores of 1.7, 2.4 and 3.1 among the same age groups (Kiwauka *et al.* 2004). In these two studies the dmft was directly proportional to the age.

Wyne *et al.* (1995) showed that in Riyadh, Saudi Arabia the mean dmft among 4 to 6 year-old preschool children was 6.9 which is very high when compared to the present study. The mean dmft increased dramatically with age, and the finding is similar to those reported by Awooda *et al.* (2013), Sufia *et al.* (2011), Begzati *et al.* (2010) and Lo *et al.* (2009).

The present study concurred with other studies that reported no gender difference in the caries prevalence in preschool children (Awooda *et al.*, 2013; Slabšinskienė *et al.*, 2010; Kiwanuka *et al.* 2004; Kerosuo and Honkala, 1991), as there was no difference in the mean dmft scores between boys (2.28) and girls (2.27).

There is conflicting evidence regarding the association between the mother's age and caries experience of the child. A study from Pakistan reported no association (Sufia *et al.* 2009), while other studies found that children with younger mothers experienced more dental caries possibly because the younger mothers paid less attention to the dental health of children (Matilla *et al.* 2000; Paunio *et al.* 1993). The present study found that children with mothers younger than 20 years old had lower dmft scores than those with mothers older than 20 years and this may be due to the fact that younger mothers had fewer children so they were better able to take care of their oral health. However, this finding may not be generalizable to the population at large since the sample size was small.

The level of a mother's education has been reported to be a risk factor for the development of dental caries in their children. Maternal education has been correlated with the occurrence and severity of ECC, in which low prevalence of dental caries and low mean dmft scores were associated with higher levels of parental education (Subramaniam and Prashanth, 2012; Traebert *et al.* 2009; Kiwanuka *et al.* 2004; Dini, *et al.* 2000; Al-Hosani and Rugg-Gunn, 1998). The contrasting finding of the present study, where the dmft of children with uneducated mothers was nearly half that of educated mothers, could be related to the small sample sizes. Nonetheless, since educated mothers have better incomes than uneducated mothers it could be that they are able to afford a more 'cariogenic' diet (sweets and chocolates) for their children. Other studies have reported that there is no significant effect of the mother education on the dental caries (Awooda *et al.* 2013; Sufia *et al.*, 2009).

Tooth brushing with fluoridated toothpaste is important for the prevention of dental caries in children. To prevent ECC in preschool children by home-based care approaches, brushing with small amount of fluoridated tooth paste is effective when it starts soon after teeth eruption (Pine *et al.* 2000). In the present study a high percentage of children were brushing their teeth on a daily basis and more than 90% of the mothers reported that they started tooth brushing for their children after they turned 3 years old. Children, whose mothers started to brush their teeth at an earlier age, showed slightly lower dmft scores than those whose mothers started to brush their teeth after 2 years, however the difference was not statistically significant. Similar findings have been reported by other studies (Pieper *et al.* 2012; Retnakumari and Cyriac, 2012; Slabsinskiene *et al.* 2010; Lo *et al.* 2009).

Furthermore, children who brushed their teeth more than once per day experienced lower caries levels. These results were similar to those of Senesombath *et al.* (2010) and Retnakumari and Cyriac. (2012). However, it was found that mother support during brushing was not associated with the mean dmft of their children. The dmft scores were similar between children who were assisted by mothers during brushing and those who were not assisted. However, these findings do not concur with the findings of the systematic review of Marinho *et al.* (2003) and with other studies (Pieper *et al.* 2012; Peretz and Eidelman, 2003).

The results of a study by Sufia *et al.* (2009) showed that the higher the level of maternal education the better the dental health practices of their children. These practices include tooth brushing at the right time using tooth brush and toothpaste and sugar consumption control. These results confirmed the data reported by Franzman *et al.* (2004) who found that parents with high educational levels were brushing their children's teeth more frequently than those with lower educational levels. Similar findings were observed in the present study. The highest proportion of children who brushed twice a day was found among the children whose mothers had a post-graduate education. Nearly all the mothers in the present study reported that they breast-fed their children and about quarter of children were bottle-fed. Nearly thirty three percent of children were both breast and bottle-fed. It has been reported that there is an increase in the prevalence of dental caries in children who are bottle fed compared with breast fed children (Qadri *et al.* 2012).

However, there is still much controversy regarding the association between breast feeding and dental caries since the latter is associated with other factors such as sugar intake and microorganism (Seow, 1998).

Some researchers have reported that prolonged breast feeding of more than one year and high frequency of feeding were risk factors for ECC (Feldens *et al.* 2010; van *et al.* 2006; Sayegh *et al.* 2002). On the other hand, the results of the present study found that the mean dmft was slightly lower in children who were breast fed. This finding was similar to that of Sankeshwari *et al.* (2012) who observed that there was a noticeable decrease in the prevalence of ECC in children with breastfed up to two years and vice versa. Rugg-Gunn (2002) and Curzon and Drummond (1987) concurred that ECC was only associated with breastfeeding when it was 'prolonged' and when feeding was 'on demand'.

A primary risk factor for ECC is the use of bottle that contains fermentable carbohydrates (Slabšinskienė *et al.* 2010; Ribeiro and Ribeiro, 2004; Hallet and O'Rourke, 2003; Karjalainen *et al.* 2001; Milgrom *et al.* 2000; Holt *et al.* 1982). The present study found no association in dmft scores between children who were bottle-fed and children who were not. Furthermore, studies have reported that bottle feeding during day and night, frequency, duration of feeding and sweetened contents were also found to be significant risk factors for ECC (Hallet and O'Rourke, 2003; Karjalainen *et al.* 2001; Milgrom *et al.* 2000). In the present study, the highest dmft scores were found among children who were bottle-fed more than three times a day, however there was no association between dmft scores and the addition of sugar to the bottle content.

Nocturnal bottle feeding has been found to be a significant risk factor in the development of dental caries in primary teeth (Slabšinskienė *et al.* 2010; Lo *et al.* 2009; Schafer and Adair, 2000) and despite the fact that the present study found higher dmft scores among children who sleep with a bottle, this finding was not statistically significant.

Most dietary factors found to be significant are related to the consumption of sugar amount, frequency and time of consumption (Harris *et al.* 2004). Studies on preschool children concluded that daily consumption of sugar containing drinks especially during night, and daily sugar intake were independent risk factors for the development

of ECC (Karjalainen *et al.* 2001; Rodrigues and Sheiham, 2000). The present study found that more than 95% of the children were eating sugars every day, and nearly 30% ate sugar-containing food more than three times per day. Sweets and chocolates were most commonly eaten. Eating sugary food was significantly associated with higher dmft scores 2.8 compared with 0.50 for children who were not eating sugary food.

The dmft scores increased with the increasing frequency of eating sugar per day ranging from 1.34 for one time per day to 2.8 for three and more times per day. This finding is supported by Moynihan (2003) and Harris *et al.* (2004). Furthermore, the different types of sugar containing food (sweets and chocolates) were associated with high dmft scores. Begzati, Birisha and Meqa. (2010) found that the frequency of sweets consumption in 93% of children was 1-3 or more times per day and this frequency was significantly associated with ECC.

Dental visits play an important part in the preventive activities and education of children and are essential before the eruption of the first tooth of the child. During the first dental visit it is necessary to counsel the parent and educate them on how to prevent dental caries (Zafar *et al.* 2009). Dental visits were not common among the study children as less than ten percent of the children had visited a dentist and the most common reason for a visit was toothache. In a study carried out in Thailand, 12% of children reported visiting a dentist mainly on account of pain (Senesombath *et al.* 2010). Slightly higher dental attendances were reported in Hong Kong by Lo *et al.* (2009) and by Kopycka- Kedzierawski and Billings (2011).

CONCLUSIONS

In summary, the present study found that the prevalence of dental caries among preschool children was high and the mean dmft was directly proportional to the age. This was mostly associated with sugar consumption and therefore calls for preventive efforts to control sugar intake.

Tooth brushing is an important preventive practice for children and a high number of children were found to be brushing their teeth regularly and children who brushed more frequently had lower dmft levels.

There was no association between dental caries and bottle-feeding. Dental visits were uncommon among the study participants and efforts are needed to improve the utilization of dental services and determine what the barriers are to attendance.

The results of the present study indicate that there are areas for further investigation. There is urgent need for a comprehensive oral preventive programme for preschool children. This programme should be designed to include counselling of mothers/guardians and kindergarten school teachers for promoting healthy behaviours of children through dietary counselling, oral hygiene instructions including supervised tooth brushing at least twice daily.



References

- Aaltonen AS. (1991). The frequency of mother-infant salivary close contacts and maternal caries activity affect caries occurrence in 4-year-old children. *Proceedings of the Finnish Dental Society*; 87: 373-82.
- Acs G, Shulman R, Ng MW, Chussid S. (1999). The effect of dental rehabilitation on the body weight of children with early childhood caries. *Pediatr Dent*; 21(2):109-13.
- Alaluusua S, Malmivirta R. (1994). Early plaque accumulation- a sign for caries risk in young children. *Community Dentistry and Oral Epidemiology*; 22:273-76.
- Alaluusua S, Renkonen OV (1983). *Streptococcus mutans* establishment and dental caries experience in children from 2-4 years old. *Scandinavian Journal of Dental Research*; 91; 453-57.
- Alanen P, Hurskainen K, Isokangas P, Pietilä I, Levänen J, Saarni UM, Tiekso J. (1994). Clinician's ability to identify caries risk subjects. *Community Dentistry and Oral Epidemiology*; 22(2): 86-80.
- Alattonen AS, Tenovuo J. (1994). Association between mother-infant salivary contacts and caries resistance in children: a cohort study. *Pediatric Dentistry*; 16: 110-16.
- Al-Hosani E, Rug-Gunn A. (1998). Combination of low parental educational attainment and high parental income related to high caries experience in pre-school children in Abu Dhabi. *Community Dentistry and Oral Epidemiology*; 26: 31-36.
- Allukian M, Horowitz AM. (2003). Effective Community Prevention Programs for Oral Diseases. In: Jong's Community Dental Health. (5th ed).
- American Academy of Pediatric Dentistry (AAPD) (2003a). Policy on Early Childhood Caries (ECC): Classification, Consequences, and Preventive Strategies.
- American Academy of Pediatric Dentistry (AAPD) (2003b). Oral health risk assessment timing and establishment of the dental home, from: <http://www.aap.org/healthtopics/oralhealth.cfm>
- American Academy of Pediatric Dentistry (AAPD). (2005). Oral health risk assessment training for pediatricians and other child health professionals, from <http://www.aap.org/healthtopics/oralhealth.cfm>.
- American Dental Association (ADA). (2009). Statements on Early Childhood Caries. Available at <http://www.ada.org/prof/resources/positions/statements>.

American Dental Association(ADA). (2006). Council on Scientific Affairs. Professionally applied topical fluoride: evidence-based clinical recommendations. *Journal of American Dental Association*; 137(8):1151–9.

Ann M J, Trumbly S, Malik G. (2011). Early Childhood Caries: Determining the Risk Factors and Assessing the Prevention Strategies for Nursing Intervention. *Pediatric Nursing*; 37(1): 9-15.

Awooda EM, Saeed SM, Elbasir EI. (2013). Caries prevalence among 3-5 years old children in Khartoum State-Sudan. *Innovative Journal of Medical and Health Science*; 3: 42-44.

Ayhan H, Suskan E, Yildirim S. (1997). The effect of nursing or rampant caries on height, body weight and head circumference. *Journal of Clinical Pediatric Dentistry*; 20:209-12.

Azevedo TD, Bezerra AC, de Toledo OA. (2005). Feeding habits and severe early childhood caries in Brazilian preschool children. *Pediatric Dentistry*; 27: 28-33.

Beck, JD. (1998). Risk revisited. *Community Dentistry and Oral Epidemiology*; 26:220-25.

Begzati A, Berisha M, Meqa K. (2010). Early childhood caries in preschool children of Kosovo – a serious public health problem. *BMC Public Health*; 10:788: 1-8.

Beltran-Aguilar ED, Barker LK, Canto MT, Dye BA, Gooch BF, Griffin SO, Hyman J, Jaramillo F, Kingman A, Nowjack-Raymer R, Selwitz RH, Wu T. (2005). Surveillance for dental caries, dental sealants, tooth retention, edentulism, and enamel fluorosis-United States, 1988-1994 and 1999-2002. *Morbidity and Mortality Weekly Report*; 54(3): 1-43.

Berkowitz RJ. (1996). Etiology of nursing caries; a microbiologic perspective. *Journal of Public Health Dentistry*; 56(1): 51-54.

Berkowitz RJ. (2003). Causes, Treatment and Prevention of Early Childhood Caries. A Microbiologic Perspective. *Journal of the Canadian Dental Association*; 69(5):304-307.

Birkeland JM, Broch L, Jorkjend L. (1997). Caries experience as predictor for caries incidence. *Community Dentistry and Oral Epidemiology*; 4: 66.

Bourne LT. (2007). South African paediatric food-based dietary guidelines. *Maternal and Child Nutrition*; 3: 227-30.

Broadbent JM, Thomson WM, Williams SM. (2005). Does caries in primary teeth predict enamel defects in permanent teeth? A longitudinal study. *Journal of Dental Research*; 84:260-264.

Burt BA, Eklund SA. (1992). *Dentistry, Dental Practice and the Community*. W. B. Saunders, Philadelphia.

Burt BA, Eklund SA. (1999). *Dentistry, Dental Practice and the Community*. (5th Ed). Philadelphia.

Burt BA, Pai S. (2001). Sugar consumption and caries risk: a systematic review. *Journal of Dental Education*; 65:1017-23.

Carino K, Shinada K, Kawaguchi Y. (2003). Early childhood caries in northern Philippines. *Community Dentistry and Oral Epidemiology*; 31:81-9.

Centers for Disease Control and Prevention (CDC). (1999). Achievements in public health, 1900–1999: fluoridation of drinking water to prevent dental caries. *Morbidity and Mortality Weekly Report*; 48(41): 933–40.

Centers for Disease Control and Prevention (CDC). (2001). Recommendations for using fluoride to prevent and control dental caries in the United States. *Morbidity and Mortality Weekly Report Recommendation Report*; 50 (14): 1-30.

Child JS. (1996). Risks for and prevention of infective endocarditis. In *Cardiology Clinics-Diagnosis and Management of Infective Endocarditis*. Child JS. Philadelphia, Pa: WB Saunders Co; 14:327-343.

Cleaton-Jones P, Williams S, Fatti LP. (2000). Surveillance of primary dentition caries in Germiston, South Africa, 1981-1997. *Community Dentistry and Oral Epidemiology*; 26: 267-73.

Curzon ME, Drummond BK. (1987). Rampant caries in an infant related to prolonged on demand breastfeeding and lacto-vegetarian diet. *Journal of Pediatric Dentistry*; 3: 25-8.

Cvetkovi A, Vulovi M, Ivanovi M. (2006). Correlation between dental health status and environmental factors: nutrition, oral hygiene and saliva in children. *Serbian Dental Journal*; 35: 217-228.

Department of Health and Children, Ireland (DHC). (2002). Forum on Fluoridation. Stationary Office Dublin. www.fluoridationforum.ie.

Department of Health and Human Services. (DHHS). (2000). Oral health in America: A report of the Surgeon General. Washington, DC.

Department of Health and Human Services (DHHS). (2000). Oral health in America: A report of the Surgeon General. Washington, DC.

Dilley GJ, Dilley DH, Machen JB. (1980). Prolonged nursing habit: A profile of patients and their families. *Journal of Dentistry for Children*; 47:102-8.

Dini EL, Holt RD, Bedi R. (2000). Caries and its association with infant feeding and oral health-related behaviors in 3-4 year-old Brazilian children. *Community Dentistry and Oral Epidemiology*; 28: 241-48.

Douglass JM, Li Y, Tinanoff N. (2008). Literature review of the relationship between Mutans streptococci in adult caregivers and Mutans streptococci and dental caries in their children. *Journal of Pediatric Dentistry*; 30:375-87.

Drury TF, Horowitz AM, Ismail AI, Maertens MP, Rozier RG, Selwitz RH. (1999). Diagnosing and reporting early childhood caries for research purposes: a report of a workshop sponsored by the National Institute of Dental and Craniofacial Research, the Health Resources and Services Administration, and the Health Care Financing Administration. *Journal of Public Health Dentistry*; 59: 192–197.

Dye BA, Tan S, Smith V, Lewis BG, Barker LK, Thornton-Evans G, Eke PI, Beltrán-Aguilar ED, Horowitz AM, Li CH. (2007). Trends in oral health status: United States, 1988-1994 and 1999-2004. National Center for Health Statistics. *Vital Health Statistics*; 11(248): 1-92.

Eckersley AJ, Blinkhorn FA. (2001). Dental attendance and dental health behavior in children from deprived and non-deprived areas of Salford, Northwest England. *International Journal of Paediatric Dentistry*; 11: 103-9.

Edelstein B, Tinanoff N. (1989). Correlation of dental caries in young children with a non laboratory method of salivary *S. mutans* sampling. *Journal of Pediatric Dentistry*; 11: 129-32.

El-Nadeef MA, Hassab H, Al-Hosani E. (2010). National survey of the oral health of 5-year-old children in the United Arab Emirates. *Eastern Mediterranean Health Journal*; 16 (1):51-5.

Estupinan SR, Baez R, Horowitz H, Warpeha R, Sutherland B, Thamer M.. (2001). Salt fluoridation and dental caries in Jamaica. *Community Dentistry and Oral Epidemiology*; 29 (4): 247-52.

Fass EN (1962). Is bottle feeding of milk a factor in dental caries? *Journal of Dentistry Child*; 29:245-51.

Featherstone JD. (2000). The science and practice of caries prevention. *Journal of American Dental Association*; 131(7):887–99.

Feldens CA, Giugliani ERJ, Vigo A, Vitolo MR. (2010). Early Feeding Practices and Severe Early Childhood Caries in Four-Year-Old Children from Southern Brazil: A Birth Cohort Study. *Caries Research*; 44:445-52.

Firestone AR. (1982). Effects of increasing contact time of sucrose solution of powdered sucrose on plaque pH in vivo. *Journal of Dental Research*; 61:1243-4.

Franzman MR, Levy SM, Warren JJ, Broffitt B. (2004). Tooth-brushing and Dentifrice Use Among Children Ages 6-60 Months. *Pediatric Dentistry*; 26(1): 87-92.

Gallagher J, Rowe J. (2001). Community nurses' contribution to oral health. *British Journal of Community Nursing*; 6 (10): 526, 528-34.

Gibson S, Williams S. (1999). Dental caries in pre-school children: associations with social class, tooth brushing habit and consumption of sugars and sugar containing foods. *Caries Research*; 33: 101-13.

Gomes V, Fonseca A, Rodrigues M. (2001). Dental health: A challenge for the health professionals. *Review Brasil Enfermagem*; 54(1): 43-47.

Gray MM, Marchment MD, Anderson RJ. (1991). The relationship between caries experience in deciduous molars at 5 years and in the first permanent molars of the same child at 7 years. *Community Dental Health*; 8:3-7.

Greenwell AL, Johnsen D, DiSantis TA, Gerstenmaier J, Limbert N. (1990). Longitudinal evaluation of caries patterns from the primary to the mixed dentition. *Pediatric Dentistry*; 12: 278-82.

Gustafsson BE, Quensel CE, Lanke LS, Lundquist C *et al.* (1954). The Vipeholm dental caries study. The effect of different levels of carbohydrate intake on caries activity in 436 individuals observed for 5 years. *Acta Odontologica Scandinavica*; 11: 232-364.

Hallett K B, O'Rourke P K. (2003). Social and behavioural determinants of early childhood caries. *Australian Dental Journal*; 48(1):27-33.

Hallett KB, O'Rourke PK. (2006). Pattern and severity of early childhood caries. *Community Dentistry and Oral Epidemiology*; 34: 25-35.

Harris R, Nicoll A, Adair P, Pine C. (2004). Risk factors for dental caries in young children: a systematic review of the literature. *Community Dental Health*; 21(1):71-85.

Hashim R, Williams SM, Thomson WM. (2009). Diet and caries experience among preschool children in Ajman, United Arab Emirates. *European Journal of Oral Sciences*; 117: 734-40.

Health Education Authority. (1989). *The Scientific Bases of Dental Health Education*. (3rd ED). Health Education Authority, London.

Holt RD, Joles D, Winter GB. (1982). Caries in preschool children. *British Dental Journal*; 153: 107.

Ismail AI. (1998). Prevention of early childhood caries. *Community Dentistry and Oral Epidemiology*; 26 (1):49–61.

Jackson RJ, Newman HN, Smart GJ, Stokes E, Hogan JJ, Brown C, Seres J. (2005). The effects of a supervised tooth brushing programme on the caries increment of primary school children, initially aged 5–6 years. *Caries Research*; 39(2):108–15.

Jacobs TK. (2006). *The Prevalence of Early Childhood Caries in the Southern Cape Karoo Region*. University of the Western Cape.

Johnsen DC, Nowjack-Raymer R. (1989). Baby bottle tooth decay (BBTD): issues, assessment, and an opportunity for the nutritionist. *J Am Diet Assoc*; 89:112-6.

Johnston T, Messer LB. (1994). Nursing caries: Literature review and report of a case managed under local anaesthesia. *Australian Dental Journal*; 39:373-81.

Karjalainen S, Söderling E, Sewón L, Lapinleimu H, Simell O. (2001). A prospective study on sucrose consumption, visible plaque and caries in children from 3 to 6 years of age. *Community Dentistry and Oral Epidemiology*; 29(2):136-42.

Kaste LM, Gift HC. (1995). Inappropriate Infant Bottle Feeding. Status of the Healthy People 2000 Objective. *Archives of Pediatrics and Adolescent Medicine*; 149(7):786-91.

Kerosuo H, Honkala E. (1991). Caries experience in the primary dentition among groups of Tanzanian and Finnish 3-7-year-old children. *Community Dentistry and Oral Epidemiology*; 19: 272-76.

Kiwanuka SN, Åstrøm AN, Trovik TA. (2004). Dental caries experience and its relationship to social and behavioral factors among 3-5-year-old children in Uganda. *International Journal of Pediatric Dentistry*; 14: 336-46.

Klein H, Palmer JD. (1937). The dental caries experience of 5-12 and 14-year-old children in Great Britain. *Community Dental Health*; 11: 45-52.

Klein SP, Bohannon HM, Bell RM, Disney JA, Foch CB, Graves RC. (1985). The cost and effectiveness of school-based preventive dental care. *American Journal of Public Health*; 75(4):382–91.

- Kopycka-Kedzierawski DT, Billings RJ. (2011). Prevalence of dental caries and dental care utilization in preschool urban children enrolled in a comparative effectiveness study. *European Archives of Paediatric Dentistry*; 12(3): 133-8.
- Kumar J V, Moss M E. (2008). Fluorides in Dental public Health Programs. *The Dental Clinics of North America*; 52(2):387-401.
- Lee C, Rezaiaimira N, Jeffcott E, Domoto P, Weinstein P. (1994). Teaching parents at WIC clinics to examine their high caries-risk babies. *Journal of Dentistry for Children*; 61 (506): 347-349.
- Li Y, Caufield PW. (1995). The fidelity of initial acquisition of *mutans* streptococci by infants from their mothers. *J Dent Res*; 74: 681-85.
- Li Y, Navia JM, Bian JY. (1996). Caries experience in deciduous dentition of rural Chinese children 3 to 5 years old in relation to the presence or absence of enamel hypoplasia. *Caries Research*; 30:8-15.
- Li Y, Navia JM, Caufield PW. Colonization by *mutans* streptococci in the mouths of 3- and 4-year-old Chinese children with or without enamel hypoplasia. *Archives of Oral Biology* 1994; 39:1057-62.
- Litt MD, Reisine S, Tinanoff N. (1995). Multi-dimensional causal model of dental caries development in low income preschool children. *Public Health Report*; 110: 607-17.
- Lo E, Loo E, Lee C. (2009). Dental health status of Hong Kong preschool children. *Hong Kong Dental Journal*; 6 (1): 6-12.
- Loesche WJ. (1969). Role of *Streptococcus mutans* in human dental decay. *Microbial Review*; 50:353-80.
- Maciel SM, Marcenes W, Sheiham A. (2001). The relationship between sweetness preferences, levels of salivary *mutans* streptococci and caries experience in Brazilian pre-school children. *International Journal of Pediatric Dentistry*; 11: 123-30.
- Manson J. (2005). Trends in Dental Public Health. In: *Concepts in Dental Public Health*.
- Marinho VC, Higgins JP, Logan S, Sheiham A. (2002). Fluoride varnishes for preventing dental caries in children and adolescents. *Cochrane Database Systematic Review*; (3):CD002279.
- Marinho VCC, Higgins JPT, Sheiham A, Logan S. (2003) Fluoride toothpastes for preventing dental caries in children and adolescents. (Cochrane Review). In: *The Cochrane Library*. Chichester: John Wiley and Sons Ltd; (1):2-103.

Marinho VCC, Higgins JPT, Sheiham A, Logan S. (2009) Fluoride toothpastes for preventing dental caries in children and adolescents. (Issue 1). The Cochrane Library. Wiley.

Matilla ML, Rautava P, Sillanpaa M, Paunio P. (2000). Caries in five-year-old children and associations with family- related factors. Journal of Dental Research; 79: 876-81.

Milgrom P, Riedy CA, Weinstein P, Tanner AC, Manibusan L, Bruss J. (2000). Dental caries and its relationship to bacterial infection, hypoplasia, diet, and oral hygiene in 6-36 month-old children. Community Dentistry and Oral Epidemiology; 28(4):295-306.

Milnes A. (1996). Description and epidemiology of nursing caries. Journal of Public Health Dentistry; 56:38-49.

Mohan A, Morse D, O'Sullivan DM, Tinanoff N. (1998). The relationship between bottle usage/content, age, and number of teeth with salivary *mutans* streptococci levels in 6-24-month-old children. Community Dentistry and Oral Epidemiology; 26: 12-20.

Moynihan P. (2003). Diet and dental caries. In: Prevention of oral Disease (Murray JJ, Ed). Oxford University Press. Oxford.

Moynihan P. (2005). The role of diet and nutrition in the aetiology and prevention of oral diseases. Bulletin of the World Health Organization; 83 (9): 694-99.

Naylor MN, Murray JJ. (1983). The prevention of dental disease. Oxford.

Oulis CJ, Raad M, Martens L. (2000). Guidelines on the use of fluoride in children: an EAPD policy document. European Journal of Pediatric Dentistry; 1:7-12.

Paunio P, Rautava P, Sillanpaa M, Kaleva O. (1993). Dental Health Habits of 3-year-old Finnish children. Community Dentistry and Oral Epidemiology; 21: 4-7.

Peressini S, Leake JL, Mayhall JT, Maar M, Trudeau R. (2004). Prevalence of early childhood caries among First Nations children, District of Manitoulin, Ontario. Pediatric Dentistry; 14:101-10.

Peretz B, Eidelman E. (2003). New Insight into Early Childhood Caries and Strategies for Prevention. In: Jong's Community Dental Health. (Fifth ed).

Peter S. (2006). Epidemiology aetiology and prevention of dental caries. In: Essentials of preventive and Community Dentistry. (3rd Ed). New Delhi.

Picton DCA, Wiltshire PJ. (1970). A comparison of the effects of early feeding habits on the caries prevalence of deciduous teeth. *Dental Practice*; 20: 170.

Pieper K, Drerrler S, Heinzl-Gutenbrunner M, Neuhäuser A, Krecker M, Wunderlich K, Jablonski-Momeni A. (2012). The influence of social status on pre-school children's eating habits, caries experience and caries prevention behavior. *International Journal of Public Health*; 57(1): 207-15.

Pine CM, McGoldrick PM, Burnside G, Curnow MM, Chesters RK, Nicholson J, Huntington E. (2000). An intervention programme to establish regular toothbrushing: understanding parents' beliefs and motivating children. *International Dental Journal*; 50 (6) supplement 2: 312-23.

Qadri G, Nourallah A, Splieth CH. (2012). Early childhood caries and feeding practices in kindergarten children. *Quintessence International*; 43 (6): 503-10.

Raadal M, Elkidir EF, Rasmussen P. (1993). The prevalence of caries in groups of children aged 4-5 and 7-8 years in Khartoum, Sudan. *International Journal of Pediatric Dentistry*; 3 (1): 9-15.

Ramos-Gomez F, Huang G, Masouredis C, Braham RL. (1996). Prevention and treatment costs of infant caries in Northern California. *Journal of Dental Child*; 63(2): 108-12.

Ramos-Gomez FJ, Weintraub JA, Gansky SA, Hoover CI, Featherstone JD. (2002). Bacterial, behavioral and environmental factors associated with early childhood caries. *Journal of Clinical Pediatric Dentistry*; 26: 165-173.

Ramya K, Prasad KVV, Niveditha H. (2011). Public oral primary preventive measures: An Indian perspective. *Journal of International Oral Health*; 3(5):7-17.

Retnakumari N, Cyriac G. (2012). Childhood caries as influenced by maternal and child characteristics in pre-school children of Kerala-an epidemiological study. *Contemporary Clinical Dentistry*; 3 (1): 2-8.

Ribeiro NM, Ribeiro MA. (2004). Breastfeeding and early childhood caries: a critical review. *Journal of Pediatrics (Rio J)*; 80: 199-210.

Rodrigues CS, Sheiham A. (2000). The relationship between dietary guidelines, sugar intake and caries in primary teeth in low-income Brazilian 3-year-olds. A longitudinal study. *International Journal of Pediatric Dentistry*; 10: 47-55.

Roeters J, Burgersdijk R, Truin GJ, Van't Hof M. (1995). Dental caries and its determinants in 2-5-year-old children. *ASDC Journal of Dental Child*; 62: 401-8.

Rugg-Gunn A. (2002). Nutrition, dietary guidelines and food policy in oral health. In: *Community Oral Health*. (Pine, C. Ed). Wright.

Sanchez OM, Childers NK. (2000). Anticipatory guidance for infant oral health: Rationale and recommendations. *American Family Physician*; 61:115-20.

Sankeshwari RM, Ankola AV, Tangade PS, Hebbal MI. (2012). Feeding Habits and Oral Hygiene Practices as Determinants of Early Childhood Caries in 3- to – 5-year-old Children of Belgaum City, India. *Oral Health and Preventive Dentistry*; 10(3); 283-90.

Sayegh A, Dini EL, Holt RD, Bedi R. (2002). Caries prevalence and patterns and their relationship to social class, infant feeding and oral hygiene in 4–5-year-old children in Amman, Jordan. *Community Dental Health*; 19: 144–151.

Schafer TE, Adair SM. (2000). Prevention of dental disease. *Pediatric Clinics of North America*; 47: 1021-42.

Schwarz, Lo, Wong. (1998). Prevention of early childhood caries-results of a fluoride toothpaste demonstration trial on Chinese preschool children after three years. *Journal of Public Health Dentistry*; 58: 12-18.

Senesombath S, Nakornchai S, Banditsing P, Lexomboon D. (2010). Early Childhood Caries and Related Factors in Vientiane, LAO PDR. *Southeast Asian Journal Tropical Med Public Health*; 41 (3): 717-25.

Seow KW. (1998). Biological mechanisms of early childhood caries. *Community Dentistry and Oral Epidemiology*; 26 (1): 8-27.

Seow WK. (1991). Enamel hypoplasia in the primary dentition: A review. *Journal of Dental Child*; 58:441–452.

Simratvir M, Moghe GA, Thomas AM, Singh N, Chopra S. (2009). Evaluation of caries experience in 3-6 year-old children, and dental attitudes amongst the caregivers in Ludhiana city. *Journal of Indian Pedodontics and preventive dentistry*; 27(3): 164-68.

Sjogren P, Birkhed S, Rangmar T. (1995). Effects of a modified tooth paste technique on a proximal caries in preschool children. *Swedish Dental Journal*; 110: 1-10.

Skeie MS, Espelid I, Skaare AB, Gimmestad A. (2005). Caries patterns in an urban preschool population in Norway. *European Journal of Pediatric Dentistry*; 6 (1): 16-22.

Slabšinskienė E, Milčiuvienė S, Narbutaitė J, Vasiliauskienė I, Andruškevičienė V, Bendoraitienė EA, Saldūnaitė K. (2010). Severe early childhood caries and behavioral risk factors among 3-year-old children in Lithuania. *Medicina (Kuanas)*; 46(2): 135-41.

Sowole A, Sote E, Folayan M. (2007). Dental caries pattern and oral hygiene related factors in Nigerian preschool children. *European Archives of Pediatric Dentistry*; 8(4): 206-10.

Steiner M, Helfenstein U, Malthaler TM. (1992). Dental predictors of high caries increment in children. *Journal of Dental Research*; 71: 1926-33.

Stephen KW. (1993). Systemic fluorides: drops and tablets. *Caries Research*; 27(1): 9.

Subramaniam P, PrashanthP. (2012). Prevalence of early childhood caries in 8- 48 month old preschool children of Bangalore city, South India. *Contemporary Clinical Dentistry*; 3(1): 15-21.

Sufia S, Chaudhry S, Izhar F, Syed A, Syed A, Mirza BA, Khan AA. (2011). Dental Caries Experience in Preschool Children – Is It Related to A child’s Place of Residence and Family Income? *Oral Health Preventive Dentistry*; 9 (4): 375-79.

Sufia S, Khan AA, Chaudhry S. (2009). Maternal Factors and Child’s Dental Health. *Journal of Oral Health Community Dentistry*; 3(3): 45-8.

Tanzer JM. (1992). Microbiology of dental caries. In: *Contemporary Oral Microbiology and Immunology*. (Slots J, Taubman MA ed).

Thitasomakul S, Thearomtree A, Piwat S , Chankanka O, Pithpornchaiyakul W, Teanpaisan R, Madyusoh S. (2006). A longitudinal study of early childhood caries in 9- to 18-month-old Thai infants. *Community Dentistry and Oral Epidemiology*; 34(6): 429-36.

Tinanoff N, Palmer CA. (2003). Dietary determinants of dental caries and dietary recommendations for preschool children. *Journal of Public Health Dentistry*; 60: 197-206.

Tinanoff N, Kanellis MJ, Vargas C. (2002). Current understanding of the epidemiology, mechanisms, and prevention of dental caries in preschool children. *Pediatric Dentistry*; 24(6):543-51.

Tinanoff N, Kaste LM, Corbin SB. (1998). Early childhood caries: a positive beginning. *Community Dentistry and Oral Epidemiology*; 26(s1): 117-19.

Tinanoff N, O’Sullivan DM. (1997). Early childhood caries: Overview and recent findings. *American Academy of Pediatric Dentistry*; 19:12-6.

Tinanoff N, Reisine, S. (2009). Update on Early childhood Caries since the Surgeon General’s Report. *Academic Pediatrics*; 9:396-403.

Traebert J, Guimaraes A, Durante EZ, Serratine AC. (2009). Low maternal schooling and severity of dental caries in Brazilian preschool children. *Oral Health Preventive Dentistry*; 7: 39-45.

Twetman S, Garcia-Godoy F, Goepferd SJ. (2000). Infant oral health. *Dental Clinics of North America*; 44:487-505.

Tyagi R. (2008). The prevalence of nursing caries in Davangere preschool children and its relationship with feeding practices and socioeconomic status of the family. *Journal of Indian Society of Pedodontics and Preventive Dentistry*: 153-57.

US Department of Health and Human Services. (USDHHS). (2000). *Oral health in America: A report of the Surgeon General*. Rockville (MD): National Institute of Health.

Vadiakas G. (2008). Case definition, Aetiology and Risk assessment of Early Childhood Caries (ECC): A revisited review. *European Archives of Paediatric Dentistry*; 9 (3):114-25.

Van Wyk PJ, Louw AJ, Du Plessis JB. (2004). Caries status and treatment needs in South Africa: Report of the 1999-2002 National Children's Oral health Survey. *South African Dental Journal*; 59(6): 238-42.

Vargas CM, Crall JJ, Schneider DA (1998). Socio-demographic distribution of pediatric dental caries. NHANES III, 1998-1994. *Journal of American Dental Association*; 129: 1229-41.

Vinckier F, Gizani S, Declerk D. (2001). Comprehensive dental care for children with rampant caries under general anaesthesia. *International Journal of Paediatric Dentistry*; 11: 25-32.

Whelton H, O'Mullane D. (2002). Public Health aspects of oral diseases and disorders. In: *Community Oral Health*. (Pine, C. Ed). Wright.

World Health Organization. (WHO). (1997). *Oral Health Surveys. Basic Methods*. 4th edition. Geneva.

World Health Organization. (WHO). (1998): *Complementary Feeding of Young Children in Developing Countries. A Review of Current Scientific Knowledge*. Geneva.

World Health Organization. (WHO). (2003). *Global strategy for infant and young child feeding*. Geneva.

World Health Organization. (WHO). (2014). *Draft guidelines on free sugars released for public consultation*.

<http://www.who.int/mediacentre/news/notes/2014/consultation-sugar-guideline/en/>.

Wyne AH, Adenubi JO, Shalan T, Khan N. (1995). Feeding and socioeconomic characteristics of nursing caries children in a Saudi population. *Journal of Paediatric Dentistry*; 17:451-454.

Zafar S, Harnekar SY, Siddiqi A. (2009). Early childhood caries: aetiology, clinical considerations, consequences and management. *International Dentistry SA*; 11(4): 24-36.



Appendix 1: Data capture sheet

Prevalence of dental caries and tooth brushing habits among preschool children in Khartoum State, Sudan

Modified WHO Oral Health Assessment Form (1997)

Form No	Data collector	Date							
<table border="1" style="display: inline-table; width: 60px; height: 20px;"> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> </table>				<table border="1" style="display: inline-table; width: 30px; height: 20px;"> <tr><td style="width: 30px; height: 20px;"></td></tr> </table>		<table border="1" style="display: inline-table; width: 60px; height: 20px;"> <tr><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td><td style="width: 20px; height: 20px;"></td></tr> </table>			

Personal information

School No

--	--	--

 School type: Public (1) Private (2)

Child name: Gender: Male (1) Female (2)

Age

--	--	--

Dentition status and treatment need

Dentition status	Treatment need
A = Sound	0 = None
B = Decayed	1 = Filling
C = Filled with decay	2 = Extraction
D = Filled no decay	3 = Pulp care
E = Missing as a result of caries	4 = Need for other care
F = Not recorded	

55	54	53	52	51	61	62	63	64	65	
										Dentition status
										Treatment need

85	84	83	82	81	71	72	73	74	75	
										Dentition status
										Treatment need

Appendix 2: Questionnaire

Questionnaire for Mothers: Prevalence of dental caries and tooth brushing habits among preschool children in Khartoum State, Sudan

Form No

--	--	--

Data collector

--

Date

--	--	--

Child name: -----

Please Tick(√) in the appropriate box

1) Age

- a) < 20

--
- b) 20 – 29

--
- c) 30 – 39

--
- d) ≥ 40

--

2) Level of education

- a) None

--
- b) Primary/basic education

--
- c) Intermediate school

--
- d) High school

--
- e) University

--
- f) Postgraduate

--

3) What causes tooth decay? (you can tick more than one box)

- a) Germs

--
- b) Too much sugar

--
- c) Bad oral hygiene

--
- d) Other (please explain):
- e) Don't know

--

4) Does your child brush his/her teeth?

- a) Yes

--
- b) No

--
- c) Sometimes

--

5) If yes or sometimes, what does the child use to brush their teeth?

- a) Toothbrush

--
- b) Miswak

--
- c) Finger

--
- d) Other (please explain):

6) If yes or sometimes, when do he/she brush? (you can tick more than one box)

- a) In the morning

--
- b) During the day

--
- c) At night

--
- d) After meals

--
- e) Other (please explain):

7) Do you help your child to brush his/her teeth?

- a) Yes
- b) No
- c) Sometimes

8) Does your child spit out the toothpaste after brushing?

- a) Yes
- b) No
- c) Don't know

9) Does your child use toothpaste that contains fluoride?

- a) Yes
- b) No
- c) Don't know

10) When did your child start brushing his/her teeth?

- a) Soon after the first tooth came out
- b) When they turned 1 year old
- c) When they turned 2 years old
- d) When they turned 3 years old
- e) Other (please explain):

11) Did you breast-feed your child?

- a) Yes
- b) No

12) Do/Did you bottle-feed your child?

- a) Yes
- b) No
- c) Sometimes

13) If yes or sometimes, what do/did you put in the bottle? (you can tick more than one box)


- a) Milk
- b) Juice
- c) Other (please explain):

14) If yes, how many times a day do/did you feed her/him the bottle?

- a) Once
- b) Twice
- c) Thrice
- d) More than three times

15) If yes, do/did you add sugar to the contents in the bottle?

- a) Yes
- b) No
- c) Sometimes



16) Does your child fall asleep with the bottle in his/her mouth?

- a) Yes
- b) No
- c) Sometimes

17) Does your child eat sugar-containing food?

- a) Yes
- b) No
- c) Sometimes

18) If yes or sometimes, what did he/she eat? (you can tick more than one box)

- a) Sweets
- b) Chocolates
- c) Biscuits and cookies
- d) Other (Please explain):

19) If yes or sometimes, How frequently does your child eat sugar-containing food per day?

- a) Once
- b) Twice
- c) Thrice
- d) More than three times

20) Have you ever taken your child to the dentist?

- a) Yes
- b) No

21) If yes, why did you take your child to the dentist?

- a) For a routine check up
- b) Toothache
- c) Other (Please explain):

22) If yes, what did the dentist do? (you can tick more than one box)

- a) Only an examination
- b) Pulled out the tooth/teeth
- c) Did a filling
- d) Other (Please explain):
- e) Don't know

THANK YOU FOR YOUR TIME!

Appendix 3: Informed consent for questionnaire

Informed consent

I am Dr.Sitana MI Elidrissi, working at Oral Health Directorate, Ministry of Health, Khartoum State. I am presently a Master's Degree student in Dental Public Health at University of the Western Cape, South Africa. I am carrying out research on the oral health of preschool children in Khartoum State.

I would like to ask you a few questions regarding your child's oral health, tooth brushing and eating habits. This will take about 15 minutes of your time. All information obtained will be treated with utmost confidentiality.

Your participation in this research is completely voluntary, you may decline to participate in the study or withdraw from the study at any time without giving any reason and this will have no adverse effects on you.

If you need any further information regarding the study, please do not hesitate to contact me (Tel: 0912874710).

Thanking you in anticipation

Dr.Sitana

I understand the information that has been provided to me and agree to participate in the study.

Name: **Signature:**

Date:

Name of Witness: **Signature:**

Appendix 3: Informed consent for information sheet


Information sheet

I am Dr.Sitana MI Elidrissi, working at Oral Health Directorate, Ministry of Health, Khartoum State. I am presently a Master’s Degree student in Dental Public Health at University of the Western Cape, South Africa. I am carrying out research into the oral health of preschool children in Khartoum State. This will be done by examining the teeth of children. This research is important and will be of great value and it will add to the existing knowledge of tooth decay in children.

The process will take about 10 – 15 minutes. There are no risks involved if your child participates in this study.

All information gathered as part of the study will be treated as strictly confidential. No one will have access to this information except the researcher. Neither your child name nor any thing that identifies him/her will be used in any reports of this study.

If you have any questions or queries regarding the study, please contact me (Tel 0912874710).



The information above has been explained to me and I agree to allow my child to participate in the research. I understand that at any time I may withdraw my child from this study without giving a reason and without it affecting my child’s normal care and education.

Name:

Signature:

Date:

Name of Witness:

Signature: