

2% decline in tree density per annum. Giraffe impacts in particular were greatest in the 3 – 5 m height class, as the mean annual growth rate of trees in the *Vachellia* woodlands was significantly lower in the 3 – 5 m class compared to the < 1 m class. Removal of giraffes from the area would cause a predicted 2% increase in tree density per annum (Birkett, 2002).

Bond and Loffell (2001) determined the impacts of extralimital giraffe within the Ithala Game Reserve, KwaZulu-Natal province, South Africa. Higher browse lines, less pods and pod bearing trees and more twig tip damage was found within the high giraffe density areas. Mortality between different *Vachellia* species, too, was evident. Regarding *Vachellia davyi*, all individuals within the high giraffe density areas were dead, whilst the live individuals were either inaccessible, or within the low giraffe density areas. This was the case for *S. caffra* and *V. karroo* as well. Mortality stood at >50% and 21.2% in the high density areas, compared to <5% and 2% in the low density areas for *S. caffra* and *V. karroo* respectively. This mortality was further illustrated by fence line contrasts. Most trees with foliage accessible to giraffes were found dead within the reserve, compared to live ones found across the fence (Bond and Loffell, 2001).

1.10. Gap in literature

As seen by the literature, numerous studies have been conducted on the diurnal and foraging behaviour of giraffes throughout their natural range. However, behavioural and diet information in areas where they are considered extralimital is lacking, especially in the Western Cape, South Africa. In the Western Cape, no behavioural studies have been conducted, with only two documented studies on the diet of extralimital giraffe in Mosaic Thicket vegetation in the southern Cape (Cornelius, 2010; Gordon *et al.*, 2016).

Within the Western Cape, particularly within the Little Karoo, viable populations of giraffes are continuously being introduced into game farms to improve tourist attractions and for recreation. Two decades ago, the amount of game farms only made up a small portion of the region, therefore the significant effect these individuals had on the environment was considered negligible. However, the number of game farms has increased considerably since then.

Extralimital species may have negative impacts on indigenous fauna and flora. However, in order to determine impacts, as well as to determine adaptability and to initiate long term monitoring, the ecology of these species needs to be fully understood. The general daily behaviour of an organism is dictated by its feeding behaviour, this being the primary driving force for survival (Dagg and Foster, 1976). Therefore, the understanding of an organism's

nutritional requirements, which, in turn, dictate the behaviour of an organism, is crucial for the formulation and implementation of effective management decisions. This ecological information will, therefore, serve as the baseline for long term monitoring of extralimital species, and for the initiation of impact assessments of extralimital species to assist in protecting natural ecosystems.

1.11. Thesis description

The general aim of this study was therefore to determine the diurnal activities and foraging behaviour of extralimital giraffe within the Little Karoo in the Western Cape. The aim was achieved through the following objectives:

1. To determine the diurnal activity budgets,
2. the diet composition and browsing levels, and
3. the availability of browse and estimated site specific browsing capacities of extralimital giraffe.

The remainder of the thesis is set out as follows:

Chapter 2: Description of study sites

This chapter describes the locality and climate of the Little Karoo, thereafter focussing on the location, history, geology, vegetation and fauna of the two study sites (Kareesbos Private Game Reserve and Tsumkwe Private Game Reserve). The study populations of each study site are given and further discussed in terms of size and length of time present.

Chapter 3: Behavioural ecology of introduced giraffe

Determination of the behaviour of giraffe within the two study sites. The activity budgets of giraffe of different sexes and for the seasons of summer, autumn, winter and spring are provided and explained.

Chapter 4: Diet composition and preference of introduced giraffe

Determination of the diet composition of giraffes within each study site. The diet of the giraffes between different sexes and seasons are compared. The level of feeding, too, is observed and compared.

Chapter 5: Determination of browsing capacities

Determination of the amount of browse available in each of the study sites, quantified by means of the Biomass Estimates of Canopy Volume (BECVOL) method. This attempts to provide an estimate on whether or not sufficient browse is available for giraffe, and whether the study sites will be able to sustain them in the long term. Site-specific browsing capacities for giraffe based on biomass estimates are also described within this chapter.

Chapter 6: General conclusions and management implications

This is the concluding chapter in which all important aspects of each of the main themes are highlighted and commented on. Management implications based on the main results are given, and suggestions for future research are provided.



CHAPTER 2 : GENERAL DESCRIPTION OF STUDY SITES

2.1. Location

Situated in the Western Cape of South Africa, the boundaries to which the Little Karoo extends have been debated for numerous years as these boundaries have been defined either geographically, biogeographically or hydrologically (Maitre *et al.*, 2009). However, the typical boundaries used to describe the Little Karoo are generally according to Vlok *et al.* (2005) based on the combination of municipal administrative and water catchment boundaries encompassing the vegetation of the area (Thompson *et al.*, 2005). It is defined as a broad east-west orientated valley, bordered by the coastal Langeberg and Outeniqua mountain ranges in the south and the inland Witteberg and Swartberg mountain ranges in the north, being geographically isolated from the Great Karoo (Thompson *et al.*, 2005; Maitre *et al.*, 2009; Mann, 2014) (Figure 2.1).




Figure 2.1: Map of the Little Karoo (Blue square illustrates the area in which the two study sites are positioned)

The study was conducted on two privately owned farms, namely Kareesbos Private Game Reserve (KPGR) and Tsumkwe Private Game Reserve (TPGR), within the Oudtshoorn area of the Little Karoo. With numerous game farms present within and around Oudtshoorn, sites were selected based on the presence of giraffe on privately owned land and the positioning of the farms within the Thicket biome.

2.2. Climate

2.2.1. Rainfall

Whilst the Witteberg and Swartberg mountain ranges in the north separate the Little Karoo from the arid and summer rainfall of the Greater Karoo, the Langeberg and Outeniqua mountain ranges in the south have a larger influence on the climatic conditions of the sub-region (Potts *et al.*, 2013). Not only do these south bordering mountain ranges separate the region from the

mesic and winter rainfall of the coastal region of the Greater Cape Floristic Region (GCFR), they block the winter rainfall frontal systems from the Atlantic and the moist rain-bearing air from the Indian Ocean (Potts *et al.*, 2013, Vlok *et al.*, 2005). The physical barriers, thus, allow the Little Karoo to have an aseasonal rainfall regime, experiencing rainfall throughout the year (Potts *et al.*, 2013).

Rainfall varies with altitude and latitude, with the Langeberg-Outeniqua mountain ranges receiving the highest amount of rainfall of up to 1650 mm per annum, compared to the 1000 mm received by the Witteberg-Swartberg mountain ranges and the 100-300 mm received by the low lying central valleys (Figure 3.2) (Maitre *et al.*, 2009). Differing pressure systems and orographic gradients distribute the seasonal rainfall experienced in the west and in the east. In the far west (Montagu), rainfall is winter dominated and more reliable due to less variability in rainfall events. Rainfall events in the west are brought about by cold fronts, together with low pressure systems and westerly winds. In contrast, in the east (Willomore), summer rainfall is experienced, dominated by moisture received from the east, and convective systems are less influenced by orographic gradients. Areas centrally positioned (Ladismith) between the far west and east borders experience bimodal rainfall (Maitre *et al.*, 2009).




Figure 2.2: Average rainfall (mm), minimum and maximum temperatures (°C) in the Oudtshoorn area throughout the study period (2014-2015)

2.2.2. Temperature

The mean daily temperatures in the Little Karoo vary tremendously, largely due to high temperatures, low relative humidity, and little or no cloud cover (Desmet and Cowling, 1999). The highest temperatures are experienced between the months of December and February (summer), with the lowest temperatures occurring between the months of June and August (winter) (Nongwe, 2008). Mean daily maximum and minimum temperatures vary between lowland, mountain and coastal areas, with mountain and coastal areas experiencing similar temperatures, while differing to those experienced in the lowlands (Maitre *et al.*, 2009). In summer months, the lowlands experience mean daily maximum temperatures greater than 30°C and mean daily minimum temperatures ranging between 14 and 16°C. In winter, lowland areas experience mean daily maxima and minima temperatures ranging between 20 and 22°C and 4 and 8°C respectively (Figure 2.2). In summer, mountain and coastal areas experience mean daily maximum temperatures in the upper 20s, and mean daily minimum temperatures ranging between 10 and 12°C. In winter, the temperatures drop dramatically, with mean daily maxima and minima ranging between 10 and 12°C and -2 and 2°C respectively (Maitre *et al.*, 2009).

2.3. Vegetation

The aseasonal rainfall regime and geographical positioning of the Little Karoo has resulted in astonishingly high levels of plant diversity and endemism despite its aridity (Cupido, 2005; Nongwe, 2008). Four of the seven vegetation biomes within southern Africa are present within its boundaries, namely the Afromontane Forest, Fynbos, Succulent Karoo and Subtropical Thicket biome (Cupido, 2005; Vlok *et al.*, 2005; Nongwe, 2008). Of these four biomes, the Succulent Karoo and the Fynbos contribute considerably to the rich diversity and are recognised as two of the 25 global biodiversity hotspots (Maitre *et al.*, 2009). Vegetation present within the study farms is described in greater detail in sections 2.4.3 and 2.5.3.

2.4. KAREESBOS PRIVATE GAME RESERVE (KPGR)

2.4.1. Location and description

KPGR (33°32'34"S; 22°01'58"E) is approximately 900 ha in size and is separated from farmlands by means of fences in the north, south and west. The entrance is located on the northeast end of the farm, bordered by an unnamed gravel road running parallel to the perennial Wynands River. The farm is used for private recreational use, with occasion hunts occurring throughout the year. Minimal human interaction is made with animal species present, with the provision of water being in the form of one large manmade dam and several smaller water

points present throughout the area, as well as supplementary feed only being provided during times of extreme dry periods, which is not often.

Several antelope species are present, including plains zebra (*Equus quagga*), eland (*Taurotragus oryx*), greater kudu (*Tragelaphus stevensoni*), gemsbok (*Oryx gazella*), common duiker (*Sylvicapra grimmia*) and impala (*Aepyceros melampus*). A total of 10 giraffes were present at the commencement of the study, with the unknown removal of one adult male during mid-2014 and a birth occurring at the end of 2014. Evidence shows the occasional occurrence of the predatory black backed jackal (*Canis mesomelas*) and leopard (*Panthera pardus*).

2.4.2. Soil and geology

The dominant geological formations on KPGR are calcrete, hardpan, conglomerate, sandstone, siltstone and mudstone, all typical of the Enon Formation of the Uitenhage Group (ENPAT, 1995). The soils are red-brown in colour (ENPAT, 1995), characteristic of the dominant coarse-grained conglomerate, with size varying between pebbles, cobbles and boulders formed with other hard rocks (Lubke and De Moor, 1998). Soils are red-yellow apedal (structureless) and approximately < 300 mm deep, allowing water to drain freely (ENPAT, 1995). However, these characteristics make this kind of geology an ineffective/poor aquifer, offering poor borehole yields with water being of a brackish and salty nature (Lubke and De Moor, 1998).

2.4.3. Vegetation

Vegetation is described at different scales according to Mucina and Rutherford (2006) and Vlok *et al.* (2005). Vegetation units are described at a broad scale of 1:1 000 000 according to Mucina and Rutherford (2006), and thereafter described in greater detail at a scale of 1:50 000, making reference to the six-tier vegetation classification system developed by Vlok *et al.* (2005).

According to Vlok *et al.* (2005), three vegetation units are present in KPGR (Table 2.1) (Figure 2.3).

Table 2.1: Vegetation units present on KPGR

	Vegetation unit		
	Oudtshoorn	Grootkop Arid	
Category	Scholtzbosveld	Spekboomveld	Volmoed Gannaveld
Biome	Succulent Karoo	Thicket	Succulent Karoo

Arid + Spekboom +			
Habitat	Scholtzbosveld	Succulent Karoo	Gannaveld
Region	Gamka	Gamka	Gamka
	Oudtshoorn		
Sector	Scholtzbosveld	Grootkop	Volmoed
Unit	Solid	Mosaic	Solid

2.4.3.1. Scholtzbosveld

Scholtzbosveld is typical of flat areas, usually on the top of hills, dominated by a single shrub species, *Pteronia pallens*, accompanied by various other shrub and succulent plant species. This single species dominance is often said to have been due to heavy grazing, and although unpalatable, occupant species are palatable and favoured by many animal species.

In KPGR particularly, the habitat type Oudtshoorn Scholtzbosveld is present. Differing from other Scholtzbosveld units due to heuweltjies often being present, prominent species include *Mesembryanthemum crystallinum*, *Lycium cinereum*, *Oncosiphon suffruticosum*, *Phyllobolus splendens* and *Psilocaulon junceum*.

2.4.3.2. Gannaveld

Located on valley bottoms, Gannaveld is often described as open plains devoid of trees, with the presence of succulent and grass plant species being uncommon. Numerous tall and small shrubs are present within this vegetation type, with the appearance of an abundance of endemic annual species after rains.

Volmoed Gannaveld is present on KPGR. Shrub species are usually short (< 1 m) and infrequent, with the abundance of shrub species differing between the higher and lower lying areas. Fewer shrub species are found in the higher lying areas with *Eriocephalus spp.*, *Euryops subcarnosus*, *Justicia orchiodes*, *Pteronia pallens*, *Tripteris sinuate* and *Zygophyllum retrofractum* being the dominant species present. The lower lying areas are more established in terms of shrubs, with *Atriplex vestita*, *Salsola spp.*, *Suaeda fruticose* and *Zygophyllum microcarpum* being the dominant species present. Although the presence of grasses and succulents are uncommon, typical of Gannaveld vegetation, *Cenchrus ciliaris* and *Fingerhuthia africana* and *Malephora spp.* and *Phyllobolus splendens* respectively, can be found. If present, heuweltjies are dominated by the invasive *Augea capensis*.

2.4.3.3. Valley Thicket with Spekboom

Located on moderate to steep hill slopes, this vegetation type is characterised by the abundance of *Portulacaria afra*, forming fairly dense and closed stands or bush clumps (Vlok *et al.*, 2005). Other trees and shrubs that may be present do not form clumps, but rather occur as single plants.

Grootkop Arid Spekboomveld is present on KPGR. *Portulacaria afra*, the indicator species for this vegetation unit, together with other woody trees and tall shrubs, are locally dominant on north facing slopes, namely *Euclea undulata*, *Lycium cinereum*, *Grewia robusta*, *Nymannia capensis*, *Searsia undulata*, *Rhigozum obovatum*, *Carissa haematocarpa* and *Gloveria intergrifolia*. *Pappea capensis* and *Dodonaea augustifolia* are sparsely present (Vlok and Euston-Brown, 2002; Vlok *et al.*, 2005). Succulent Karoo vegetation patches are common, with *Pteronia spp.* and the succulent *Aloe*, *Cotyledon* and *Tylecolon spp.* species being abundant. The annual grass species *Aristida congesta* and *Enneapogan desvauzii* are occasionally present (Vlok and Euston-Brown, 2002; Vlok *et al.*, 2005).






Figure 2.3: Vegetation map of KPGR

2.5. TSUMKWE PRIVATE GAME RESERVE (TPGR)

2.5.1. Location and description

TPGR (33°33'41" S; 22°21'51" E) is approximately 1500 ha in size, bordered by the N12 in the south and by farmland fences in the west, east and northeast. Used for educational and tourism purposes, several animal species are present, including springbok (*Antidorcas marsupialis*), eland (*Taurotragus oryx*), greater kudu (*Tragelaphus strepsiceros*), gemsbok (*Oryx gazella*), wildebeest (*Connochaetes gnou*), impala (*Aepyceros melampus*) and giraffe (*Giraffa camelopardalis giraffa*). Water is supplied by several small waterholes throughout the farm and supplementary feed is provided for smaller antelope species, particularly springbok. Six giraffe were present at the start of the study period, but reduced to five at the end. One giraffe was killed due to an unknown medical condition at the beginning of 2014, one translocated to avoid inbreeding in August 2014, and one birth occurred in May 2015. Predator species present include the black backed jackal (*Canis mesomelas*) and leopard (*Panthera pardus*).

2.5.2. Soil and geology

Similar soil and geological formations are present on TPGR, as described in KPGR. See section 2.4.2.

2.5.3. Vegetation

According to Mucina and Rutherford (2006), TPGR encompasses vegetation that is positioned within the Albany Thicket, Fynbos and Succulent Karoo biomes. The Albany Thicket biome is the smallest compared to the Fynbos and Succulent Karoo biomes, making up only 2.2% of the area of South Africa, with the majority positioned within the Eastern Cape (Strydom and King, 2009). With approximately 1600 plant species, of which 20% are endemic to South Africa, overstocking has said to have severely degraded approximately 60% of the biome. TPGR is located in the Gamka Thicket (AT 2) vegetation unit (Mucina and Rutherford, 2006), occurring on the lower mountain slopes and river valleys between the Swartberg and Outeniqua mountain ranges bordering the Little Karoo centering around the Oudtshoorn area. Classified as least threatened, the main impact on this vegetation is fire, causing fragmentation.

Although the Fynbos biome only encompasses approximately 6.7% of South Africa, this biome possesses a higher number of plant species (approximately 7 500 vascular plant species) than any other biome occurring in the region, of which 70% are endemic (Mucina and Rutherford,

2006). Located in the Limestone Renosterveld Bioregion, particularly the Kango Limestone Renosterveld (FRI 1) vegetation unit (Mucina and Rutherford, 2006), it extends from Calitzdorp in the north to the Cango Caves region in the east, including the De Rust area (Mucina and Rutherford, 2006). Classified as least threatened, the main risk to this vegetation unit is cultivation, with approximately 14% already transformed (Mucina and Rutherford, 2006).

According to Vlok *et al.* (2005), three vegetation units are present in TPGR (Table 2.2) (Figure 2.4).

Table 2.2: Vegetation units present on TPGR

	Vegetation units			
	Mons Ruber	Kruisrivier	Grootkop	
Category	Waboom-Thicket	Spekboom-Pruimveld	Apronvel d	Olifants River and floodplain
Biome	Thicket	Thicket	Karoo	Drain
Habitat	Mosaic	Valley + Spekboom + Succulent Karoo	Gravel	River
Region	Gamka	Gamka	Gamka	Gamka
Sector	Mons Ruber	Kruisrivier	Grootkop	Olifants
Unit	Mosaic	Mosaic	Solid	Brack

2.5.3.1. Mons Ruber Waboom-Thicket

Located on moderate to steep hill slopes, this habitat type is characterised by the dominance of Albany Thicket elements on north facing slopes and in fire protected areas on south facing slopes where soils are deep and/or nutrient rich, and the dominance of Fynbos elements in shallow soils low in clay content on south facing slopes (Vlok and Euston-Brown, 2002, Vlok *et al.*, 2005). Albany Thicket elements include *Carissa haematocarpa*, *Euclea undulata*, *Gloveria integrifolia*, *Searsia undulata* and *Portulacaria afra* on the north facing slopes, as well as *Cassine euclaeformis*, *Gymnosporia buxifolia*, *Maytenus acuminata*, *Maytenus oleoides*, *Nymannia capensis*, *Olea europaea subsp. africana* and *Pterocelastrus tricuspidatus*

on the south facing slopes, whereas the Fynbos elements include *Agathosma recurvifolia*, *Erica karooica*, *Ishyrolepis gaudichaudiaris*, *Passerina obtusifolia*, *Protea nitida*, *Rhodocoma arida* and *Rhodocoma gracilis* (Vlok and Euston-Brown, 2002, Vlok *et al.*, 2005).

2.5.3.2. Grootkop Apronveld

Located in flat lying areas, these habitat types are patchy, however they have a large variety of species, with a dominance of asteraceous and succulent shrubs, aloes and geophytes, and with trees and tall shrubs being less prominent (Vlok *et al.*, 2005).

Dominant asteraceous shrubs include *Elytropappus rhinocerotis*, *Eriocephalus ericoides*, *Euryops lateriflorus*, *Euryops oligoglossus*, *Euryops subcarnosus*, *Hertia cf. pallens*, *Hirpicium alienatum*, *Pteronia incana*, *Pteronia pallens*, *Pteronia paniculata*, *Rosenia humilis* and *Tripteteris sinuata*. Succulent species include prominent leaf succulent shrubs, namely *Cotyledon orbiculata*, *Drosanthemum giffenii*, *Lampranthus haworthii*, *Ruschia cf. ceresiana* and *Tylecolon wallichii*, and smaller succulent species, namely *Adromischus triflorus*, *Conophytum truncatum* and *Glottiphyllum depressum*. Locally dominant grasses, *Digitaria argyrograpta* and *Fingerhuthia africana*, dominate these flat areas, whereas geophyte species, namely *Babiana sambucina*, *Freesia refracta*, *Gladiolus permeabilis*, *Lapeirousia pyramidalis*, *Moraea polyanthos*, *Moraea polystachya*, *Ornithogalum dubium*, *Ornithogalum juncifolium* and *Tritonia securigera*, dominate the patchy gravel areas. The less prominent tree and tall shrub species include *Carissa haematorcarpa*, *Euclea undulata*, *Searsia undulata*, *Lycium cinereum* and *Nymannia capensis*.

2.5.3.3. Kruisrivier Spekboom-Pruimveld

Located on flat lying areas and on the bottom of slopes, this vegetation unit is characterised by the dominance of *Portulacaria afra* and other woody trees and shrubs in a matrix of Succulent Karoo, on sandy and gravel soils.

Dominant woody trees and shrubs include *Carissa haematorcarpa*, *Euclea undulata*, *Gloveria integrifolia*, *Nymannia capensis*, *Pappea capensis*, *Grewia robusta*, *Searsia longispina*, *Rhigozhum obovatum*, *Pteronia incana*, *Eriocephalus africana*, *Eriocephalus ericoides* and *Eriocephalus punctulatus*. Stem and leaf succulent and geophyte species are also abundant in this vegetation unit and include *Aloe microstigma*, *Aloe ferox*, *Cotyledon orbiculare*, *Gasteria brachyphylla*, *Pelargonium peltatum*, *Pelargonium tetragonum*, *Quaqa pillansii*, *Ruschia ceresiana*, *Sarcostemma viminale*, *Tylecolon cacalioides*, *Tylecolon paniculata*, *Chlorophytum*

comosum, *Dipcadi viride*, *Drimia capensis*, *Hessea stellaris*, *Nerine humilis* and *Ornithogalum graminifolium*. Grass species are usually abundant after heavy rains, with species including *Cenchrus ciliaris*, *Erharta calycina*, *Panicum deustum*, *Sporobolus africanus* and *Stipa dregeana*.

2.5.3.4. Olifant's River and floodplain

Classified as a riverine unit, the indicator of this vegetation type is *Vachellia karroo*, with the distinctive prominence of *Salsola aphylla*. Herb species, such as *Ballota africana*, occur together with *Vachellia karroo*, with *Senegalia caffra*, *Chrysocoma oblongifolia* and *Cyperus congestus* occasionally present. These areas are highly subjected to soil erosion in the long term.






Figure 2.4: Vegetation map of TPGR

CHAPTER 3 : DIURNAL ACTIVITY BUDGETS OF EXTRALIMITAL GIRAFFE

3.1. INTRODUCTION

Understanding the social and non-feeding related behaviour of animals is of a great importance to understanding the way in which individuals not only interact with one another, but with the environment. For extralimital species within game farms, this knowledge is extremely beneficial, as animal behaviour may be altered and restricted in confined spaces, compared to open areas. In addition, it may help in providing information about the welfare of the populations, as well as the facilitation of managerial decisions regarding the maintenance of populations and habitats.

Giraffes have been able to successfully exploit different habitats through their physiological and behavioural adaptations (Fennessy, 2004). Physiologically, the nasal cooling of the brain enables giraffes to cope in arid environments (Fennessy, 2004; Dagg, 2014). On the other hand, giraffes are able to adjust their behaviour, such as the time spent on different activities (activity budgets), depending on various factors, such as the time of day, season, habitat type, foraging and social strategies (Dagg, 2014; Deacon, 2015). Activity budgets are discussed in greater detail in Chapter 1. However, in short, activity budgets have been documented by a few studies. Browsing is a dominant activity, comprising approximately 30-80% of giraffes' activity budgets (Pellew, 1983; Fennessy, 2004; Blomqvist and Renberg, 2007; Adolfsson, 2009; Deacon, 2015). Furthermore, activity budgets are shown to differ between sex and season (Pellew, 1983; Fennessy, 2004; Blomqvist and Renberg, 2007; Adolfsson, 2009; Deacon, 2015). For example, males are known to spend more time travelling, whereas females tend to spend more time foraging than males (Pellew, 1983; Fennessy, 2004; Blomqvist and Renberg, 2007; Adolfsson, 2009; Deacon, 2015). Seasonally, giraffes are shown to spend more time on energy conserving activities, such as resting, ruminating and vigilance during hotter seasons, and more time on energy consuming activities, such as walking and browsing during the cooler seasons (Fennessy, 2004; Deacon, 2015).

With only a few studies documenting the activity budgets of giraffes, information on the behaviour of extralimital giraffe in South Africa is limited, with no literature available on the behaviour of extralimital giraffe in Albany Thicket vegetated areas. Therefore, the results reported in this chapter are the first to be documented on extralimital giraffe within the Albany Thicket biome within South Africa.

The objectives and hypotheses of this chapter are as follows:

1. To describe the diurnal activity budgets of giraffes in Kareesbos Private Game Reserve (KPGR) and Tsumke Private Game Reserve (TPGR)
 - Hypothesis 1: Browsing will be the dominant activity displayed by giraffes
2. To compare the activity budgets of giraffes of differing sex
 - Hypothesis 2: Activity budgets will differ between males, females and juveniles
3. To compare activities of giraffes between different seasons
 - Hypothesis 3: Activity budgets will differ between the different seasons, with giraffes spending more time on energy consuming activities during cooler seasons and spending more time on energy conserving activities during the warmer seasons

3.2. METHODS

3.2.1. Field observations

Prior to the commencement of field observations, familiarization with individuals was done using photo identification (Appendix A and B). Field observations were conducted from a vehicle and occasionally on foot, at an approximate distance of 100-400 m where animals were observed to continue with 'natural' behaviour such as browsing. No absolute distance was decided upon due to the distance varying depending on the study area in terms of vegetation cover and height, the presence of other animals and environmental variables such as rain or fog. The herd was followed, either by foot or vehicle, with the number of individuals, estimated age and gender being noted. Observations, completed using binoculars if individuals were too far to observe with the naked eye, were conducted using a modified version of the interval scan sampling method (Altmann, 1974). The method involved observations being conducted on the same herd every five minutes from 06:00-18:00 on all visible individuals, with night observations being excluded from the study. This results in approximately 1 200 (maximum) observations per day depending on farm, herd size, and limiting factors such as climatic conditions and accessibility (terrain, slope, etc.). An observation was defined as one activity being conducted by one individual during one scan (Parker and Bernard, 2005; Deacon, 2015). To avoid observer bias and subjectivity, the first observed activity displayed by an individual

during a scan was recorded, with the next observation taken after five minutes. Observations were completed for each farm per season (winter, spring, summer and autumn).

Activity budgets were based on the following displayed behaviours (Fennessy, 2004; Deacon, 2015):

- Browsing: Defined as an individual foraging on a plant. Foraging to the point of chewing (not ruminating) and swallowing was included.
- Vigilance: Defined as an individual focussing on external stimuli such as predators or other animals. Individuals are seen as vigilant when eyes and ears are focused in the direction of the stimulus.
- Standing: Different to vigilance and can be defined as an individual resting in a standing position without doing any specific activity.
- Walking: Defined as an individual walking without foraging or ruminating.
- Rumination: Defined as an individual standing or walking and ruminating simultaneously. Ruminating should not be confused with chewing.
- Galloping: Defined as an individual running.

Activities observed at a lesser extent were:

- Defecation.
- Urination.
- Grooming: Defined as an individual licking itself or scratching against plants or other objects.
- Sexual behaviour: Includes flehmen response, sniffing or mounting displayed by males.
- Suckling: Whereby juvenile individuals nurse from any female.
- Osteophagia: Defined as an individual feeding on the bones of carcasses found within the study area.

3.2.2. Statistical analysis

The frequency of each activity was determined by expressing the number of each record for each activity as a percentage of the frequency of occurrence per season. All descriptive statistics and graphs calculated and displayed respectively were generated in Microsoft Excel 2015.

To evaluate seasonal differences in giraffe activities, proportion (percentage) data was used and checked for normality using the Shapiro-Wilk normality tests. The majority of the data was shown not to be normally distributed; hence, results were further tested using generalized linear mixed models (GLMM) to cater for non-normal distributions (IBM Corporation, 2012). The Poisson loglink model was applied for count data. The analysis was used to test for differences between behaviours displayed by giraffes of differing sex, as well as how behaviour differed between different seasons. Pairwise comparisons were tested at a 5% significance level.



3.3. RESULTS

3.3.1. Kareesbos Private Game Reserve

3.3.1.1. Activity budgets




Figure 3.1: Diurnal activity budgets of males (n = 3006) (top left), females (n = 2346) (top right) and juveniles (n = 725) (bottom) in Kareesbos Private Game Reserve during the entire study period

Hypothesis 1: Browsing will be the dominant activity displayed by giraffes

The overall activity budgets for males, females and juveniles in KPGR are shown in Figure 3.1. The most common activity displayed by males was browsing (37%), followed by walking (24%), rumination (20%) and vigilance (10%). However, resting (6%) accounted for a small portion of the activity budget.

Browsing (46%), too, was the most common activity displayed by females in KPGR. This was followed by walking (21%), rumination (16%) and vigilance (10%). Resting (4%) by females comprised a small portion of the diurnal activity budgets.

Juveniles, however, spent most of their time on vigilance (29%) throughout the study period. Walking (28%), browsing (19%) and rumination (12%) were observed to be the second, third and fourth most displayed activities, with resting (8%) occurring to a lesser extent.

3.3.1.2. Sex differences in activity budgets

Hypothesis 2: Activity budgets will differ between males, females and juveniles

Certain activities were shown to differ between sexes. Males (6% vs. 4%) were shown to rest ($F_{2,199} = 3.3346, p = 0.031$), walk (24% vs. 21%) ($F_{2,199} = 14.598, p < 0.001$) and ruminate (20% vs. 16%) ($F_{2,199} = 28.119, p < 0.001$) more than females. Males, too, ruminated (20% vs. 12%) ($p < 0.001$) and browsed (37% vs. 19%) ($F_{2,199} = 25.443, p < 0.001$) more than juveniles. Females, however, were shown to browse more than males (46% vs. 37%) ($p < 0.001$) and juveniles (46% vs. 19%) ($p < 0.001$), as well as ruminate (16% vs. 12%) more than juveniles ($p < 0.001$).

Although males and females spent relatively the same amount of time on vigilance ($p > 0.05$), juveniles were shown to be more vigilant than both males ($p < 0.001$) and females ($p < 0.001$). Juveniles, too, walked more than males ($p < 0.001$) and females ($p = 0.004$).

3.3.1.3. Seasonal differences in activity budgets

Hypothesis 3: Activity budgets will differ between the different seasons, with giraffes spending more time on energy consuming activities during cooler seasons and spending more time on energy conserving activities during the warmer seasons.




Figure 3.2: Behaviour of males (top left), females (top right) and juveniles (bottom) during winter, spring, summer and autumn in Karesbos Private Game Reserve

3.3.1.3.1. Energy conserving activities

Females rested ($F_{3,199} = 14.461$, $p < 0.001$) and ruminated ($F_{3,199} = 5.0384$, $p = 0.001$) more during the cooler seasons than the warmer seasons. Females rested less during summer than winter ($p < 0.001$), autumn ($p < 0.001$) and spring ($p = 0.002$). Rumination occurred the most during autumn, decreasing in winter ($p = 0.043$) and spring ($p = 0.022$), where rumination occurred the least. Females, however, were more vigilant ($F_{3,199} = 28.965$, $p < 0.001$) during the warmer seasons, with vigilance occurring less frequently during autumn than the remaining seasons ($p < 0.001$).

Males, too, rested ($F_{3,199} = 24.91$, $p < 0.001$) and ruminated ($F_{3,199} = 5.146$, $p = 0.002$) more during the cooler seasons than the warmer seasons. Males rested more during autumn than spring ($p < 0.001$) and summer ($p < 0.001$) and ruminated more during winter than spring ($p < 0.001$) and summer ($p = 0.017$). Males, however, were more vigilant during the warmer seasons ($F_{3,199} = 35.8442$, $p < 0.001$), with males being less vigilant in autumn than the remaining seasons ($p < 0.001$).

Juveniles rested ($F_{3,199} = 6.331$, $p < 0.001$) and ruminated ($F_{3,199} = 8.62$, $p < 0.001$) more during the cooler seasons than warmer seasons. More specifically, juveniles rested more during winter and autumn than during summer and spring ($p < 0.05$). In addition, juveniles ruminated more during winter than during spring ($p < 0.001$) and summer ($p = 0.001$) and ruminated the most in autumn ($p = 0.006$). Vigilance, however, occurred more during the warmer seasons ($F_{3,199} = 42.22$, $p < 0.001$), with juveniles being less vigilant in autumn than the remaining seasons ($p < 0.001$).

3.3.1.3.2. Energy consuming activities

No seasonal differences were shown for walking by females ($F_{3,199} = 2.521$, $p = 0.059$), males ($F_{3,199} = 0.432$, $p = 0.73$) and juveniles ($F_{3,199} = 0.83$, $p = 0.479$). However, females ($F_{3,199} = 4.335$, $p = 0.006$) browsed more during the cooler seasons, with more browsing occurring during winter and spring and autumn ($p < 0.001$). Males, however, browsed more during warmer seasons ($F_{3,199} = 3.211$, $p = 0.024$), with more browsing occurring during summer than during winter ($p = 0.025$) and autumn ($p = 0.003$). No seasonal variation in browsing was shown for juveniles ($F_{3,199} = 0.016$, $p = 0.997$).

3.3.2. Tsumkwe Private Game Reserve

3.3.2.1. Activity budgets




Figure 3.3: Diurnal activity budgets of males (n = 3101) (top left), females (n = 2090) (top right) and juveniles (n = 160) (bottom) in Tsumkwe Private Game Reserve during entire study period

Hypothesis 1: Browsing will be the dominant activity displayed by giraffes

The overall activity budgets for males, females and juveniles in TPGR are shown in Figure 3.3. The most common activity displayed by males was browsing (37%), followed by rumination (22%), walking (19%) and vigilance (11%). Resting accounted for 10% of the activity budget.

Browsing (47%), too, was the most common activity displayed by females in TPGR. This was followed by walking (18%), rumination (18%) and resting (9%). Vigilance (7%) by females comprised a small portion of the diurnal activity budget.

The activity budgets of juveniles were only determined during autumn. Vigilance was the most common activity (47%), followed by walking (26%), resting (16%) and browsing (10%). Juveniles were shown not to ruminate.

3.3.2.2. Sex differences in activity budgets

Hypothesis 2: Activity budgets will differ between males and females

Certain activities were shown to differ between sexes. Males were shown to walk more than females (19% vs. 18%) ($F_{1,127} = 11.174, p = 0.001$), with females spending more time browsing than males (47% vs. 37%) ($F_{1,127} = 78.314, p < 0.001$). Males and females spent relatively the same amount of time on resting ($F_{1,127} = 2.088, p = 0.151$), rumination ($F_{1,127} = 0.41, p = 0.523$) and vigilance ($F_{1,127} = 0.915, p = 0.341$) throughout the study period.

3.3.2.3. Seasonal differences in activity budgets

Hypothesis 3: Activity budgets will differ between the different seasons, with giraffes spending more time on energy consuming activities during cooler seasons and spending more time on energy conserving activities during the warmer seasons

3.3.2.3.1. Energy conserving activities

Females rested ($F_{3,127} = 2.615, p = 0.049$) and ruminated ($F_{3,127} = 9.421, p < 0.001$) more during the warmer seasons than during the cooler seasons. Females rested more during summer than during winter ($p = 0.01$), spring ($p = 0.008$) and autumn ($p = 0.014$). Rumination occurred more during summer than during winter ($p < 0.001$) and autumn ($p < 0.001$). No seasonal variation in vigilance was shown for females ($F_{3,127} = 2.233, p = 0.088$).

Males, too, rested ($F_{3,127} = 13.935, p < 0.001$) more during the warmer seasons compared to the cooler seasons. Males rested more during summer than during spring ($p < 0.001$) and autumn

($p < 0.001$). Males, however, ruminated ($F_{3,127} = 6.02$, $p = 0.001$) more during winter than during autumn ($p < 0.001$) and spring ($p = 0.042$) and more during summer than autumn ($p = 0.001$). Males were also shown to be more vigilant ($F_{3,127} = 5.986$, $p = 0.001$) during autumn than during winter ($p = 0.001$) and summer ($p = 0.002$).




Figure 3.4: Behaviour of males (top) and females (bottom) during winter, spring, summer and autumn in Tsumkwe Private Game Reserve

3.3.2.3.2. Energy consuming activities

No differences were shown in walking by females ($F_{3,127} = 9.421$, $p < 0.001$) and males ($F_{3,127} = 6.02$, $p = 0.001$). Browsing by females ($F_{3,127} = 14.685$, $p < 0.001$) decreased from the cooler to the warmer seasons, with females browsing more in winter than during summer ($p < 0.001$) and autumn ($p < 0.001$) and more during spring than summer ($p < 0.001$) and autumn ($p < 0.001$). Browsing by males ($F_{3,127} = 10.705$, $p < 0.001$), however, increased dramatically from

winter to spring ($p = 0.005$) and decreased from spring to autumn, with more browsing occurring in spring than in summer ($p = 0.001$) and autumn ($p < 0.001$).

3.4. DISCUSSION

Browsing will be the dominant activity displayed by giraffes

Activity budgets of giraffes have been documented by a few studies (Pellew, 1983; Fennessy, 2004; Blomqvist and Renberg, 2007; Adolfsson, 2009; Deacon, 2015). In all these documented cases, browsing was shown to be the dominant diurnal behaviour. In Mokolodi Nature Reserve, Botswana, giraffes spent approximately 36% of their time browsing (Blomqvist and Renberg, 2007). In Serengeti National Park, Tanzania (Pellew, 1983), females and males were shown to browse 72.4% and 55.34% of their time respectively. Similar dominance was shown in the Namib Desert, Namibia and Masai Mara National Reserve, with females and males spending 59% and 51% and 39% and 36% of their time browsing, respectively. As shown, the dominance of browsing is a norm for giraffes.

Activity budgets will differ between males and females

Cows spending more time browsing than males is considered as the norm in numerous studies conducted within natural ranges (Pellew, 1984a; Du Toit, 1990; Blomqvist, 2007; Fennessy, 2004; Deacon, 2015), as well as on other African herbivores (Spinage, 1968, Leuthold and Leuthold, 1978; Ben-Shahar and Fairall, 1987). According to Ginnett (1997), sexual dimorphism and reproductive concerns are two main factors influencing foraging differences in males and females. In terms of sexual dimorphism, Ginnett and Demment (1997) explain that total basal metabolism increases as a fractional exponent of body mass, therefore males, who tend to be larger than females, require more energy per unit time. In addition, a linear function exists between the gut capacity of herbivores and body mass. Herbivores with a greater body mass will have a higher gut volume to metabolic requirement ratio (“food processing capacity”), thus having the capability to retain food in the gut for longer periods of time, and in turn increasing the efficiency of nutrient extraction per unit of intake (Ginnett and Demment, 1997). Therefore, males have the ability to incorporate low quality forage within their diet, and meet metabolic and energy requirements in shorter time periods than females by increasing feeding rates (Ginnett and Demment, 1997). Reproductive requirements of either sex may also play a role in foraging differences (Parker, 2004).

Males were also shown to rest, walk and ruminate more than females. Males browsing less than females, thus allowing more time for other activities, may explain this. Fennessy (2004)

indicated that increased vigilance comes at a cost to foraging and reduces the risk of predation. However, predators are at no risk to giraffes in KPGR and TPGR, due to their absence. Therefore, increased vigilance by males may be socially related (Ginnett and Demment, 1997). For example, older males may be more vigilant in order to compete for females, whereas smaller males may be more vigilant to avoid aggression (Pellew, 1984; Artiss and Martin, 1995; Cameron and Du Toit, 2005). Furthermore, males tend to ruminate more than females due to their increased feeding rates and their need to extract more nutrients (Leuthold and Leuthold, 1978; Pellew, 1984).

Activity budgets will differ between the different seasons, with giraffes spending more time on energy consuming activities during cooler seasons and spending more time on energy conserving activities during the warmer seasons

Foraging time is dependent on numerous factors, such as body size, metabolic and energetic requirements of species and individuals, the availability of digestible food, food intake rates, habitat, as well as abiotic factors such as ambient temperatures and wind (Knight, 1991). The amount of digestible food plays a huge role in the proportion of time herbivores invest in browsing, with seasonal variations usually as a result of the change in food quantity and quality (Beckman and Prins, 1989; Knight, 1991). Seasonal variation in giraffe browsing has been frequently documented (Parker, 2004; Theron, 2005; Cornelius, 2010; Deacon, 2015). These variations have been associated with the change of the phenology of deciduous plant species that dominate these regions, with food quantity and quality decreasing during the drier seasons and foraging time increasing (Janecke and Smit, 2011). However, in this study, browsing by males increased from the cooler to the warmer seasons (winter to summer), whereas female browsing decreased from the cooler to the warmer seasons. This trend may be explained not by the quantity of available forage, but the quality. In both study sites, the majority of available forage species are evergreen, except for small patches of *Vachellia karroo* and *Lycium cinereum* (Chapter 3). However, since quality was not investigated within this study, it is suggested that differences in food quantity may be minimal between the different seasons, but changes in the quality of these plant species may cause the change in browsing frequency between seasons.

Females in TPGR were shown to rest more during the warmer seasons (summer) than the cooler seasons. This may be explained by ambient temperatures, with giraffes often resting more when ambient temperatures are high (Fennessy, 2004; Deacon, 2015). In addition, males ruminated

more during winter than the rest of the seasons in TPGR. This increase in rumination may be due to their browsing patterns and possible lower quality of browse during the winter seasons, thus compensating for their need to extract more nutrients.

3.5. CONCLUSION

Browsing was shown to be the dominant diurnal activity of males, females and juveniles in both study sites. Marked differences were observed in activity budgets of different sexes, with females browsing more than males, and males ruminating, walking and allocating more time to vigilance than females. Females likely browse more than males due to males requiring more energy per unit time, thus increasing feeding rates and incorporating lower quality forage in their diet to meet metabolic requirements in a shorter time. Males therefore ruminate more than females, due to this increase in feeding rates and low quality forage and the need to extract more nutrients. Furthermore, males spending less time browsing allows them to allocate more time to other activities, such as walking and vigilance, than females.

Season did not have as great an effect on all diurnal activities as within their natural ranges, possibly due to the differences in factors such as rainfall, temperature and other environmental factors. These factors need to be studied in greater depth to understand why the behaviour of giraffe in different localities differs according to season. Browsing is of great importance, particularly due to the extralimital status of giraffe within the area, therefore the identification and understanding of forage species, browsing behaviour and forage availability is essential and will be explored in Chapter 4 and Chapter 5 respectively.

CHAPTER 4 : DIET AND BROWSING LEVELS OF EXTRALIMITAL GIRAFFE

4.1. INTRODUCTION

The relationship between herbivores and plant communities is best described by a herbivore's foraging behaviour (Cornelius, 2010), and is essential for and influenced by, various reasons and factors respectively. Foraging behaviour is influenced by both plant and animal factors, such as plant availability, chemical composition and defence, and animal digestibility and body size (Owen-Smith, 1982; Pellew, 1984a; Bryant *et al.*, 1991; Cornelius, 2010). These factors result in herbivores continuously modifying their foraging behaviour according to seasonal variation in plant species availability and quality.

Giraffes are considered megaherbivores, with an adult female having to consume approximately 2.1% of their live weight per day, which approximates to 16 kg of digestible dry matter (DM⁻¹) (Dagg and Foster, 1976; Deacon, 2015). Regarded as browsers, giraffes feed on the leaves and shoots of trees and shrubs (Hofmann, 1989; Du Toit, 1988, Parker *et al.*, 2003). *Vachellia* species dominate the diet of giraffes within their natural ranges, resulting in an observed dietary shift from a deciduous dominated diet within the wet season, to a semi-deciduous, evergreen diet during the dry season (Leuthold and Leuthold, 1972; van Aarde and Skinner, 1975; Sauer *et al.*, 1977; Parker, 2004; Theron, 2005). Preferences, however, differ between areas, with plant species of higher water and protein content, as well as those with more newly produced shoots and leaves, being favoured (Pellew, 1984a; Hall-Martin and Basson, 1975; Cornelius, 2010).

It is said that the neck of the giraffe has developed in such a way that browsing occurs higher than other browsers, thus reducing the risk of competition (Cameron and Du Toit, 2007). Males are larger than females (Deacon, 2015), documented to browse higher in plant canopies than females (Parker, 2004). However, numerous studies (Leuthold and Leuthold, 1972; Sauer *et al.*, 1977; Woolnough and Du Toit, 2001; Blomqvist and Renberg, 2007; O'Connor, 2013) have observed giraffe to browse lower than three metres (see Chapter 2). This promotes niche overlap with other herbivores, hence increasing the risk of interspecific competition. For example, kudu and eland have been documented to have a maximum browsing height of 2.5 m (Dayton, 1978; Watson, 1999; Smit, 2004; Deacon, 2015).

Giraffe diet assessments, therefore, are crucial for management decisions regarding natural rangelands, providing the basis for understanding resource and habitat requirements of species

(Deacon, 2015), maintaining and regulating animal populations (Cornelius, 2010), identifying potential competition with other herbivore species, as well highlighting potential impacts giraffes may have on the environment (Parker, 2004; Pienaar, 2013; Deacon, 2015). Only two studies has been conducted on the diet composition of giraffes within the Western Cape (Cornelius, 2010; Gordon *et al.*, 2016), with no detailed studies being conducted on browsing levels within the area. Therefore, it is imperative to understand these ecological aspects of extralimital giraffe within the Western Cape, and to manage them accordingly to prevent negative impacts on thicket vegetation, and to ensure the welfare of giraffe populations.

Therefore, the aim of this chapter was to determine the diet composition and browsing postures of extralimital giraffe within the Western Cape. This chapter addresses the following objectives and hypotheses:

- To determine diet composition of giraffes
 - Hypothesis 1: Giraffes of differing sex will consume the same species
- To compare diet composition between different seasons
 - Hypothesis 2: Diet composition will not differ between different seasons due to the majority of the plant species being evergreen
- To determine the browsing levels of giraffes and compare between different seasons
 - Hypothesis 3: Giraffes will browse at low levels that will not differ seasonally

4.2. METHODS

4.2.1. Field observations

4.2.1.1. Diet observations

Giraffe observations were completed using the interval scan method, whereby observations were conducted every five minutes on each individual in a herd from sunrise to sunset for four days each season (winter, spring, summer and autumn) from August 2014 to May 2015. Observations, done by means of binoculars if individuals were too far to observe with the naked eye, were conducted mostly from a vehicle; however, observations were conducted occasionally on foot if obstacles such as vegetation cover or environmental factors such as mist or rain restricted views. When observed as browsing, sex of the individual, as well as the

browse species consumed and level of browsing was recorded. Before the commencement of the study, plant species were familiarised with.

4.2.1.2. Level of browsing

The general browsing level of giraffe was determined using methods by Theron (2005) and Deacon (2015). The level of browsing is generally categorised according to the height of browsing in relation to body posture, with six levels of feeding existing for giraffe. These levels of browsing range from ground height to head height, with each level and approximate heights illustrated in Figures 4.1 and 4.2 respectively, and are defined as follows:




Figure 4.1: Pictures depicting the different browsing levels by giraffes (Pictures taken by: Clement Cupido and Nico Smit)




Figure 4.2: Picture illustrating the approximate height measurements of an adult female giraffe as reported in literature (Patterson *et al.*, 1965; Hargens *et al.*, 1987; Mitchell and Skinner, 1993; Deacon, 2015)

- Level 1 (ground height) is defined as browsing with the head positioned lower than the knees, with knees bent or stretched outwards (height less than 1 m).
- Level 2 (knee height) is defined as browsing with the head positioned lower than the underside of the abdomen, with the exception of feeding at ground height with knees bent (level 1) (height approximately 1 m).
- Level 3 (chest height), is defined as browsing with the neck positioned horizontally or lower, with the neck not being positioned lower than the underside of the abdomen (height approximately 2 m).
- Level 4 (neck height) is defined as browsing with the neck positioned lower than 45° from the horizontal neck position, with the neck not extended lower than the base of the neck (height approximately 3 m).
- Level 5 (head height) is defined as browsing at the normal upright standing position, with the neck positioned at least 45° from the horizontal neck position (height approximately 3.5 m).

- Level 6 (stretch height) is defined as browsing with mouth and head reaching upwards higher than the base of the horns (height approximately 4 m and higher).

4.2.1.3. Statistical analysis

The diet composition and browsing levels of giraffe categories during the study period and different seasons were considered as the frequency of each browsed species represented as a percentage (%). The comparisons of diet composition and browsing levels between seasons and sex were further analysed using generalized linear mixed models (GLMM) to cater for non-normal distributions (IBM Corporation, 2014). The Poisson with loglink model was applied for count data. Pairwise comparisons were tested at a 5% significance level using the Tukey least significant difference (LSD) method. All tests were conducted at a 0.05 significance level using IBM SPSS Statistics (IBM Corporation, 2014).

4.3. RESULTS

4.3.1. Kareesbos Private Game Reserve (KPGR)

4.3.1.1. Diet composition

Hypothesis 1: Giraffes of differing sex will consume the same species

Table 4.1: The diet composition of all categories of the giraffe herd in KPGR throughout the study period

Species	Plant family	Proportion in diet (% \pm 1 SD)		
		Males	Females	Juveniles
<i>Pappia capensis</i>	Sapindaceae	44.95 \pm 5.88	47.71 \pm 8.59	54.40 \pm 12.01
<i>Portulacaria afra</i>	Portulacaceae	16.06 \pm 2.49	16.46 \pm 4.41	16.00 \pm 10.39
<i>Euclea undulata</i>	Ebenaceae	15.89 \pm 5.17	15.23 \pm 6.92	6.40 \pm 5.21
<i>Lycium spp.</i>	Solanaceae	12.58 \pm 6.61	11.88 \pm 8.42	15.20 \pm 10.02
Other				
<i>Salsola</i>				
<i>glabrescens</i>	Chenopodiaceae	2.73 \pm 3.19	1.58 \pm 0.79	1.60 \pm 2.77
<i>Searsia</i>				
<i>longispina</i>	Anacardiaceae	2.24 \pm 3.32	2.11 \pm 2.07	-

<i>Rhigozhum</i>				
<i>obovatum</i>	Bignoniaceae	1.57 ± 3.54	2.20 ± 4.21	1.60 ± 4.16
<i>Vachellia karroo</i>	Fabaceae	0.99 ± 1.37	0.70 ± 1.25	4.00 ± 6.25
<i>Leipoldtia</i>				
<i>schultzei</i>	Aizoaceae	0.75 ± 0.92	0.62 ± 0.69	-
<i>Augea capensis</i>	Zygophyllaceae	0.91 ± 1.68	0.35 ± 0.88	-
<i>Asparagus spp.</i>	Asparagaceae	0.58 ± 1.24	0.62 ± 0.57	0.80 ± 1.38
<i>Carissa</i>				
<i>haematocarpa</i>	Apocynaceae	0.17 ± 0.18	0.35 ± 0.38	-
<i>Euphorbia</i>				
<i>mauritanica</i>	Euphorbiaceae	0.17 ± 0.35	0.18 ± 0.32	-
<i>Schinus mollie</i>	Anacardiaceae	0.33 ± 0.45	-	-
<i>Malephora lutea</i>	Mesembryanthemaceae	0.08 ± 0.18	-	-

A total of 15 species from 14 plant families were consumed by giraffes in KPGR throughout the study period (Table 4.1). However, only four species, namely *P. capensis*, *P. afra*, *E. undulata* and *Lycium spp.* were considered important species, contributing approximately 89%, 91% and 92% to the diet of male, female and juvenile giraffes respectively.

Hypothesis 2: Diet composition will not differ between different seasons due to the majority of the plant species being evergreen

Pappia capensis was the most important species consumed by males, with significantly more being consumed in spring and summer than during winter and autumn ($F_{3,87} = 2.864$, $p = 0.041$). *Portulacaria afra* was the second most important species, with consumption remaining relatively the same between the different seasons ($F_{3,87} = 0.871$, $p = 0.459$). *Euclea undulata* was the third most important species consumed, with consumption decreasing in spring compared to the other seasons ($F_{3,87} = 5.853$, $p = 0.001$). Although the fourth most important species, the consumption of *Lycium spp.* only increased from spring to autumn ($F_{3,87} = 6.604$, $p < 0.001$). Consumption of the remaining species remained low during the study period. *Rhigozhum obovatum* and *V. karroo* were only consumed by males during spring and summer respectively. The consumption of *S. longispina* ($F_{3,87} = 3.463$, $p = 0.02$) and *S. glabrescens* ($F_{3,87} = 3.417$, $p = 0.021$) was greatest during winter, with less being consumed during the remaining seasons. *Carissa haematocarpa* (other category) was only consumed in spring and

summer, with *S. mollie* and *M. lutea* consumed in summer and spring respectively. *Euphorbia mauritanica* and *Asparagus spp.*, were only consumed in autumn, with *A. capensis* being only consumed in winter and autumn.

Pappea capensis was the most important species consumed by females, with consumption decreasing from winter to autumn ($F_{3,82} = 6.938, p < 0.001$). *Portulacaria afra* was the second most important species, with significantly less being consumed during spring than the remaining seasons ($F_{3,82} = 3.099, p = 0.031$). *Euclea undulata* was the third most important species consumed, with its importance decreasing in spring compared to the other seasons ($F_{3,82} = 5.145, p = 0.003$). The importance of *Lycium spp.*, the fourth most important species consumed by females, increased from spring to autumn ($F_{3,82} = 13.668, p < 0.001$). The importance of the remaining species remained low during study period, with *R. obovatum* only being consumed in autumn and spring. *Vachellia karroo* was only consumed during summer. No seasonal differences in the consumption of *S. glabrescens* was observed ($F_{3,82} = 0.728, p = 0.538$). *Searsia longispina* was not consumed during summer ($F_{3,82} = 2.834, p = 0.043$). Species in the “other” category in Figure 5.3, namely *A. capensis* and *E. mauritanica* were only consumed in autumn and winter respectively. *Carissa haematocarpa* was consumed only in summer and spring, whereas *L. schultzei* was only consumed in winter and summer.

Pappea capensis was the most important species consumed by juveniles, with juveniles consuming less during summer than during the remaining seasons ($F_{3,33} = 4.198, p = 0.023$). *Portulacaria afra* was the second most important species, with consumption decreasing from winter to autumn ($F_{3,33} = 4.421, p = 0.009$). *Lycium spp.*, the third most important species, was only consumed from spring, with consumption increasing till autumn ($F_{3,33} = 4.398, p = 0.01$). *Euclea undulata*, the fourth most important species, was consumed in winter. However, consumption of *E. undulata* ceased in spring, and increased in summer and autumn ($F_{3,33} = 2.909, p = 0.049$). Juveniles did not consume *S. longispina*. *Rhigozhum obovatum*, *V. karroo* and *S. glabrescens* were only consumed in spring, summer and autumn respectively.




Figure 4.3: Diet composition of male (top left), female (top right) and juvenile (bottom) giraffes in KPGR during the different seasons (% + 1 SE)

4.3.1.2. Browsing levels

Hypothesis 3: Giraffes will browse at low levels that will not differ seasonally




Figure 4.4: Browsing levels of males, females and juveniles in KPGR during the study period (% + 1 SE)

Males browsed more frequently at L3 (55%) (Figure 4.4) throughout the study period. Males decreased their browsing height levels from winter to autumn, with males browsing more at L2 ($F_{3,1316} = 7.843, p < 0.001$) than L3 ($F_{3,1316} = 4.925, p = 0.002$) during these seasons (Figure 4.5). In addition, males did not browse at L5 and L6 during winter (Figure 5.5).

Females browsed more frequently at L2 (41%) (Figure 4.4) throughout the study period. However, females increased their browsing height levels in winter, with females shown to browse significantly more at L3 ($F_{3,1286} = 5.057, p = 0.002$) during this season (Figure 4.5). Females, too, increased their browsing levels during spring, with females browsing more at L4 ($F_{3,1286} = 12.746, p < 0.001$) and L5 ($F_{3,1286} = 4.591, p = 0.003$). In addition, females only browsed at L6 during autumn (Figure 4.5).

Juveniles browsed more frequently at L3 and L4 (31.2%) (Figure 4.4) throughout the study period. Juveniles increased their browsing level during spring and summer, with juveniles shown to browse more at L4 ($F_{3,551} = 1.325, p = 0.026$) during these seasons (Figure 4.5). As the seasons moved into autumn, juveniles adjusted their browsing level, with juveniles browsing more at L2 ($F_{3,551} = 1.319, p = 0.049$) during this season (Figure 4.5). In addition, juveniles only browsed at L6 during autumn.




Figure 4.5: Browsing levels of male (top left), female (top right) and juvenile (bottom) giraffes in KPGR during the different seasons (% + 1 SE)

4.3.2. Tsumkwe Private Game Reserve (TPGR)

4.3.2.1. Diet composition

Hypothesis 1: Giraffes of differing sex will consume the same species

Table 4.2: The diet composition of all categories of the giraffe herd in TPGR throughout the study period

Species	Plant family	Proportion in diet (% \pm 1 SD)		
		Males	Females	Juveniles
<i>Searsia longispina</i>	Anacardiaceae	25.95 \pm 10.81	31.00 \pm 10.62	80.00
<i>Euclea undulata</i>	Ebenaceae	18.67 \pm 8.02	17.19 \pm 5.39	15.64
<i>Pappaea capensis</i>	Sapindaceae	8.94 \pm 3.46	12.06 \pm 4.11	4.36
<i>Vachellia karroo</i>	Fabaceae	18.51 \pm 16.34	8.44 \pm 8.41	-
<i>Grewia robusta</i>	Tiliaceae	8.09 \pm 6.12	10.62 \pm 7.53	-
<i>Lycium spp.</i>	Solanaceae	6.18 \pm 3.48	9.42 \pm 1.71	-
<i>Nymannia capensis</i>	Meliaceae	4.60 \pm 1.31	4.67 \pm 1.2	-
<i>Rhigozum obovatum</i>	Bignoniaceae	3.81 \pm 2.46	3.83 \pm 3.51	-
Other				
<i>Dodonaea viscosa</i>	Sapindaceae	2.11 \pm 0.84	1.30 \pm 0.88	-
<i>Euphorbia mauritanica</i>	Euphorbiaceae	1.39 \pm 2.02	1.21 \pm 1.05	-
<i>Salsola glabrescens</i>	Chenopodiaceae	1.20 \pm 1.76	1.89 \pm 1.09	-
<i>Zygophyllum spp.</i>	Zygophyllaceae	0.56 \pm 0.76	0.47 \pm 0.39	-

A total of 12 species from 12 plant families were consumed by giraffes in TPGR throughout the study period (Table 4.2). However, five species, namely *S. longispina*, *E. undulata*, *P. capensis*, *V. karroo* and *G. robusta* were considered important species, contributing approximately 80% and 79% to the diet of male and female giraffes respectively. Juveniles only browsed on three species, namely *S. longispina*, *E. undulata* and *P. capensis*.

Hypothesis 2: Diet composition will not differ between different seasons due to majority of the plant species being evergreen

Searsia longispina ($F_{3,41} = 14.301$; $p < 0.001$) and *E. undulata* ($F_{3,41} = 13.11$; $p < 0.001$) were the first and second most important species consumed by males respectively, with consumption of both species decreasing significantly from winter to autumn. *Pappea capensis* was the third most important species during winter, with its importance decreasing from spring to autumn ($F_{3,41} = 6.493$, $p = 0.001$) and the importance of *V. karroo* ($F_{3,41} = 68.705$, $p < 0.001$) and *G. robusta* ($F_{3,41} = 35.362$, $p < 0.001$) increasing during these seasons. *Lycium spp.* was the sixth most important species, with male consumption decreasing significantly from summer to autumn ($F_{3,41} = 18.652$, $p < 0.001$). The importance of *N. capensis*, *R. obovatum*, *D. viscosa*, *E. mauritanica*, *S. glabrescens* and *Zygophyllum spp.* remained low during all seasons. The consumption of *N. capensis* ($F_{3,41} = 9.88$, $p < 0.001$) differed seasonally, with males consuming *N. capensis* more during winter than the remaining seasons. The consumption of *D. viscosa* increased from winter to autumn ($F_{3,41} = 5.507$, $p = 0.003$). The consumption of *R. obovatum* increased significantly during the spring and summer seasons compared to autumn and winter ($F_{3,41} = 3.702$, $p = 0.019$). *Euphorbia mauritanica* ($F_{1,41} = 3.918$, $p = 0.028$) and *Zygophyllum spp.* were only consumed during summer and autumn, whereas *S. glabrescens* was only consumed during spring and autumn.

Similar results were observed for females, with *S. longispina* ($F_{3,41} = 30.605$; $p < 0.05$) and *E. undulata* ($F_{3,41} = 12.197$; $p < 0.001$) being the first and second most important species respectively, with consumption decreasing significantly from winter to autumn. *Pappea capensis* was the third most important species during winter, with its importance decreasing from spring to autumn ($F_{3,41} = 20.854$, $p < 0.001$) and the importance of *V. karroo* ($F_{3,41} = 16.015$, $p < 0.001$) and *G. robusta* ($F_{3,41} = 99.095$, $p < 0.001$) increasing during these seasons. *Lycium spp.* was the sixth most important species, with females' consumption increasing significantly from summer to autumn ($F_{3,41} = 3.025$, $p = 0.04$). The importance of *N. capensis*, *R. obovatum*, *D. viscosa*, *E. mauritanica*, *S. glabrescens* and *Zygophyllum spp.* remained low during all seasons. The consumption of *N. capensis* ($F_{3,41} = 1.528$, $p = 0.222$) did not differ seasonally. However, the consumption of *R. obovatum* increased significantly during the spring and summer seasons compared to autumn and winter ($F_{3,41} = 14.33$, $p < 0.001$). The consumption of *D. viscosa* ($F_{3,41} = 6.296$, $p = 0.001$) and *S. glabrescens* ($F_{2,41} = 8.244$; $p = 0.001$) increased during spring and autumn. *Euphorbia mauritanica* was only consumed in summer and autumn, with consumption increasing significantly in autumn. *Zygophyllum spp.* was only consumed in summer. Only one juvenile was present during the study period on TPGR during the season of autumn. However, since the individual was born in autumn, not

much browsing occurred. However, when browsing, the juvenile consumed only three species, namely *S. longispina*, *E. undulata* and *P. capensis* (Table 4.2).



UNIVERSITY *of the*
WESTERN CAPE





Figure 4.6: Diet composition of males (top) and females (bottom) in TPGR during the different seasons (% + 1 SE)

4.3.3.2. Browsing levels

Hypothesis 3: Giraffes will browse at low levels that will not differ seasonally




Figure 4.7: Browsing levels of males and females in TPGR during the study period (% + 1 SE)

Males browsed more frequently at L3 (55%) (Figure 5.7) throughout the study period. However, males adjusted their browsing level from winter to autumn, shown by the decrease in browsing at L3 ($F_{3,41} = 7.71, p < 0.001$) from winter to autumn, and the subsequent increase in browsing at L5 ($F_{3,41} = 6.16, p = 0.001$) from spring to autumn (Figure 4.8). Males were shown not to browse at L6 (Figure 4.7).

Females browsed more frequently at L3 (57%) (Figure 4.7) throughout the study period. However, females decreased their browsing levels during spring and autumn, with females browsing more at L2 ($F_{3,41} = 5.891, p = 0.002$) during these seasons (Figure 4.8). In addition, females were shown not to browse at L5 and L6 (Figure 4.7).





Figure 4.8: Browsing levels of males (top) and females (bottom) in TPGR during the different seasons (% + 1 SE)

4.4. DISCUSSION

Numerous studies have shown that giraffes tend to consume more than 20 species within their diet (Leuthold and Leuthold, 1972; Hall-Martin, 1974; van Aarde and Skinner, 1975; Sauer *et al.*, 1977, Sauer *et al.*, 1982, Parker, 2004). The only study conducted within Mosaic Thicket within the southern cape reported 20 species to have been consumed by giraffes (Cornelius, 2010). The wide variety of species consumed by giraffes in larger open areas is explained by the ability of giraffe to travel long distances and their high metabolic requirements (Parker, 2004). In addition, their high metabolic requirements allow less time for giraffes to be selective, thus including more plant species in their diet to compensate for their need to consume high quantities of food per day (Innis, 1958; Bell, 1971; Pellew, 1984a, b; Skinner and Smithers, 1990; Parker, 2004).

However, within the smaller, confined areas such as KPGR and TPGR, the number of species consumed was less than that reported in the literature, specifically 14 and 12 in KPGR and TPGR respectively. The vegetation types found within the two reserves may explain this. Discussed in greater detail in Chapter 2, the largest vegetation type taking up the majority of the area of KPGR is Grootkop Arid Spekboomveld (see Chapter 2), whereby the indicator species *P. afra* and a few other tall shrub and woody tree species, namely *E. undulata*, *L. cinereum*, *G. robusta*, *N. capensis*, *S. undulata*, *R. obovatum*, *C. haematocarpa*, *Gloveria integrifolia*, *P. capensis* and *Dodonaea augustifolia* are present (Chapter 2). In addition, approximately half of TPGR lies within the Kruisrivier Spekboom-Pruimveld vegetation type, a mixture of Mosaic Thicket and Fynbos elements (Vlok *et al.*, 2005) (Chapter 2). Dominant species present are few, namely *C. haematocarpa*, *E. undulata*, *G. integrifolia*, *N. capensis*, *P. capensis*, *G. robusta*, *S. longispina*, *R. obovatum*, *Pteronia incana*, *Ericcephalus africana*, *Eriocephalus ericoides* and *Eriocephalus punctulatus* (Chapter 2). Therefore, a single vegetation type occupying a large area may have resulted in this low diversity of species consumed compared to other studies due to a low diversity of species present within both reserves.

Although many species were present within the diet in both farms, only four and five species formed approximately 90% of the giraffes' diet in KPGR and TPGR respectively. Plants of the *Vachellia* genus are known to be principal diet species, and have been well documented within the literature (Innis, 1958; Oates, 1970; Leuthold and Leuthold, 1972; Hall-Martin, 1974; Stephens, 1975; van Aarde and Skinner, 1975; Field and Ross, 1976; Kok and Opperman, 1980; Parker, 2004; Cornelius, 2010). Cornelius (2010) showed giraffe to consume 60.5% of

V. karroo, which is much higher than that found within this study, even though being conducted within the same biome within the same region. Although less was consumed in both KPGR and TPGR, *V. karroo* was still consumed when present. *Vachellia karroo* is not a dominant species found within the vegetation types of both reserves (Chapter 2), which may explain the low proportion in the diet.

Seasonal differences in the importance of principal diet species were observed. In both reserves, the importance of *R. obovatum* and *Lycium spp.* in the case of KPGR, and the deciduous *V. karroo* and *G. robusta* in the case of TPGR, which all produce flowers after spring rains (Anderson, 2004) increased during spring and summer. Giraffes are known to adjust their diet during leaf bearing and flowering seasons, due to their preference for new shoots due to the higher water, protein, and mineral content (Fennessy, 2004), as well as their preference for flowers and pods (Sauer, 1983; Cooper *et al.*, 1988; Du Toit, 1990; Parker, 2004).

As seen in the results, the majority of browsing occurred at level 3 and below in both KPGR and TPGR. As seen in Figure 4.2, adults browsing at level 3, or chest height, occurs at an approximate height of 2 metres (Deacon, 2015). Therefore, younger adults or juveniles browsing at the same level will be browsing at a height lower than 2 m. Taking this into consideration, most of the browsing by giraffes on both reserves occurred below two metres. A possible reason for the low browsing can be due to the low height of vegetation present within the area (J Paulse, personal observation). This is lower than what has been documented in other studies (see Chapter 1), where browsing occurred mostly below three metres. Kudus and eland, present on both farms, have a maximum browsing height of 2.5 m (Dayton, 1978; Watson, 1999; Deacon, 2015). Therefore, with the continuous low levels of browsing by giraffe, as well as browsing by elands and kudus, niche overlaps occur with these three species, which are common (Leuthold and Leuthold, 1978; Deacon, 2015). This poses a risk of interspecific competition if there is a lack of browse available higher up in tree canopies.

4.5. CONCLUSION

Pappia capensis and *Searsia longispina* were the most important species forming the bulk of the giraffes' diet in KPGR and TPGR respectively. Although known as a highly preferred genus, *Vachellia* species formed a small proportion of giraffes' diet in KPGR due to its scarcity within the area. However, *V. karroo* was the third most important species consumed in TPGR, increasing in spring and summer due to its deciduous nature.

Giraffe browsing in both KPGR and TPGR occurred mostly below two metres, which may compete with other herbivores, such as kudu and eland present at both study sites. This may pose a risk to giraffe and other herbivore populations, as well as plant species, due to the possible increase in interspecific competition within the two reserves. It therefore has a management implication to keep numbers low and according to the browsing capacity to avoid losses of animals or the deterioration of vegetation.



CHAPTER 5 : QUANTIFICATION OF POTENTIAL BROWSE AND BROWSING CAPACITY

5.1. INTRODUCTION

Wildlife management requires proper management of both vegetation and soil (Bothma *et al.*, 2004). However, knowledge of various factors, such as the quantity and quality, growth processes and the grazing and browsing potential of plants, the causes and effects of overgrazing, fire and bush encroachment, soil dynamics, as well as seasonal variation, is required to achieve proper management (Bothma *et al.*, 2004). There are several measurable approaches to better understand these aspects and enforce proper management. Browsing capacity is one of these approaches, by managing vegetation through the regulation of large herbivore numbers.

It is important to distinguish between browse and available browse. The browse can be seen as the total plant material of the woody species that are potentially used as food by a specific group of animals (Rutherford, 1979). Seasonal changes such as the moisture content and palatability of species can also influence the definition of browse. The available browse is usually a more restricted quantity than browse. The availability of this browse is usually based on the maximum height above the ground which an animal can utilize. Some species like the giraffe and the black rhinoceros (*Diceros bicornis*) change their feeding height according to factors such as the time of year as well as social behaviour (Stuart-Hill, 1991).

Browsing capacity is a useful planning tool, providing guidelines as to how many large herbivores an area can sustain (Bothma *et al.*, 2004). It is defined as the productivity of the browsable portion of a homogeneous unit of vegetation expressed as the area of land required to maintain a single animal unit over an extended number of years without the deterioration of vegetation or soil (AU/ha) (Trollope and Trollope, 1990). Various factors may influence the browsing capacity of an area, such as the density, species composition, palatability, digestibility, growth potential and phenology of woody plants, as well as their chemical and structural defences (Bothma *et al.*, 2004; Cornelius, 2010; Janecke and Smit, 2011; Deacon, 2015). The combination of these factors, as well as the effect of predation and water availability, often leads to herbivores distributing themselves throughout the area, with certain areas being preferred and/or avoided (Deacon, 2015).

Although non-territorial, giraffes tend to travel within a specific home range, often overlapping with the home ranges of other giraffes and herbivores (Kok and Opperman, 1980). This,

together with giraffes having high metabolic requirements (Bell, 1971), may result in the overexploitation of resources if concentrated in certain areas, particularly within fenced areas (Deacon and Smit, 2012). The overexploitation of food resources, as well as the negative effects giraffes have on the ecosystem within their natural and extralimital ranges, is discussed in greater detail in Chapter 1.

Subtropical thicket is said to be resilient to indigenous herbivores (Stuart-Hill, 1992; Sigwela *et al.*, 2009). For example, in the Eastern Cape, South Africa, goats had a more detrimental effect on Kaffrarian Succulent Thicket, such as through the decline in plant species, even when stocked at similar stocking densities as elephants (Stuart-Hill, 1992). Elephants, on the other hand, kept the succulent thicket in a relatively uniform state (Stuart-Hill, 1992). However, it is not known how subtropical thicket is impacted by extralimital species. Due to this vegetation evolving in the absence of giraffes (Parker, 2004; Cornelius, 2010), Subtropical Thicket may be impacted by giraffe herbivory if not managed properly. In the Western Cape particularly, giraffes are being introduced for aesthetic and tourism reasons, despite their extralimital status, availability of preferred plant species, browsing capacities and giraffe impacts being unquantified. Considering these impacts, information on browsing capacities is imperative for the sustainable management of Thicket vegetation and giraffe populations alike. The following objectives and hypotheses for this study were thus as follows:

- Objective 1: To determine browsing capacity for each utilization area
 - Hypothesis 1: Browsing capacity will not differ between the different areas.
- Objective 2: To determine the browsing capacity for each height stratum
 - Hypothesis 2: Browsing capacity will decrease as the height stratum increases.
- Objective 3: To determine browsing capacity for the different seasons of the year
 - Hypothesis 3: Browsing capacity will remain the same between the different seasons.
- Objective 4: To determine the number of browsers that can be sustained by each farm

5.2. METHODS

5.2.1. Selection of habitats

Utilization areas were divided into four categories, namely no, low, medium and high utilization areas. The selection was based on the amount of time the giraffe herd spent within each area during the study period. GPS coordinates, with a coordinate taken once a giraffe was located and thereafter every time it moved to a different location, determined the amount of time spent within each area. GPS locations were taken for each of the four days per season, totalling 16 days. The vegetation type of each utilization area is described in Chapter 2.






Figure 5.1: Habitat preferences determined by giraffe movement on KPGR throughout the study period




Figure 5.2: Habitat preferences determined by giraffe movement on TPGR throughout the study period

5.2.2. Quantification of potential browse

Potential browse was quantified using the BECVOL procedure (Smit, 1989). BECVOL allows for the quantification of available plant biomass between the different height strata of woody plants. Evaluating the strata illustrates that the browsing capacity for browser game species is far more complex than calculating the grazing capacity for grazing game species. The following may influence the calculation of the browsing capacity for a specific browser species:

- Accessibility of the plant species to browsers;
- Height distribution of the browse material;
- Phenology of the plant species (whether they are evergreen or deciduous);
- Seasonal presence of flowers and pods or seeds with a high nutrient content.

Three 100 x 2.5 m (250 m²) transects were sampled within each utilization area within each farm, totalling 12 transects per farm (3 000 m²). For each transect, the dimensions of all plants, above 0.5 m, were measured. Plants with a height less than 0.5 m were regarded as saplings, and included in the number of plants per hectare (plants ha⁻¹). The dimensions measured for each woody plant included the following: (A) maximum tree height; (B) height of maximum canopy diameter; (C) height of first leaves/leaf bearing shoots; (D) maximum canopy diameter (considering a theoretical canopy being circular, maximum canopy diameter was taken as the average of two measurements (D1 and D2) taken perpendicular to each other); (E) base diameter at height of first leaves (application of same principle used for maximum canopy diameter).

Figure 5.3 is schematic diagram illustrating a typical woody plant and measurements in the BECVOL procedure. Some Thicket species do not follow this growth form, but the program differentiates between various shapes and sizes. In addition, it is important to note that the regressions (see below) incorporated into the BECVOL do not include all the species from the thicket biome. However, since there are no alternative methods to quantify potential browse, general regressions were used (Professor Nico Smit, University of the Free State, personal communication).




Figure 5.3: Schematic diagram of a typical tree and measurements involved in the BECVOL method (Smit, 2014)

The original BECVOL model was developed from harvested woody plants, which includes a regression equation that relates the spatial volume to the actual leaf volume and leaf dry mass (Smit, 1989; Smit, 1996). The BECVOL-3 model (Smit, 2014), the latest version, however, allows for the available browse to be calculated from the plant canopy volume. These estimates include those of leaves and young shoots (< 0.5 cm in diameter), an addition to the previous model (version 2) (Smit, 1996), which only allowed the estimates of the leaves (leaf volume and leaf dry mass) (Smit, 2014). The BECVOL-3 model, is therefore considered to be more accurate, as some browsers are selective feeders, browsing on leaves, as well as young shoot regrowth and fresh plant material (Penderis and Kirkman, 2014). More importantly, the BECVOL-3 model allows for the estimation of dry biomass in different height strata (Smit, 2014).

The BECVOL-3 model allows the following to be computed (described and detailed in Smit, 2014):

- Tree density (plants ha⁻¹)
- Evapotranspiration Tree Equivalents (ETTE/ha⁻¹) (An Evapotranspiration Tree Equivalent (ETTE) is defined as the leaf volume equivalent of a 1.5 m single-stemmed tree).

- Total leaf dry mass (DM ha⁻¹)
- Leaf biomass below a browsing height of 1.5 m (DM ha⁻¹)
- Leaf dry mass below a browsing height of 2.0 m (DM ha⁻¹)
- Leaf dry mass below a browsing height of 5.0 m (DM ha⁻¹)
- Shoot dry mass - shoots <0.5 cm below a browsing height of 1.5 m (DM ha⁻¹),
- Shoot dry mass - shoots <0.5 cm below a browsing height of 2.0 m (DM ha⁻¹),
- Shoot dry mass - shoots <0.5 cm below a browsing height of 5.0 m (DM ha⁻¹),
- Stem dry mass - stems >0.5-20 cm in diameter (DM ha⁻¹),
- Wood dry mass - wood >20 cm in diameter (DM ha⁻¹),
- Total wood dry mass (all fractions) (DM ha⁻¹),
- Total tree dry mass - leaves and wood combined (DM ha⁻¹).

Leaf and shoot DM ha⁻¹ was additionally calculated at a stratified height of below 1.5, 2 and 5 m. These represent the mean heights of impala (*Aepyceros melampus*) and Boer goat (*Capra aegagrus hircus*), kudu (*Tragelaphus strepciseros*) and giraffe respectively (Dayton, 1978, Skinner and Smithers, 1990).

5.2.3. Browsing capacity calculations

The average browsing capacity for the year for each utilization area, as well as the total area for each study site, was calculated using the following formula (Smit, 2014):

$$y = d \div \left[\frac{(DM1 \times f1 \times p1) + (DM2 \times f2 \times p2) + (DM3 \times f3 \times p3) + (DMx \times fx \times px)}{r} \right]$$

Where:

y = browsing capacity (ha BU⁻¹)

d = number of days in year (365)

DM = total leaf dry mass yield ha⁻¹

f = utilization factor

p = phenology

r = daily leaf dry mass required per BU (2.5% of body mass)

The substitution for specific species in terms of their daily leaf dry mass per grazer or browser unit (GU/BU) can be seen in Table 5.1 below (Smit, 2006). A browser unit is defined as the metabolic equivalent of a kudu (mean mass of 140 kg) (Smit, 2006). A giraffe is equivalent to 5.2 BU's.

Table 5.1: Approximate substitution values of a few game species in terms of grazer units (GU) and browser units (BU)

Game species	Aver. Mass (kg)	Intake (% of mass)	% grass	% leaves	GU	BU
Gemsbok	210	2.7	100	0	1.3	0
Black wildebeest	140	2.5	100	0	0.8	0
Burchell's zebra	216	4.1	100	0	1.9	0
Roan antelope	235	2.8	100	0	1.5	0
Springbok	37	3.0	70	30	0.1	0.1
Impala	52	2.7	70	30	0.2	0.1
Eland	460	2.4	30	70	0.7	2.2
Kudu	140	2.5	0	100	0	1.0
Giraffe	828	2.2	0	100	0	5.2

Phenology values were allocated for each species measured. These values vary from 100% (p = 1.0) for evergreen species, to 0% (p = 0.0) for deciduous species. Phenology values of 1, therefore, were given to all plant species in this study due to their evergreen nature, except for *Lycium cinereum* and *Vachellia karroo* (Table 5.2).

The utilization factor was applied, and refers to the edible parts of the leaf and shoots material to avoid the assumption that browsers feed in the centre of the foliage. The actual consumption

of this material, however, is limited due to the browsing utilizations of the animals (Deacon, 2015).

The number of giraffe individuals that each study area can maintain was therefore calculated as follows:

$$BU = \frac{ha}{BC}$$

Where:

BU = browser unit

ha = total area of study site

BC = browsing capacity (ha/BU)

5.2.4. Statistical analysis

Results from the BECVOL-3 model were further tested using generalized linear models (GLMM) to cater for non-normal distributions (IBM Corporation, 2012). The gamma with loglink model was applied for continuous data produced by the BECVOL-3 model. The analysis was used to test for differences between browsing capacities between utilization areas, height strata, as well as between seasons. Pairwise comparisons were tested at a 5% significance level using the Sequential Bonferroni method to control for alpha rates.

5.3. RESULTS

5.3.1. Species list

A total of 23 and 27 plant species were measured using the BECVOL method on Kareesbos Private Game Reserve (KPGR) and Tsumkwe Private Game Reserve (TPGR) respectively (Table 5.1). Of these species, 877 and 947 individuals were measured on KPGR and TPGR respectively. As observed in Chapter 5 (diet chapter), of the 23 species sampled in KPGR, only 14 were present within the giraffe diet. For TPGR, only 12 of the 27 plant species sampled were present within the diet. The majority of the plant species were evergreen, with only two plant species, namely *Lycium cinereum* and *Vachellia karroo* being deciduous (Table 5.2).

Table 5.2: List of all plant species present within the BECVOL sampling plots in KPGR and TPGR

Species	Family	Growth form	Phenology	Study site
<i>Galenia fruticosa</i>	Aizoaceae	Shrub	Evergreen	Both
<i>Tetragonia fruticosa</i>	Aizoaceae	Shrub	Evergreen	Kareesbos
<i>Searsia longispina</i>	Anacardiaceae	Tree	Evergreen	Tsumkwe
<i>Carissa haematocarpa</i>	Apocynaceae	Shrub	Evergreen	Both
<i>Asparagus spp.</i>	Asparagaceae	Shrub	Evergreen	Both
<i>Euryops spp.</i>	Asteraceae	Shrub	Evergreen	Both
<i>Pteronia pallens</i>	Asteraceae	Shrub	Evergreen	Both
<i>Pentzia incana</i>	Asteraceae	Shrub	Evergreen	Kareesbos
<i>Chrysocoma spp.</i>	Asteraceae	Shrub	Evergreen	Tsumkwe
<i>Elytropappus rhinocerotus</i>	Asteraceae	Shrub	Evergreen	Tsumkwe
<i>Eriocephalus africanus</i>	Asteraceae	Shrub	Evergreen	Tsumkwe
<i>Rhigozum obovatum</i>	Bignoniaceae	Tree	Evergreen	Kareesbos
<i>Salsola glabrescens</i>	Chenopodiaceae	Shrub	Evergreen	Kareesbos
<i>Cotyledon spp.</i>	Crassulaceae	Shrub	Evergreen	Both
<i>Crassula spp.</i>	Crassulaceae	Shrub	Evergreen	Kareesbos
<i>Tylecodon wallichii</i>	Crassulaceae	Shrub	Evergreen	Tsumkwe
<i>*Euclea undulata</i>	Ebenaceae	Tree	Evergreen	Both
<i>Euphorbia mauritanica</i>	Euphorbiaceae	Shrub	Evergreen	Both
<i>Aspalathus spp.</i>	Fabaceae	Shrub	Evergreen	Kareesbos
<i>Calobota sericea</i>	Fabaceae	Shrub	Evergreen	Tsumkwe
<i>Nymmania capensis</i>	Meliaceae	Tree	Evergreen	Tsumkwe
<i>Drosanthemum spp.</i>	Mesembryanthemaceae	Shrub	Evergreen	Both
<i>Ruschia robusta</i>	Mesembryanthemaceae	Shrub	Evergreen	Both
<i>Psilocaulon spp.</i>	Mesembryanthemaceae	Shrub	Evergreen	Kareesbos

	Mesembryan-			
<i>Aridaria spp.</i>	themaceae	Shrub	Evergreen	Tsumkwe
* <i>Portulacaria afra</i>	Portulacaceae	Shrub	Evergreen	Kareesbos
<i>Thamnochortus spp.</i>	Restionaceae	Restio	Evergreen	Tsumkwe
* <i>Pappea capensis</i>	Sapindaceae	Tree	Evergreen	Both
<i>Dodonaea viscosa</i>	Sapindaceae	Tree	Evergreen	Tsumkwe
* <i>Lycium cinereum</i>	Solanaceae	Shrub	Deciduous	Both
* <i>Lycium ferocissimum</i>	Solanaceae	Shrub	Evergreen	Both
<i>Hermannia amoena</i>	Sterculiaceae	Shrub	Evergreen	Kareesbos
<i>Struthiola ciliata</i>	Thymelaeaceae	Shrub	Evergreen	Tsumkwe
* <i>Grewia robusta</i>	Tiliaceae	Tree	Evergreen	Tsumkwe
<i>Zygophyllum spp.</i>	Zygophyllaceae	Shrub	Evergreen	Both

*indicates species forming the bulk of the diet composition of the two giraffe populations (see Chapter 4)

5.3.2. Kareesbos Private Game Reserve

5.3.2.1. Available browse



Table 5.3: Plant density (plants ha⁻¹), evapotranspiration tree equivalents per hectare (ETTE ha⁻¹) and total dry mass per hectare (kg DM ha⁻¹) for all species measured using the BECVOL method in the different utilization areas in KPGR

Utilization area	Plant density (Plants ha ⁻¹)	Evapotranspiration tree equivalents (ETTE ha ⁻¹)	Total browse mass (< 5 m) (kg DM ha ⁻¹)
No utilization	4 349	4 606	1 868
Low utilization	2 440	2 698	1 087
Medium utilization	2 401	3 456	1 417
High utilization	2 879	3 717	1 502

Table 5.4: Plant density (plants ha⁻¹), evapotranspiration tree equivalents per hectare (ETTE ha⁻¹) and total dry mass per hectare (kg DM ha⁻¹) for all important diet species measured using the BECVOL method in the different utilization areas in KPGR

Species	Plant density (Plants ha ⁻¹)				Evapotranspiration tree equivalents (ETTE ha ⁻¹)				Total browse mass (< 5 m) (kg DM ha ⁻¹)			
	Utilization area				Utilization area				Utilization area			
	No	Low	Moderate	High	No	Low	Moderate	High	No	Low	Moderate	High
<i>Pappea capensis</i>	0	93	40	133	0	487	348	667	0	201	146	274
<i>Portulacaria afra</i>	667	67	107	413	2 529	343	680	1 251	1 042	140	283	508
<i>Euclea undulata</i>	227	173	107	267	861	719	463	549	352	294	190	222
<i>Lycium cinereum</i>	347	267	160	67	157	269	242	36	61	107	98	14
<i>Lycium ferocissimum</i>	67	80	80	40	190	137	151	53	77	55	61	21
Total	1 308	680	494	920	3 737	1 955	1 884	2 556	1 532	797	778	1 039

The results of the biomass assessment of all plant species recorded in the KPGR survey plots are presented in Table 5.3. From this data, it is notable that plant density per hectare was the lowest in the low utilization area and highest in the no utilization area. However, important diet plant density per hectare was highest and lowest in the no and moderate utilization areas respectively (Table 5.4).

Evapotranspiration tree equivalents per hectare was the highest and lowest in the no and low utilization areas respectively (Table 5.3). Evapotranspiration tree equivalents of important diet species per hectare, however, was highest in the no utilization area, but lowest in the moderate utilization area (Table 5.4).

Total browse per hectare, i.e. total dry matter per hectare, was highest in the no utilization area, with the second highest and lowest dry matter per hectare occurring within the high and low utilization areas respectively (Table 5.3). Similar results were observed regarding the important diet species (Table 5.4).

Pappea capensis, was observed to be absent within the no utilization area, compared to the high utilization area where plant density per hectare, ETTE per hectare and total dry matter mass was highest in the high utilization area. *Portulacaria afra* had the highest plant density per hectare, ETTE per hectare and total dry matter mass within the no utilization area compared to the other areas (Table 5.4). The presence of *L. cinereum* per hectare, too, was highest in the no utilization areas; however, both *Lycium* species showed the lowest amounts of plants per hectare, ETTE per hectare and total dry matter mass in the high utilization area. The ETTE per hectare and total dry matter mass of *E. undulata* was highest in the no utilization area, but more individuals per hectare were found within the high utilization area.

5.3.2.2. Browsing capacity




Figure 5.4: Browsing capacity of the (top left) 1.5 m, (top right) < 2 m and (bottom) < 5 m height stratum within each utilization area in KPGR

Hypothesis 1: Browsing capacity will not differ between the different areas.

It is evident that browsing capacity decreased the more the area was utilised in all height strata, due to the number of hectares required to sustain a browser unit being greatest in the low utilization area for each height stratum (Figure 5.4), compared to the no ($p < 0.001$), medium ($p < 0.001$) and high ($p < 0.001$) utilization areas. This may be due to plant density (Table 5.3) and important diet plant density (Table 5.4) increasing from the low to the high utilization area, within which the number of important diet species was greatest.

However, the no utilization browsing capacity was lowest in all height strata compared to low ($p < 0.001$), medium ($p < 0.001$) and high ($p < 0.001$) utilization areas. Steeper slopes (personal observation), as well as the absence of important diet species, such as *P. capensis* (Table 5.3) in the no utilization area, may cause giraffes not to occupy these areas.

Hypothesis 2: Browsing capacity will decrease as the height stratum increases.

It is evident that less hectares are required per browser unit as the height stratum increases (Figure 5.4). More hectares are required per browser unit relying on browse in the less than 1.5 height stratum, compared to the less than 2 ($p < 0.001$) and 5 m ($p < 0.001$) height strata.

Hypothesis 3: Browsing capacity will remain the same between the different seasons.

Browsing capacity differed between seasons, with more hectares required per browser unit in KPGR during winter than during summer ($p < 0.001$) and autumn ($p < 0.001$). Browsing capacity was relatively similar, however, between winter and spring ($p = 0.295$), as well as between the remaining seasons ($p > 0.05$) (Figure 5.4).

5.3.2.3. Number of browser units sustained

The number of calculated browser units that can be sustained, as well as the current browsers currently found on KPGR can be observed in Table 5.6 and 5.7 respectively.

Table 5.5: Table representing the browsing capacity, browser and giraffe numbers that can be sustained in KPGR

	Browsing capacity (ha/BU)			Total browser units sustained			Number of giraffe sustained		
	< 1.5 m	< 2 m	< 5 m	< 1.5m	< 2m	< 5 m	< 1.5 m	< 2 m	< 5 m
January	6.4	4.8	4.0	141	188	225	27	36	43

February	6.4	4.8	4.0	141	188	225	27	36	43
March	6.4	4.8	4.0	141	188	225	27	36	43
April	6.4	4.8	4.0	141	188	225	27	36	43
May	6.5	4.9	4.1	138	184	220	27	35	42
June	6.6	4.9	4.1	136	184	220	26	35	42
July	6.7	5	4.2	134	180	214	26	35	41
August	6.8	5	4.2	132	180	214	25	35	41
September	6.8	5	4.2	132	180	214	25	35	41
October	6.5	4.9	4.1	138	184	220	27	35	42
November	6.4	4.8	4.0	141	188	225	27	36	43
December	6.4	4.8	4.0	141	188	225	27	36	43

As observed in Table 5.6, the following browser units that can be sustained by browse at different heights are as follows:

- 1.5 m: average of 138 BU
- < 2 m: average of 185 BU
- < 5 m: average of 221 BU
 - < 5 m = 221 BU = 43 giraffes, therefore browsing capacity = 900 ha/221 BU = 21 ha per giraffe (< 5 m)

Table 5.6: The number of browsing animals currently occupying KPGR (numbers supplied by owner)

Species	Height (m)			Animals	BU representation
	< 1.5	< 2	< 5		
Giraffe	Y	Y	Y	11	57
Eland	Y	Y	N	45	99
Kudu	Y	Y	N	50	50
Zebra	Y	N	N	12	0
Impala	Y	N	N	7	1

Gemsbok	Y	N	N	4	0
Total				129	207

At the time of the study, KPGR was stocked with 129 animals of six different species, which is equivalent to 207 browser units (Table 5.7). All species are able to utilize available browse below 1.5 m. However, only 138 BU can be sustained. All represented species, too, are able to browse up to 2 m (Table 5.7). However, an estimate of only 185 BU can be sustained within this height stratum. Available browse between 2 and 5 m, heights at which giraffes are capable of reaching, allows an additional 36 BU that can be supported.

5.3.3. Tsumkwe Private Game Reserve

5.3.3.1. Available browse

From Table 5.5, it is observed that the no and high utilization areas had the lowest and highest plant density per hectare respectively. This was similar in the case of important diet species, with plant density being the lowest and highest in the no and high utilization areas respectively. (Table 5.6). Evapotranspiration tree equivalents per hectare was lowest in the no utilization area and highest in the medium utilization area (Table 5.5). However, ETTE of important diet species per hectare was highest in the low utilization area, with the lowest ETTE per hectare found within the no utilization area (Table 5.6).

Total dry matter mass was lowest in the no utilization area, compared to the medium utilization area where it was the highest (Table 5.5). However, total dry matter mass of important diet species per hectare was highest in the low utilization area, with total dry matter mass being second highest within the high utilization area (Table 5.6).

Grewia robusta was absent in the no utilization area. However, *G. robusta* plant density, ETTE and total dry matter mass per hectare was highest in the high utilization area (Table 6.6). Similar results were observed for *S. longispina*. *Euclea undulata*, however, had the highest plant density, ETTE and total dry matter mass per hectare in the low utilization area. Similar results were observed for *L. ferocissimum*. *Lycium cinereum*, however, had the highest plant density, ETTE and total dry matter mass per hectare within the medium utilization area.

Table 5.7: Plant density (plants ha⁻¹), evapotranspiration tree equivalents per hectare (ETTE ha⁻¹) and total dry mass per hectare (kg DM ha⁻¹) for all species measured using the BECVOL method in the different utilization areas in TPGR

Utilization area	Plant density (Plants ha ⁻¹)	Evapotranspiration tree equivalents (ETTE ha ⁻¹)	Total browse mass (< 5 m) (kg DM ha ⁻¹)
No utilization	1 759	1 055	417
Low utilization	1 946	2 468	1 000
Medium utilization	4 854	3 950	1 571
High utilization	4 050	3 158	1 245

Table 5.8: Plant density (plants ha⁻¹), evapotranspiration tree equivalents per hectare (ETTE ha⁻¹) and total dry mass per hectare (kg DM ha⁻¹) for all important diet species measured using the BECVOL method in the different utilization areas in TPGR

Species	Plant density (Plants ha ⁻¹)				Evapotranspiration tree equivalents (ETTE ha ⁻¹)				Total browse mass (< 5 m) (kg DM ha ⁻¹)			
	Utilization area											
	No	Low	Moderate	High	No	Low	Moderate	High	No	Low	Moderate	High
<i>Euclea undulata</i>	13	147	40	80	105	1271	299	278	43	529	123	114
<i>Grewia robusta</i>	0	27	27	53	0	79	87	221	0	27	32	78
<i>Lycium cinereum</i>	147	13	160	93	27	11	168	56	10	4	67	22
<i>Lycium ferocissimum</i>	40	53	27	27	185	297	67	91	76	122	27	37
<i>Searsia longispina</i>	0	0	40	93	0	0	177	379	0	0	72	156
Total	200	440	294	346	317	1 658	798	1 025	129	682	321	407

5.3.3.2. Browsing capacity




Figure 5.5: Browsing capacity of the (top left) 1.5 m, (top right) < 2 m and (bottom) < 5 m height stratum within each utilization area in TPGR

Hypothesis 1: Browsing capacity will not differ between the different areas.

In TPGR, more hectares are required to sustain a browser unit in the no utilization area, compared to the low ($p < 0.001$), medium ($p < 0.001$) and high ($p < 0.001$) utilization areas, due to the low density of important diet species within the no utilization area (Table 5.6). Although the density of principal diet species tends to increase from the no to the high utilization areas (Table 5.6), fewer hectares are required to sustain a browser unit within the low utilization area, than within the no ($p < 0.001$), medium ($p < 0.001$) and high ($p < 0.001$) utilization areas (Figure 5.5). The high utilization area contains the highest principal diet species density per hectare (Table 5.6), with fewer hectares required to sustain a single browser unit within this area, than what is required in the low ($p < 0.001$) and medium ($p < 0.001$) utilization areas.

Hypothesis 2: Browsing capacity will decrease as the height stratum increases.

It is evident that as the strata height increases, there is a decrease in browsing capacity (Figure 5.5). In other words, more hectares are required to sustain a single browser unit browsing at a height less than 1.5 m than a browser unit browsing at a height less than 2 m ($p < 0.001$) and 5 m ($p < 0.001$).

Hypothesis 3: Browsing capacity will remain the same between the different seasons.

No seasonal change in browsing capacity was evident ($X^2 = 5.079$; $p = 0.166$) (Figure 6.5).

5.3.3.3. Number of browser units sustained

The number of calculated browser units that can be sustained, as well as the different browsers currently found in TPGR can be observed in Tables 5.8 and 5.9 respectively. However, the number of animals currently occupying the area were not supplied by the owner. Giraffe numbers (5 giraffe) were the only numbers available due to the study being based on these individuals.

Table 5.9: Table representing the browsing capacity, browser and giraffe numbers that can be sustained in TPGR

Month	Browsing capacity (ha/BU)			Total browser units sustained			Number of giraffe sustained		
	< 1.5 m	< 2 m	< 5 m	< 1.5m	< 2m	< 5 m	< 1.5 m	< 2 m	< 5 m
January	28.5	20.5	17.5	53	73	86	10	14	16

February	28.5	20.5	17.5	53	73	86	10	14	16
March	28.5	20.5	17.5	53	73	86	10	14	16
April	28.5	20.5	17.5	53	73	86	10	14	16
May	28.9	20.7	17.7	52	72	85	10	14	16
June	29.1	20.8	17.8	52	72	84	10	14	16
July	29.5	21	17.9	51	71	84	10	14	16
August	29.8	21.2	18	50	71	83	10	14	16
September	29.8	21.2	18	50	71	83	10	14	16
October	28.9	20.7	17.7	52	72	85	10	14	16
November	28.5	20.5	17.5	53	73	86	10	14	16
December	28.5	20.5	17.5	53	73	86	10	14	16

As observed in Table 5.8, the following browser units that can be sustained by browse at different heights are as follows:

- 1.5 m: average of 53 BU
- < 2 m: average of 73 BU
- < 5 m: average of 86 BU
 - < 5 m = 86 BU = 17 giraffes, therefore browsing capacity = 1500 ha/221 BU = 88 ha per giraffe (< 5 m)




Table 5.10: Animal species currently present on TPGR

Species	Height (m)			BU representation
	< 1.5	< 2	< 5	
Giraffe	Y	Y	Y	26
Eland	Y	Y	N	2.2
Kudu	Y	Y	N	1
Springbok	Y	N	N	0
Wildebeest	Y	N	N	0

Gemsbok	Y	N	N	0
Roan antelope	Y	N	N	0
Total				29

As observed in Table 5.9, 26 BU (5 giraffe) currently occupy the property. Giraffe have the ability to browse below 1.5 m, 2 m and 5 m. Therefore, a total of 26 BU needs to be supported up to 5 m.

5.4. DISCUSSION

Browsing capacity will not differ between the different areas.

Browsing capacity decreased the more the area was utilised in both KPGR and TPGR, due to the increase in density of principal diet species. On the same scale, the selection of plants displays diet selection, with diet selection ultimately leading to habitat selection on a larger scale (Duncan and Gordon, 1999). However, forage is one factor, amongst many, that determines habitat selection (Janecke and Smit, 2011).

Herbivores tend to spend more time in certain areas due to the limitations in energy requirements (Fabricius, 1989). Ungulates tend to limit their time spent travelling in search of forage by selecting areas with an adequate amount of preferred species (Fabricius, 1989). Therefore, more time is spent feeding and less time travelling (Senft *et al.*, 1987; Fabricius, 1989). For example, in the Northern Cape, eland, which are mixed feeders, prefer areas with a high available biomass of leaves (Fabricius, 1989).

In this study, the utilization of habitats by giraffes is similar. Giraffes tend to congregate and utilise areas more where there is an availability and abundance of preferred plant species (Berry, 1973; van der Jeugd and Prins, 2000; Fennessy, 2004; Deacon, 2015). In Lake Manyara National Park, Tanzania, although giraffe densities differed between different habitats, higher densities were found in the most vegetated areas (van der Jeugd and Prins, 2000). Giraffe populations in Luangwa Valley, Zambia were similar (Berry, 1978). A narrow, elongated home range existed along the Luangwa River, favoured due to the variety and quantity of food throughout the year. Areas lacking in palatable or preferred plant species were less favoured (Berry, 1973).

Utilization of certain areas by giraffe populations in KPGR and TPGR may, too, be due to the presence of principal diet species. In KPGR, a high abundance of the three species forming the bulk of the giraffes' diet, namely *P. capensis*, *P. afra* and *E. undulata* (Chapter 5), were found within the high utilization areas. This area is found within the Grootkop Arid Spekboomveld vegetation type, dominated by the indicator species *P. afra*, as well as the other key species (Chapter 3). TPGR was similar, with the two principal species, *G. robusta* and *S. longispina*, found in greater abundances within the medium and high utilization areas. These areas are positioned within the Kruisrivier Spekboom-Pruimveld vegetation type (Chapter 2), which is dominated by the presence of these species.

In contrast, the no or low utilization of certain areas were a result of the absence of preferred species. In KPGR, *P. capensis*, the most preferred species, was absent within the no utilization area. However, *P. afra*, was highest within this area. This could be due to herbivores not utilizing the area due to its positioning at the bottom of a rocky, steep slope. Giraffes tend avoid rocky habitats (Fennessy, 2004). In TPGR, even though principal diet species, such as *E. undulata* and both *Lycium* species, are present within the no utilization area, the two most preferred species, *G. robusta* and *S. longispina* are absent.

Browsing capacity will decrease as the height stratum increases.

Browse capacity in both study sites decreased as the height stratum of plants increased. This is not surprising, due to the structure of plants, particularly large shrubs and trees having canopies with the majority of their biomass positioned higher above the ground. In South Africa, Woolnough and Du Toit (2001) concluded that leaf dry mass increased with increasing height up the canopy. Giraffe browsing further supported this, with giraffe feeding efficiency increasing higher up the canopy and giraffe obtaining a higher plant matter dry mass per bite. Similar results were observed in KwaZulu-Natal, South Africa, where browse production was highest in the 2.5 – 5 m height stratum, compared to the 0.0 – 0.5 m height stratum which had the lowest browse production (Penderis and Kirkman, 2014).

Browsing capacity will remain the same between the different seasons.

Forage availability, a determinant of habitat selection, may be influenced by seasonal and phenological change in plant species (Fennessy, 2004; Janecke and Smit, 2011; Penderis and Kirkman, 2014; Deacon, 2015). Within giraffes' natural ranges, often dominated by deciduous species, food availability decreases and increases within the dry and wet seasons respectively

(Eamus, 1999). To counter this problem, giraffes need a larger area to meet their daily requirements, and tend to expand their home ranges in search of food (Deacon, 2015).

However, in game ranches, herbivores do not have the advantage of being able to roam over vast areas in search of food during these critical periods due to confined spaces (Janecke and Smit, 2011). Ranchers often resort to hunting or supplementary feeding to compensate for the shortage in browse. However, KPGR and TPGR do not experience this problem, due to the majority of plants within the area being of an evergreen nature. Unlike deciduous plant species shedding leaves as a response to low moisture availability and high evaporative demands during the dry season, evergreen species maintain water transport to their canopies at a sufficient rate (Penderis and Kirkman, 2014). This allows evergreen species to maintain a full canopy all year round, with a less than 10% canopy biomass lost during the dry season (Penderis and Kirkman, 2014). This is highly advantageous, due to browse being available all year round to herbivore populations, and, thus, browsing capacity remaining constant seasonally, as in the case of KPGR and TPGR.

Several recommended stocking rates exist for giraffe in Thicket vegetation. However, the majority of these stocking rates are not based on quantitative research. Within the Western Cape, stocking rates of 19 properties ranged from 30 to 1650 ha per giraffe (Cornelius, 2010). These calculations were based on the size of the properties, and not taking the suitability of the habitat into consideration. Cornelius (2010), however, based browsing capacity estimates within the Western Cape on giraffe observations and vegetation surveys, with a recommended 78.5 ha being the minimum area required to sustain one giraffe. Browsing capacity estimates within KPGR (25 ha/giraffe < 2 m and 21 ha/giraffe < 5 m) and TPGR (107 ha/giraffe < 2 m and 88 ha/giraffe < 5 m) are observed to be lower and higher respectively. However, the recommendation of less than two metres should be considered due to the continuous browsing of both giraffe populations within this height stratum, even though their height allows them to browse up to five metres (Deacon, 2015). In addition, browsing capacities should be adjusted accordingly to the number of other herbivores present on the property. This is suggested to avoid the possibility of interspecific competition (Leuthold and Leuthold, 1978; Deacon, 2015).

5.5. CONCLUSION

There is much confusion regarding the stocking rate of giraffe due to their extralimital status in the Western Cape. This study found KPGR to have a browsing capacity of 25 ha and 21 ha

per giraffe at a browse height stratum of two and five metres respectively. In contrast, TPGR rendered browsing capacities of 107 and 88 ha per giraffe at a browse height stratum of two and five metres respectively. Although these farms are situated in close proximity in the same biome, these values are far from similar. It is therefore seen that giraffe browsing capacities need to be based on quantifiable research and need to be site specific due to the vast differences and heterogeneity of areas within the same vegetation biome.

Both these game farms need knowledge and information to be proactive in their decisions. The academic implications of this study need to be considered as an on-going process as new information becomes available. This project was aimed at developing new skills and techniques for research to address complex animal-environment interactions with applicability to other similar regions of southern Africa.

Combining research and management ideas will develop conservation management plans that will enable scientists to conduct future research and demonstrate that management options are determined by the intention to limit giraffes from future mortalities because of low quality foliage.



CHAPTER 6 : GENERAL CONCLUSIONS, MANAGEMENT IMPLICATIONS AND FUTURE RESEARCH

6.1. INTRODUCTION

The general aim of the study was to determine the general and foraging behaviour, as well as site-specific browsing capacities of extralimital giraffe in the Little Karoo of the Western Cape, South Africa. Literature available on this subject is lacking, regardless of giraffe continuously being introduced into the area despite the area being outside their natural distribution. The study therefore sought to achieve the following objectives:

1. To determine the diurnal activity budgets,
2. the diet composition and browsing levels, and
3. the availability of browse and estimated site specific browsing capacities of extralimital giraffe.

6.2. EMPIRICAL FINDINGS

Objective 1: To determine the diurnal activity budgets of extralimital giraffe

In both study sites, browsing was the dominant behaviour displayed by giraffes, followed by walking, rumination, rest and vigilance. However, contrasting results were shown between the study sites regarding sex and seasonal differences. In KPGR, males rested, walked and ruminated more than females, with females browsing more than males. However, in TPGR, males were only found to walk more than females, with females browsing more than males.

Contrasting seasonal differences were also shown between the two study sites. In KPGR, females, males and juveniles spent more time on energy conserving activities, such as resting and rumination during the cooler seasons, whereas these activities occurred more during the warmer seasons in TPGR. Females, however, spent more time on energy consuming activities, such as browsing during the cooler seasons, whereas males browsed more during the warmer seasons in both study sites.

Objective 2: To determine the diet composition and browsing levels of extralimital giraffe

Fifteen plant species were consumed in KPGR, with four principal species, namely *P. capensis*, *P. afra*, *E. undulata* and *Lycium spp.* forming approximately 90% of their diet. The importance of *Lycium spp.* and *R. obovatum* increased during spring and summer. On the other hand, 12 plant species were consumed in TPGR, with five principal species, namely *S. longispina*, *E.*

undulata, *P. capensis*, *V. karroo* and *G. robusta* forming approximately 90% of their diet. The importance of *V. karroo*, *R. obovatum* and *G. robusta* increased during spring and summer.

In both KPGR and TPGR, males browsed more frequently at L3. However, in KPGR, males lowered their browsing level and browsed more at L2 from winter to autumn, whereas in TPGR, males raised their browsing level and browsed more at L5 from winter to autumn. Females in KPGR generally browsed most frequently at L2, but raised their dominant browsing levels to L3 and L4 during winter and spring respectively. Females in TPGR, however, browsed more frequently at L3, decreasing their browsing level and browsing more at L2 during spring and autumn.

Objective 3: To determine the availability of browse and estimated site-specific browsing capacities of extralimital giraffe

The calculated browsing capacity for KPGR and TPGR was 25 and 21 and 107 and 88 ha per giraffe respectively for the respective browse height strata of two and five metres. KPGR and TPGR are therefore within the estimated browsing capacities, however, the recommended browsing capacity of less than two metres should be considered due to the continuous browsing of both giraffe populations at lower levels, and should additionally be adjusted according to the presence of other browsers.

6.3. RESEARCH IMPLICATIONS

This thesis reveals that giraffes are able to adapt to new environments regardless of their historical distribution. By introducing giraffes into new areas, giraffes have taken advantage of browse species different to those consumed within their natural ranges (Hall-Martin, 1974; Dagg and Foster, 1976; Sauer *et al.*, 1977; Pellew, 1983; Caister *et al.*, 2003; Fennessy, 2004; Deacon, 2015). In addition, their survival and ability to produce offspring is an indicator that the quality of available browse is suitable. However, the absence of a distinct dietary shift between seasons in both KPGR and TPGR (Chapter 4) may pose a problem. Within their natural range, plant species composition consists of a mixture of deciduous and evergreen species (Deacon, 2015), resulting in a distinct dietary shift and allowing “rest” of plant species once the quantity of leaves decrease. However, in the study region, the continuous utilization of the same plant species throughout the year might have some negative effects. Therefore, this study suggests continuous examination of the utilization of principal browse species by extralimital giraffe.

Monitoring is an effective tool if used correctly (Hurt and Hardy, 1989). Since the area is outside historical ranges, and giraffes have been shown to cause negative impacts (Foster and Dagg, 1972; Hall-Martin, 1974; Van Aarde and Skinner, 1975; Sauer *et al.*, 1977; Pellew, 1983; Augustine and McNaughton, 1998; Birkett, 2002; Brenneman *et al.*, 2009), monitoring will be extremely beneficial, as it will provide indications of browse lines, heavy decrease in leaf quantity, or the stunted growth of plant individuals (Cornelius, 2010). It will also indicate any niche overlaps and low browsing by extralimital giraffe (Chapter 5), resulting in possible interspecific competition with indigenous herbivores.

The study also reveals that browsing capacities for extralimital giraffe can not be generalised and need to be site specific. Although KPGR and TPGR are situated relatively close to one another, the vegetation types present on both farms are different, although situated in the same biome. Taking this, as well as the low browsing and potential impact of extralimital giraffe into consideration, it is suggested that the minimal policy regarding this species be revised. Requirements for the ownership of giraffe is only adequate fencing, as stated in fencing specifications as per Cape Nature's minimum specifications for adequate enclosure (dated 14 October 1976, as amended). To avoid the deterioration of vegetation, and to prevent negative effects on indigenous herbivores, this policy needs to be revised and stronger measures put in place.

6.4. RECOMMENDATIONS FOR FUTURE STUDIES

Upon completion of the study, it can be seen that there is a need for further research to facilitate the attainment of the goal of the study. Recommendations for future studies are as follows:

6.4.1. Monitoring

Long term monitoring is required for more detailed information on the behaviour and diet of extralimital giraffe, as well as more accurate browsing capacity calculations. In addition, further studies should include the monitoring of other extralimital giraffe throughout the Western Cape and beyond. Furthermore, monitoring should include vegetation monitoring. Vegetation monitoring will help determine changes in vegetation, and provide information on giraffe impacts.

6.4.2. Quantification of available browse and browsing capacity

Easier vegetation sampling methods for quantifying available browse and browsing capacity within Thicket areas needs to be developed due to the dense and intertwined nature of plant

species. Simplified and repeatable techniques will be beneficial especially to farmers who do not have time for tedious methods.

6.5.3. Habitat quality

Future studies on habitat quality are required to provide information on why extralimital giraffe prefer certain plant species and avoid others. These diet preference studies will also determine habitat preferences, allowing for the gaps on the spatial ecology of extralimital species to be filled. In addition, determining habitat qualities may help in habitat and vegetation conservation.

6.5.4. Policies

All research should try and be implemented into policy. More policy on the transportation and ownership of extralimital giraffe will help maintain genetic integrity and prevent inbreeding of the South African giraffe.

6.5. LIMITATIONS OF THE STUDY

The study has offered important baseline knowledge on the adaptive behaviour and diet selection of extralimital giraffes in the Western Cape. However, because of the methodology used, the study encountered two major limitations. Firstly, logistics only allowed for four days of observations per site every three months. This resulted in a relatively small sample size per season. Future studies should try to increase the frequency of observations.

The second limitation was accessibility. Vehicles could not get into certain rocky, steep areas. Therefore, observations in these areas had to be conducted on foot. Reaching these areas by foot often took time, and as a result, observations were lost.

6.6. CONCLUSION

This study has provided previously lacking baseline data about the activity budgets, diet selection and proposed site-specific browsing capacities for extralimital giraffe. In addition, it has provided a basis for future studies to build on. Farmers can reap the benefits this species brings, if managed correctly, and if stronger policies are put in place to avoid potential and negative effects on indigenous flora and fauna.

REFERENCE LIST

- Adolfsson UG. 2009. Diurnal behaviour and utilization of shade in Masai giraffes (*Giraffa camelopardalis tippelskirchi*). Degree project. Swedish University of Agricultural Sciences, Uppsala.
- Altmann J. 1974. Observational study of behaviour: sampling methods. *Behaviour* 49: 227-267.
- Anderson TA. 2004. A beginner's guide to the plants of Kimberley and surrounds, with special reference to Magersfontein Battlefield. McGregor Museum & ShareNet.
- Artiss T, Martin K. 1995. Male vigilance in white-tailed ptarmigan, *Lagopus leucurus*: mate guarding or predator detection? *Animal Behaviour* 49:1249–1258.
- Aucamp A, Tainton N. 1984. *Veld management in the valley bushveld of the Eastern Cape*. Government Printer. Pretoria.
- Augustine DJ, McNaughton SJ. 1998. Ungulate effects on the functional species composition of plant communities: Herbivore selectivity and plant tolerance. *Journal of Wildlife Management* 62:1165-1183.
- Baer DJ, Oftedal OT, Fahey GC. 1985. Feed selection and digestibility by captive giraffe (*Giraffa Camelopardalis*). *Zoo Biology* 4(1):57-64.
- Beckman JH, Prins HHT. 1989. Feeding strategies of sedentary large herbivores in East Africa, with emphasis on the African buffalo, *Syncerus coffer*. *African Journal of Ecology* 27(2): 129-147.
- Ben-Shahar R, Fairall N. 1987. Comparison of the diurnal patterns of blue wildebeest and red hartebeest. *South African Journal of wildlife research* 17(2).
- Bell RHV. 1971. A grazing ecosystem in Serengeti. *Scientific American* 225: 86-93.
- Berry PSM. 1973. The Luangua valley giraffe. *Puku* 7:71-92.
- Berry PSM, Bercovitch FB. 2012. Darkening coat colour reveals life history and life expectancy of male Thornicroft's giraffes. *Journal of Zoology* 287 (3): 157-160.
- Birkett A. 2002. The impact of giraffe, rhino and elephant on the habitat of a black rhino sanctuary in Kenya. *African Journal of Ecology* 40:276-282.

- Blomqvist, PA, Renberg L. 2007. Feeding behaviour of Giraffe (*Giraffa camelopardalis*) in Mokolodi Reserve, Botswana: University of Uppsala, Minor Field Study:1653-5634.
- Bond WJ, Loffell D. 2001. Introduction of giraffe changes acacia distribution in a South African savanna. *African Journal of Ecology* 39:286-294.
- Boshoff AF, Kerley RM, Cowling RM, Wilson SL. 2001. Conservation planning in the Greater Addo National Park: The potential distributions, and estimated spatial requirements and population sizes, of the medium- to large-sized mammals. Terrestrial Ecology Research Unit Report no. 33. University of Port Elizabeth, Port Elizabeth.
- Bothma J, Van Rooyen N, Van Rooyen MW. 2004. Using diet and plant resources to set wildlife stocking densities in African savannas. *Wildlife Society Bulletin* 32(3): 1-12.
- Brenneman, RA, Bagine, RK, Brown DM, Ndeti R, Louis EE. 2009. Implications of closed ecosystem conservation management: the decline of Rothschild's giraffe (*Giraffa camelopardalis rothschildi*) in Lake Nakuru National Park, Kenya. *African Journal of Ecology* 47: 711-719.
- Bryant JP, Provenza FD, Pastor J, Reichardt PB, Clausen TP, du Toit JT. 1991. Interactions between woody plants and browsing mammals mediated by secondary metabolites. *Annual Review of Ecology and Systematics* 22: 431-446.
- Caister LE, Shields WM, Gosser A. 2003. Female tannin avoidance: a possible explanation for habitat and dietary segregation of giraffes (*Giraffa camelopardalis perlata*) in Niger. *African Journal of Ecology* 41:201-210.
- Cameron EZ, Du Toit JT. 2007. Winning by a neck: Tall giraffes avoid competition with shorter browsers. *American Naturalist* 169:130-135.
- Carnaby T. 2010. *Beat About The Bush Mammals*. Jacana Publishers. pp 376
- Caroline AD, Adhiambo WPJ. 2012. Movement patterns and home range sizes of the Rothschild's giraffes (*Giraffa Camelopardalis rothschildii*) translocated to Ruma National Park, Kenya. *The International Journal of Engineering and Science* 2(7):14-22.
- Castley JG, Boshoff AF, Kerley GIH. 2001. Compromising South Africa's natural biodiversity-inappropriate herbivore introductions. *South African Journal of Science* 9(7):344-348.

- Chapuis J, Bousses P, Pisanu B, Reale D. 2001. Comparative rumen and faecal diet microhistological determinations of European mouflon. *Journal of Range Management* 54:239-242.
- Chenje M, Mohamed-Katerere J. 2006. Invasive alien species. *Africa Environment Outlook 2*. Malta: Progress Press Ltd. pp 331-372.
- Cilliers, G., & Kok, O.B. 1994. Feeding behaviour and activity aspects of giraffe (*Giraffa camelopardalis*) in the Franklin Nature Reserve, South Africa. Department of Zoology and Entomology, University of the Orange Free State, South Africa.
- Ciofolo I, Le Pendu Y. 2002: The feeding behaviour of giraffe in Niger. *Mammalia* 66: 183-194.
- Clavero M, Garcia-Berthou E. 2005. Invasive species are a leading cause of animal extinctions. *Trends in Ecology and Evolution* 20: 110.
- Coe MJ. 1967. "Necking" behaviour in the giraffe. *J. Zool. Lond.* 151:313-321.
- Cooke HBS. 1974. The fossil mammals of Cornelia, OFS, South Africa. *Mem. Nat. Mus. Bloem.* 9:63-88.
- Cooper SM, Owen-Smith N, Bryant JP. 1988. Foliage acceptability to browsing ruminants in relation to seasonal changes in the leaf chemistry of woody plants in a South African savanna. *Oecologia* 75: 336-342.
- Cornelius AJ. 2010. The management of extralimital giraffe (*Giraffa camelopardalis*) in the Mosaic Thicket of the southern Cape, South Africa. MSc thesis. Nelson Mandela Metropolitan University, South Africa.
- Cupido CF. 2005. Assessment of veld utilisation practices and veld condition in Little Karoo. MSc thesis. University of Stellenbosch, South Africa.
- Daehler CC, Gordon DR. 1997. To introduce or not to introduce: Trade-offs of non-indigenous organisms. *Trends in Ecology and Evolution* 12: 424-425.
- Dagg AI. 1960. Food preferences of the giraffe. *Zoology Society of London* 135:640-642.
- Dagg AI. 1971. *Giraffa Camelopardalis*. *Mamm Species* 5:1-8.
- Dagg AI. 2014. *Giraffe: Biology, Behavior, and Conservation*. Cambridge: Cambridge University Press

- Dagg AI, Foster JB. 1976. *The giraffe, its biology, behaviour, and ecology*. New York: Van Nostrand Reinhold Company.
- Dagg AI, Foster JB. 1982. *The Giraffe: its Biology, Behavior and Ecology*. Krieger: Malabar, Florida.
- Danckwerts JE. 1981. A technique to assess the grazing capacity of sweetveld with particular reference to the False Thornveld areas of the Ciskei. MSc thesis, University Natal, Pietermaritzburg, South Africa.
- Dayton BR. 1978. Standing crops of dominant *Combretum* species at three browsing levels in the Kruger National Park. *Koedoe* 21: 67-76.
- Deacon F. 2015. The spatial ecology, habitat preferences and diet selection of giraffe (*Giraffa camelopardalis giraffa*) in the Kalahari region of South Africa. PhD thesis. University of the Free State, South Africa.
- Deacon F, Smit GN. 2012. Movement patterns and impact of giraffe (*Giraffa camelopardalis*) on the woody plants of a fenced area in the central Free State. *Giraffa* 6:2-22.
- Desmet PG, Cowling RM. 1999. The climate of the karoo – a functional approach. In: Dean WRJ, Milton SJ (eds), *The Karoo – Ecological Patterns and Processes*. Cambridge: Cambridge University Press. pp 3–16.
- Duncan AJ, Gordon IJ. 1999. Habitat selection according to the ability of animals to eat, digest and detoxify foods. *Proceedings of the Nutrition Society* 58: 799-805.
- Du Toit JT. 1988. Patterns of resource use within the browsing ruminant guild in the central Kruger National Park. PhD thesis, University of the Witwatersrand, Johannesburg.
- Du Toit JT. 1990. Home range – body mass relations: a field study on African browsing ruminants. *Oecologia* 85: 301-303.
- Eamus D. 1999. Ecophysiological traits of deciduous and evergreen woody species in the seasonally dry tropics. *Tree* 4:11-16.
- ENPAT: Environmental Potential Atlases. 1995. User's Reference and Database Guides. Department of Environmental Affairs and Tourism.
- Evans NEM. 1970. The reaction of a group of Rothchild's giraffe to a new environment. *East African Wildlife Journal* 8:53-62.

- Fabricius C. 1989. Habitat suitability assessment for indigenous browsing ungulates in the Northern Cape. MSc thesis. University of the Witwatersrand, South Africa.
- Fairall N. 1968. The reproductive seasons of some mammals in the Kruger National Park. *Zoologica Africana* 3:189-210.
- Fennessy J. 2009. Home range and seasonal movements of *Giraffa camelopardalis angolensis* in the northern Namib Desert. *African Journal of Ecology* 47: 318-327.
- Ferreira N. 1970. Game digest for giraffe, *Giraffa camelopardalis* Linn. B.Sc. - Hons. Project. University of Pretoria, South Africa.
- Field CR, Ross IC. 1976. The savanna ecology of Kidepo National Park II. Feeding ecology of elephant and giraffe. *East African Wildlife Journal* 14:1-15.
- Foster JB. 1966. The giraffe of the Nairobi National Park: Home range, sex relations, the herd and food. *East African Wildlife Journal* 4:139-148.
- Foster JB, Dagg AC. 1972. Notes on the biology of the giraffe. *East African Wildlife Journal* 10:1-16.
- Fourie PF. 1977. Enkele aspekte van die identiteit, verspreiding, gedrag en voeding van die kameelperd *Giraffa camelopardalis giraffa* (Boddaert 1785) in die Nasionale Krugerwildtuin. MSc thesis, University of Potchefstroom, South Africa.
- Fritz H, Duncan P, Gordon IL, Illius AW. 2002. Megaherbivores influence trophic guilds structure in African ungulate communities. *Oecologia* 131(4): 620-625.
- Furstenburg D. 1991. The influence of plant tannin on the feeding ecology of giraffe (*Giraffa camelopardalis*). MSc Dissertation. University of Pretoria, South Africa.
- Furstenburg D. 2003. Kameelperd, *Giraffa camelopardalis*. Wild en Jag, Jul.8-13.
- Furstenburg D, Van Hoven W. 1994 Condensed tannin as anti-defoliate agent against browsing by giraffe (*Giraffa camelopardalis*) in the Kruger National Park. *Comp. Biochem. Physiol.* 107A, 425-431.
- Ginnett T, Demment MW. 1997. Sexual segregation in giraffe foraging behavior at two spatial scales. *Oecologia* 110:291-300.
- Griesel J. 1961. A new game sanctuary for the Orange Free State, The Willem Pretorius Game Reserve. *Afr. Wild Life* 15: 121-125.

- Hall-Martin AJ. 1974. A note on the seasonal utilisation of different vegetation types by giraffe. *South African Journal of Science* 70:122-123.
- Hall-Martin AJ. 1975. Studies on the Biology and Productivity of the Giraffe, *Giraffa camelopardalis*. MSc. Thesis. University of Pretoria, South Africa.
- Hall-Martin AJ, Basson WD. 1975. Seasonal chemical composition of the diet of Transvaal Lowveld giraffe. *Journal of Southern Africa Wildlife Management Association* 5:19-21.
- Hargens AR, Millard RW, Pettersson K. 1987. Gravitational haemodynamics and oedema prevention in the giraffe. *Nature* 329: 59-60.
- Hilton-Taylor C. 1996. Patterns and characteristics of the flora of the Succulent Karoo biome, southern Africa. In: Van der Maesen LJG, Van der Burgt XM, Van Medenbachde Rooy JM (eds), *The Biodiversity of African Plants*. Dordrecht: Kluwer Academic Publishers. pp 58-72.
- Hoffman MT. 1989. A preliminary investigation of the phenology of subtropical thicket and Karroid shrubland in the lower Sundays River Valley, SE Cape. *South African Journal of Botany* 55:586-597.
- Hoffman MT, Cowling RM. 1990. Desertification in the lower Sundays River Valley. *South African Journal of Arid Environments* 19: 105-117.
- Hofmann RR (eds). 1973. *The ruminant stomach (stomach structure and feeding habits of East African game ruminants)*. East African Monographs in Biology, vol 2. East African Literature Bureau, Nairobi
- Hofmann RR. 1989. Evolutionary steps of ecophysiological adaptation and diversification of ruminants: A comparative view of their digestive system. *Oecologia* 78: 443-457.
- Hudson RJ, White RG (eds). 1985. *Bioenergetics of Wild Herbivores*. CRC Press.
- IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp.
- Innis A. 1958. The behaviour of the giraffe, *Giraffa camelopardalis*, in the eastern Transvaal. *Proceedings of the Zoological Society of London* 131: 245-278.
- Jacobs EP. 2008. Diet and feeding effects of introduced giraffe in the Eastern Cape. MSc thesis. Nelson Mandela Metropolitan University, South Africa.

- Janecke BB, Smit GN. 2011. Phenology of woody plants in riverine thicket and its impact on browse availability to game species. *African Journal of Range & Forage Science*, 28(3): 139–148.
- Johnson DH. 1980. The comparison of usage and availability measurements for evaluating resource preference. *Ecology* 61: 65-71.
- Klasen SAH. 1963. Giraffe are strictly browsers. *Black Lechwe* 23-25.
- Kingdon J (eds). 1979. *East African Mammals: An Atlas of Evolution in Africa (Vol. 3, Part B: Large Mammals)*. Chicago, University of Chicago Press. pp 231-249.
- Kirk HD. 1966. Progress of Mlilwane Sanctuary, Swaziland, *Afri. Wild. Life* 20:313-320.
- Knight MH. 1991. Ecology of the gemsbok *Oryx gazella gazella* (Linnaeus) and blue wildebeest *Connochaetes taurinus* (Burchell) in the southern Kalahari. PhD thesis. University of Pretoria, South Africa.
- Kok OB, Opperman DPJ. 1980. Feeding behaviour of giraffe (*Giraffa camelopardalis*) in the Willem Pretorius Game Reserve, Orange Free State. *South African Journal of Wildlife Research* 10: 45-55.
- Kok OB, Opperman DPJ. 1985. Voerbesikbaarheid en voedingswaarde van die belangrikste voedselplante van die kameelperd *Giraffa camelopardalis* (Linnaeus, 1758) in die Willem Pretoriuswildtuin, Oranje-Vrystaat. *Koedoe* 28:17-34.
- Langman, VA. 1978. Giraffe pica behaviour and pathology as indicator of nutritional stress. *The Journal of Wildlife Management* 42, 141-147.
- Le Pendu Y, Ciofolo, I, Gosser A. 2000. The social organisation of giraffes in Niger. *African Journal of Ecology* 38: 78-85.
- Lechmere-Oertel RG, Kerley GIH, Cowling RM. 2005. Patterns of degradation in semi-arid Succulent Thicket, South Africa: What is the new stable state? *Journal of Arid Environments* 62: 459 – 474.
- Leuthold BM. 1979. Social organisation and behaviour of giraffe in Tsavo East National Park. *African Journal of Ecology* 19: 9-34.
- Leuthold BM, Leuthold W. 1978. Ecology of the giraffe in Tsavo East National Park, Kenya. *East African Wildlife Journal* 16: 1-20.

- Lubke R, De Moor IJ. 1998. *Field guide to the Eastern and Southern Cape Coasts*. Juta and Company Ltd.
- Maitre DL, Colvin C, Maherry A. 2009. Water resources in the Klein Karoo: the challenge of sustainable development in a water-scarce area. *South African Journal of Science* 105: 39-48.
- Mann G. 2014. The ecology of leopards (*Panthera pardus*) in the Little Karoo, South Africa. PhD thesis. Rhodes University, South Africa.
- Matson TK. 2003. Habitat use and conservation of the vulnerable black-faced impala (*Aepyceros melampus petersi*) of Namibia. PhD Thesis. University of Queensland, Australia.
- Mitchell G, Skinner JD. 2003. Giraffe myths and cravings. Scientists solve another evolutionary riddle. *African Wildlife* 57(1):12-13.
- Mooring MS, Hart BL. 1995. Costs of allogrooming in impala: distraction from vigilance. *Animal Behaviour*. 49: 1414-1416.
- Mucina L, Rutherford MC. (eds) 2006. *The Vegetation of South Africa, Lesotho and Swaziland, Strelitzia 19*. South African National Biodiversity Institute, Pretoria.
- Muller Z, Bercovitch F, Brand R, Brown D, Brown M, Bolger D, Carter K, Deacon F, Doherty JB, Fennessy J, Fennessy S, Hussein AA, Lee D, Marais A, Strauss M, Tutchings A, Wube T. 2016. *Giraffa camelopardalis* (errata version published in 2017). The IUCN Red List of Threatened Species 2016: e.T9194A109326950. <http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T9194A51140239.en>.
- Naeem S, Thompson LJ, Lawler SP, Lawton JH, Woodfin RM. 1994. Declining biodiversity can alter the performance of ecosystems. *Nature* 386: 734-736.
- Nesbit Evans EM. 1970. The reaction of a group of Rothchild's giraffe to a new environment. *East African Wildlife Journal* 8:53-62.
- Nongwe N. 2008. A systematic conservation assessment of habitat transformation and degradation in the Little Karoo, South Africa. MSc thesis. University of Cape Town, South Africa.
- Nowak RM, Paradiso JL. (eds) 1983. 4th edition. *Walker's mammals of the world Volume II*. The John's Hopkins University Press. Baltimore.
- Oates LG. 1970. Food preferences of giraffe in Transvaal Mopane woodland. *J. S. Afr. Wildl. Manage. Ass* 2:21-23.

- O'Connor DAF. 2013. Foraging ecologies of giraffe and camels in northern Kenya: Effects of habitat structure and possibilities for competition. MSc Thesis, University of Michigan, United States of America.
- Omphile, U.J. 1997. Seasonal diets, diet quality and habitat preferences of large, savanna ungulates in the Chobe National Park, Botswana. unpublished PhD thesis. University of Wyoming. Laramie.
- Owen-Smith N. 1982. *Factors influencing the consumption of plant products by large herbivores*. In Huntly, B.J. and Walker, B.H. (Eds.) *Ecology of Tropical Savannas*. Springer, Berlin.
- Owen-Smith N (eds.). 1988. *Megaherbivores: The influence of very large body size on ecology*. Cambridge University Press, Cambridge.
- Owen-Smith N. 1999. *The animal factor in veld management*. In: Tainton NM (eds) *Veld Management in South Africa*. University of Natal Press, Pietermaritzburg. pp 117-138.
- Parker DM, Bernard RTF, Colvin SA. 2003. The diet of a small group of extralimital giraffe. *African Journal of Ecology* 41: 245-253.
- Parker DM. 2004. The feeding biology and potential impact of introduced giraffe (*Giraffa camelopardalis*) in the Eastern Cape Province, South Africa. MSc thesis. Rhodes University, South Africa.
- Parker DM, Bernard RTF. 2005. The diet and ecological role of giraffe (*Giraffa camelopardalis*) introduced to the Eastern Cape, South Africa. *Journal of Zoology* 267: 203-210.
- Patterson JL, Goetz RH, Doyle JT, Warren JV, Gauer OH, Detweiler DK, Said SI, Hoernicke H, McGregor M, Keen EN, Smith MH, Hardie EL, Reynolds M, Flatt WP, Waldo DR. 1965. Cardiorespiratory dynamics in the ox and giraffe, with comparative observations on man and other mammals. *Annals of the New York Academy of Sciences* 127:393-413.
- Pellew RA. 1983. The impacts of elephant, giraffe and fire upon the *Acacia tortilis* woodlands of the Serengeti. *African Journal of Ecology* 21:41-74.
- Pellew RA. 1984a. Food consumption and energy budgets of the giraffe. *Journal of Applied Ecology* 21:141-159.
- Pellew RA. 1984b. The feeding ecology of a selective browser, the giraffe (*Giraffa camelopardalis tippelskirchi*). *J. Zool. Lond.* 202:57-81.

- Penderis CA, Kirkman KP. 2014. Savanna browse production. 1: Determinants and measurement. *African Journal of Range and Forage Science* 31: 13-24.
- Pienaar RC. 2013. The feeding ecology of extralimital nyala (*Tragelaphus angasii*) in the arid mosaic thicket of the southern Cape. Masters thesis. Nelson Mandela Metropolitan University, South Africa.
- Pierce SM, Cowling RM. 1983. Phenology of fynbos, renosterveld and subtropical thicket in the south eastern Cape. *South African Journal of Botany* 3:1-16.
- Potts AJ, Hedderson TA, Vlok JHJ, Cowling RM. 2013. Pleistocene range dynamics in the eastern Greater Cape Floristic Region: A case study of the Little Karoo endemic *Berkheya cuneate* (Asteraceae). *South African Journal of Botany* 88: 401-413.
- Pratt DM, Anderson VH. 1982. Population, distribution and behaviour of giraffe in the Arusha National Park, Tanzania. *Journal of Natural History* 16:481-489.
- Pratt DM, Anderson VH. 1985 Giraffe social behaviour. *Journal of Natural History* 19:771-781.
- Renouf D, Lawson JW. 1986. Harbour seal vigilance? Watching for predators or mates? *Biol Behav* 11:44-49.
- Sauer JJC. 1983. Food selected by giraffes in relation to changes in chemical composition of the leaves. *South African Journal of Animal Science* 13: 40-43.
- Sauer JJC, Theron GK, Skinner JD. 1977. Food preferences of giraffe *Giraffa camelopardalis* in the arid bushveld of the western Transvaal. *South African Journal of Wildlife Research* 7(2):53-59.
- Sauer JJC, Skinner JD, Neitz AW. 1982. Seasonal utilization of the leaves by giraffes *Giraffa camelopardalis*. And the relationship of the seasonal utilization to the chemical composition of the leaves. *South African Journal of Wildlife Research* 10: 45-55.
- Seeber PA, Ciofolo I, Ganswindt A. 2012. Behavioural inventory of the giraffe (*Giraffa camelopardalis*). *BMC Research Notes* 2012, 5:650.
- Senft RL, Coughenour MB, Bailey DW, Rittenhouse LR, Sala OE, Swift DM. 1987. Large herbivore foraging and ecological hierarchies: Landscape ecology can enhance traditional foraging theory. *Biological Science* 37(11): 789-799.
- Siegfried WR. 1980. Vigilance and group size in springbok. *Madoqua* 12(3): 151-154.

- Sigwela A, Kerley GIH, Mills AJ, Cowling RM. 2009. The impact of browsing-induced degradation on the reproduction of subtropical thicket canopy shrubs and trees. *South African Journal of Botany* 75: 262–267.
- Skead CJ. 1987. Vol. 2. Historical mammal incidence in the Cape Province (Eastern half of the Cape Province). Chief Directorate of Nature and Environmental Conservation. Cape Town.
- Skinner JD, Chimimba CT (eds). 2005. *The mammals of the southern African subregion*. Cambridge University Press, Cape Town.
- Sicks F. 2011. REM sleep in giraffe. Presented at the 1st Wild International Giraffe congress held at the Etosha Safari Lodge, Namibia, 4-7 July 2011.
- Singer R, Bone EL. 1960. Modern giraffes and the fossil giraffids of Africa. *Ann. S. Afr. Mus.* 45:375-548.
- Smit GN. 1989. The Qualitative tree formula: A comprehensive approach to the quantification of woody plant communities. *Journal of the Grassland Society of Southern Africa* 6: 186-191.
- Smit GN. 1996. BECVOL: Biomass Estimates from Canopy VOLUME (version 2) - users guide. Unpublished Manual. University of the Free State, Bloemfontein.
- Smit GN, Richter CGF, Aucamp AJ. 1999. *Bush encroachment: an approach to understanding and managing the problem*. In: Tainton NM (eds) *Veld management in South Africa*. University of Natal Press, Pietermaritzburg. pp 246-260.
- Smit GN. 2004. An approach to tree thinning to structure southern African savannas for long-term restoration from bush thickening. *Journal of Environmental Management*. 71: 179-191.
- Smit GN. 2006. *Calculation of grazing capacity and browse capacity for game species*. In: Schutte F (eds) *Game ranching in central South Africa*. Alberts Design and Marketing, Brandfort. pp 18-28.
- Smit GN. 2014. BECVOL 3: an expansion of the aboveground biomass quantification model for trees and shrubs to include the wood component. *African Journal of Range & Forage Science* 31(2).
- Spinage CA. 1968. *The book of the giraffe*. Collins, London.
- Stark PTB, Slabach BL. 2012. The Scoop on Eating Dirt. *Scientific American*, 306, 30-32

- Stephens JM. 1975. Food selected by a giraffe in Sable Park, Rhodesia. *Arnoldia Rhodesia* 28:1-7.
- Strydom HA, King ND (eds). 2009. *Fuggle and Rabie's Environmental Management in South Africa*. Juta and Company Ltd.
- Stuart-Hill GC. 1992. Effects of elephants and goats on the Kaffrarian succulent thicket of the Eastern Cape, South Africa. *Journal of Applied Ecology* 29,: 699–710
- Stuart-Hill GC. 1999. Succulent Bushveld. In: Tainton N M. (eds) *Veld management in South Africa*. University of Natal Press, Pietermaritzburg.
- Sweetapple PJ, Nugent G. 1998. Comparison of two techniques for assessing possum (*Trichosurus vulpecula*) diet from stomach contents. *New Zealand Journal of Ecology* 22:181-188.
- Theron ME. 2005. Voedingsgedrag van kameelperde (*Giraffa camelopardalis*) in die sentrale Vrystaat. MSc Thesis. University of the Free State, South Africa.
- Thompson MW, Vlok J, Cowling RM, Cundill SL, Mudau N. 2005. A land transformation map for the Little Karoo. Final Report Version 2. Critical Ecosystems Protection Fund, CAPE, Cape Town.
- Valeix M, Fritz E, Dubois S, Kanengoni K, Alleaume S, Said S. 2007. Vegetation structure and ungulate abundance over a period of increasing elephant abundance in Hwange National Park, Zimbabwe. *Journal of Tropical Ecology* 23:87-93.
- Van Aarde RJ, Skinner JD. 1975. The food and feeding behaviour of the giraffe (*Giraffa camelopardalis*) in the Jack Scott Nature Reserve. *Publications of the University of Pretoria* 97: 59-68.
- Van Der Jeugd HP, Prins, HHT. 2000 Movements and group structure of giraffe (*Giraffa camelopardalis*) in Lake Manyara National Park, Tanzania. *Journal of Zoology* 251: 15–21.
- Van Hoven W. 2010. *The natural diet of wild herbivores*. In: Bothma J du P and Du Toit JG (eds.) *Game ranch management*, 5th Edition. Van Schaik Publishers, Pretoria. pp 347-367.
- Van de Koppel J. and Prins HHT. 1998. The importance of herbivore interactions for the dynamics of African savannah woodlands: a hypothesis. *Journal of Tropical Ecology* 14:565–576.

- Vlok JHJ, Euston-Brown DIW. 2002. Biological Survey Report (plants and Birds). Unpublished Report for the Subtropical Thicket Ecosystem Planning Project (STEP). Terrestrial Ecology Research Unit, Department of Zoology. University of Port Elizabeth, South Africa. www.zoo.upe.ac.za/step.
- Vlok JHJ, Cowling RM, Wolf T. 2005. A vegetation map for the Little Karoo. Succulent Karoo Ecosystem Planning (SKEP) Project (Grant no-1064410304).
- Ward D, Young TP. 2002. Effects of large mammalian herbivores and ant symbionts on condensed tannins of *Acacia drepanolobium* in Kenya. *Journal of Chemical Ecology* 28:913-929.
- Watson A. 1999. Conservation of Biodiversity Resource Areas (COBRA) Project: Kenya (1992–1998), Washington, DC: Development Alternatives Incorporated.
- Woolnough AP, Du Toit JT. 2001. Vertical zonation of browse quality in tree canopies exposed to a size-structured guild of African browsing ungulates. *Oecologia* 129:585-590.
- Wyatt JR. 1969: The feeding ecology of giraffe (*Giraffa camelopardalis* L.) in Nairobi National Park, and the effect of browsing on their main food plants. MSc thesis. University of East Africa, Nairobi.
- Yáber MC, Herrera EA. 1994. Vigilance, group size and social status in Capybaras. *Animal Behaviour* 48(6):1301-1307
- Zinn AD, Ward D, Kirkman K. 2007 Inducible defences in *Acacia sieberiana* in response to giraffe browsing. *African Journal of Range and Forage Science* 24:123-129

APPENDICES

Appendix A




Figure A: Individuals of giraffe herd on Kareesbos Private Game Reserve

Appendix B




Figure B: Individuals of giraffe herd on Tsumkwe Private Game Reserve