



International Journal of Fisheries and Aquatic Studies

E-ISSN: 2347-5129

P-ISSN: 2394-0506

(ICV-Poland) Impact Value: 5.62

(GIF) Impact Factor: 0.549

IJFAS 2020; 8(3): 372-378

© 2020 IJFAS

www.fisheriesjournal.com

Received: 10-03-2020

Accepted: 12-04-2020

Ayofe M Hamed

Department of Fisheries,
Faculty of Science, Lagos State
University. P.O. Box 0001,
LASU Post Office, Ojo, Lagos,
Nigeria

Albert O Amosu

Department of Agricultural
Science, School of Vocational &
Technical Education, Adeniran
Ogunsanya College of Education,
PMB 007 Festac Town,
Otto/Ijanikin, Lagos, Nigeria

James Ayuk

Risk & Vulnerability Science
Centre, University of Fort Hare,
1 King Williams Town Road,
Private Bag X1314, Alice,
South Africa

The effects of detergent and fertilizer toxicity on growth rate, nutrient utilization and survival of *Clarias gariiepinus* (Burchell) fingerlings reared under controlled condition

Ayofe M Hamed, Albert O Amosu and James Ayuk

Abstract

Study on the effects of metals on growth rate and survival of *Clarias gariiepinus* (Burchell 1822) fingerlings bred in tanks containing water concentrated with inorganic fertilizer and detergent was conducted. 300 fingerlings of *C. gariiepinus* (7.5 ± 0.24) were stocked into different tanks of 5000m³ capacity water and fed with formulated diet for 8 weeks. The tanks were treated with 0ml, 50ml, 100ml, 150ml, 200ml and 250 ml of inorganic detergent and fertilizer respectively. The metal content in the fish was determined through wet acid digestion and extraction procedure using Atomic Absorption Spectrophotometer, while growth was determined using indices method. The values obtained for Cu, Zn, Mg, P, Cd, Pb, Mn, K and Na respectively were generally below the Federal Environmental Protection Agency (FEPA) limits. The survival rate recorded for fingerlings treated with fertilizer was 99% for all concentrations but a range between 88% to 0% survival was recorded for fingerlings stocked in detergent treated water. Toxicity behavior response observed in this study were, slow movement, gulping for atmospheric oxygen, darting and erratic swimming. The results obtained indicated that concentration of 50ml detergent and 100ml of fertilizer were the suitable and tolerable limits in terms of growth and survival for culturing *C. gariiepinus* fingerlings.

Keywords: Growth rate, *C. gariiepinus*, concentration, detergent, fertilizer, fingerlings, metals, survival

1. Introduction

Globally, most coastal nations are undergoing a rapid industrial revolution ^[1, 2], these developmental changes may lead to several environmental deterioration such as land, air and aquatic pollution ^[3, 4]. Aquatic pollution and over exploitation of coastal fish stock are detrimental to sustainable fish production. Therefore, an alternative source of fish production through aquaculture for a sustainable animal protein ^[5], best available cultural practices, bodes well for the aquaculture industry and ever increasing population ^[6]. Yet these aquatic resources provides livelihoods for about 3 billion people and about 10–12% of the world population engaged in the fisheries and aquaculture ^[7-9]. The population of the world is about 7.5 billion and requires renewable resources for sustainable existence. Of which over 1.3 billion people still suffer from hunger, this comprises the poor, especially those in rural areas, who lives below one dollar and 150g (50–60%) of protein required of adult per day) of protein per day ^[10]. Although 53% marine fishery resources of most fishing territories and water bodies have reached their maximum potential for fisheries production, while 32% is overfished, depleted, or recovering from depletion ^[11-14]. This is due to the fact that several millions of people around the world derived a source of income and livelihood in the coast and fisheries sector. This fish stock recruitment problem is characterized with increase in capture of large exotic species, negative aquaculture practices and other detrimental effects like pollution/eutrophication, climate change on fisheries production ^[15].

Aquaculture remains one of the fastest-growing agro-industrial activities in the last four decades and projected to outpace population growth. As the world population increases geometrically, there is a consequent demand for more food production, employment and goods and services ^[16]. These gave rise to an increase in industrialization, which has resulted into the release of more waste materials of various kinds into the immediate environment, especially

Corresponding Author:

Ayofe M Hamed

Department of Fisheries,
Faculty of Science, Lagos State
University. P.O. Box 0001,
LASU Post Office, Ojo, Lagos,
Nigeria

aquatic environment [17]. The problems of bioaccumulation of heavy metals in aquatic organisms, which ultimately serve as food to man therefore, deserve an urgent attention. In Africa generally and Nigeria in particular, it has been found that there is unregulated discharge of effluents [18] from agro-industrial activities and communities into water [19, 20]. Concern has been expressed in some circles about the role and fate of the trace metals derived from these sources [21].

Many causes of pollution including fertilizer contain nutrients such as nitrates and phosphates [22]. In excess levels, nutrients over stimulate the growth of aquatic renewable resources. However, excessive growth of these types of organism consequently clogs the water ways, use un-dissolved oxygen as they decompose, in turn, and prove very harmful to aquatic organism as it affects the respiration ability of fish [23]. On the other hand metallic pollution may be acute sources of pollution not just because they are toxic above relatively low concentration [24], but because they are persistent in the environments long after the primary source of pollutant has been removed [35]. Effluents from industries and agro-allied facilities have been identified as one of those introducing heavy metals along with other pollutants into the Nigeria aquatic environment [24, 26]. In addition, the effect of pollutants

at the cellular level can accumulate and eventually alter the functional integrity of an organ or tissue system, which potentially affect life [27].

Due to the economic value and resistance nature of *C. gariepinus* as a potential culturable fish species in Nigeria, the aims and objectives of this research was to investigate the best conditions for cultivating *C. gariepinus*. Also to determine the effects of inorganic fertilizer and detergent as pollutants in the culture medium with resultant effects on growth performance and survival of *Clarias gariepinus* fingerlings.

2. Materials and Methods

2.1 Experimental fish samples

A total of three hundred (300) fingerlings were obtained from the Aquaculture/fisheries unit of the Agricultural Science department, Adeniran Ogunsanya College of Education, Oto/Ijanikin, Lagos State, and transported in aerated plastic bags to the *In-situ* hatchery facilities of Fisheries department, Lagos State University, Ojo, Lagos for the experiment (Figure 1). The weight of the fingerlings collected ranged between average mean weights of $7.5 \pm 0.24\text{g}$, and were four weeks old at the time of collection. They were acclimatized for 24 hours prior to the commencement of the experiment.




Fig 1: Locations of the fish collection site and *In-situ* experimental research facility.

2.2 Experimental Design

The fingerlings were grouped into different tanks with 5000m^3 of water, same age group were used in each of three treatments {(Control (C)_{1,2,3}, Fertilizer (F)_{1,2,3} and Detergent, (D)_{1,2,3}} and three replicates were weighed and stocked in 5000m^3 culturing bowls. 50g each of fertilizer and detergent were separately weighed and dissolved in 5 litres of water and allowed to stay for 48 hours to dissolve completely. The solution of the fertilizer (F) and detergent (D) were measured into the different tanks labeled as F1_{1,2,3} to F5_{1,2,3} and D1_{1,2,3} to D5_{1,2,3} while F₀ and D₀ are control for fertilizer and detergent respectively. The fishes were introduced into various already prepared concentrations of fertilizer and detergent (0ml, 50ml, 100ml, 150ml, 200ml and 250ml) after 24 hours acclimatization.

2.3 Feed Formulation & Determination of Growth and Nutrient Utilization

Fish feed of 30% crude protein level (fish meal, soybean meal, white maize, palm oil, Ca_2SO_4 and premix) was formulated using the Pearson's square method as described in [28] and reported by [23], while the preparations of the diet were based on the description by [29]. The feed was administered twice (per day at 08:30 and 17:30 hr) daily at 4% body weight for 8 weeks. Weight changes were measured every three days and the feeding rate adjusted accordingly to accommodate the change [30]. The growth rate was carried out using growth indices method as described by [31, 32].

2.4 Physicochemical Parameters and Chemical Evaluation of Experimental fish

Samples of experimental diets and fingerlings were analyzed for proximate composition, according to the methods of [33]. Water quality indicators were monitored closely by determining pH, temperature, dissolved oxygen and total ammonia levels in the culture medium as described by [34, 35].

2.5 Survival

Percentage mortality was calculated using this formula

$$\% \text{Survivability} = \frac{\text{Number of fish stocked} - \text{Number of mortality} \times 10}{\text{Number of fish stocked}}$$

2.6 Determination of wet acid digestion of P, Na, Mg, Cd, Mn, Pb, Zn and Cu

0.5g of grinded fish sample material (oven dry 60 °C) was weighed into a 125ml Erlenmeyer flask which has been previously washed with distilled water. Perchloric acid of about 4ml, 25ml conc. HNO₃ and 2ml conc. H₂SO₄ was added under a fume hood. The contents were mixed and heated gently at low to medium heat on a hot plate under perchloric acid fume hood. The sample was allowed to cool and 40-50ml of distilled water was added. Boil for another half a minute on the same plate at medium heat. The sample was cooled and the solution is filter completely with a wash bottle into a

100ml Pyrex volumetric flask. The solution was stored P and Fe, determinations of Zn, Na by calorimetric and Mg was determined by atomic absorption spectrometry (AAS) [36].

2.7 Statistical analysis

All data collected were analyzed for significant differences ($p < 0.05$) (ANOVA) on Graph Pad Prism 5. The results were expressed as mean $\bar{X} \pm SD$ (standard deviation) and pictorial representation. Determined differences among treatments were partitioned by the Least Significant Difference (LSD) and the Duncan's New Multiple Range Test (DNMRT) [37].

3. Results

Proximate and gross compositions of the diet with different ingredients used are presented in Table 1, weight gain of fish treated in detergent and fertilizer were showed in Table 2 and 3 respectively. While metals concentration in both fish treated detergent and fertilizer were presented in Table 4 and 5. The water quality parameters of both culture medium detergent and fertilizer are showed in Table 6 and 7 respectively. The percentage survival of fish and toxicity behavior respond in detergent solution are presented in Table 8. Table 1: Shows the proximate and gross composition of feed ingredient formulated at 30% crude protein, with the gross composition of the diet is high for carbohydrate, with soybean and fishmeal contributed 22.4% each of the diet.

Table 1: % Proximate and gross composition of feed ingredients

Proximate composition 30% C.P		Gross composition of feed ingredient 30% C.P	
Moisture content	19.30	Fishmeal	22.40
Dry matter content	80.87	Soybean	22.40
Fat content	4.04	Maize	48.60
Ash content	9.53	Vitamin Premix	1.00
Crude fibre content	10.32	Oil	5.00
Crude protein content	7.62	Dicalcium phosphate	0.50
Nitrogen free extract	49.36		

The summary of the growth in Table 3: Shows the biweekly weight gain of fish in detergent tank. All concentration and control are not different in initial weight but the final weight

gain are significantly different with D1 having highest weight gain. No weight gain recorded for D4 and D5 beyond week 2 due to death of the fish.

Table 2: Biweekly weight gain (WTG) of fish treated with detergent

Days	Concentrations of detergent (ml)					
	Control	50	100	150	200	250
14	0.35 ± 0.30 ^a	0.25 ± 0.07 ^b	0.30 ± 0.14 ^b	0.30 ± 0.14 ^b	0.35 ± 0.07 ^c	0.45 ± 0.07 ^d
28	0.40 ± 0.00 ^a	0.10 ± 0.00 ^b	0.10 ± 0.00 ^b	0.40 ± 0.28 ^c	-	-
42	0.85 ± 0.49 ^a	0.80 ± 0.14 ^b	0.70 ± 0.14 ^c	0.55 ± 0.21 ^c	-	-
56	1.10 ± 0.14 ^a	1.25 ± 0.90 ^b	0.85 ± 0.00 ^b	-0.25 ± 0.21 ^c	-	-

Figures in the same horizontal row having the same superscript are not significant different ($p > 0.05$)

Table 3: Biweekly weight gain (WTG) of Fish treated with Fertilizer

Days	Concentrations of in-organic fertilizer (ml)					
	Control	50	100	150	200	250
14	0.30 ± 0.14 ^a	0.30 ± 0.00 ^a	0.30 ± 0.14 ^a	0.25 ± 0.07 ^b	0.20 ± 0.00 ^b	0.35 ± 0.07 ^c
28	0.30 ± 0.14 ^a	0.35 ± 0.21 ^b	0.30 ± 0.28 ^a	0.60 ± 0.14 ^c	0.45 ± 0.35 ^c	0.35 ± 0.21 ^b
42	1.05 ± 0.78 ^a	0.80 ± 0.56 ^b	0.80 ± 0.48 ^b	0.80 ± 0.57 ^b	0.65 ± 0.49 ^c	0.70 ± 0.50 ^c
56	1.20 ± 0.42 ^a	0.85 ± 0.21 ^b	1.50 ± 0.70 ^c	1.45 ± 0.07 ^c	1.30 ± 0.57 ^d	1.25 ± 0.35 ^d

Figures in the same horizontal row having the same superscript are not significant different ($p > 0.05$)

Table 4: Concentration of metals in *Clarias gariepinus* treated with detergent

Conc. (ml)	Metals (mg.g ⁻¹)								
	Na	K	P	Mg	Cu	Zn	Cd	Pb	Mn
Control	4.08±0.01	6.05±0.07	4.57±0.07	3.24±0.20	0.08±0.05	0.87±0.01	0.04±0.03	2.21±0.02	0.13±0.07
50	3.82±0.03	5.61±0.01	3.68±0.05	2.88±0.01	0.06±0.01	0.83±0.04	0.05±0.02	2.09±0.04	0.13±0.03
100	1.05±0.04	1.71±0.01	3.47±0.01	3.77±0.07	0.07±0.00	0.75±0.02	0.04±0.01	2.29±0.03	0.12±0.02
150	4.24±0.01	6.41±0.40	4.2±0.01	3.62±0.04	0.06±0.01	0.93±0.01	0.04±0.01	2.46±0.01	0.13±0.00
200	3.11±0.07	5.40±0.14	4.90±0.12	3.72±0.02	0.08±0.02	1.08±0.01	0.05±0.01	2.19±0.02	0.14±0.01
250	2.85±0.02	5.20±0.07	5.0±0.06	3.98±0.06	0.07±0.00	0.88±0.05	0.03±0.01	1.94±0.01	0.11±0.02

Table above shows the summary of heavy metals, Cu, Zn, Cd, Pb, Mn and nutrient, Na, P, K in *Clarias gariepinus* cultured in detergent solution with K having the highest value $6.41 \pm 0.14 \text{ mg.g}^{-1}$ while Cd has the lowest value $0.03 \pm 0.01 \text{ mg.g}^{-1}$.

Table 5: Concentration of metals (mg.g⁻¹) in *Clarias gariepinus* treated with fertilizer

Conc.(ml)	Na	K	P	Mg	Cu	Zn	Cd	Pb	Mn
50	4.03±0.07	5.95±0.07	4.72±0.02	3.11±0.07	0.07±0.02	0.95±0.02	0.04±0.01	2.00±0.01	0.12±0.02
100	4.17±0.01	6.1±0.14	5.06±0.01	3.85±0.07	0.07±0.07	0.93±0.01	0.03±0.00	2.50±0.01	0.14±0.01
150	3.95±0.01	5.61±0.05	3.89±0.02	3.86±0.00	0.07±0.01	0.88±0.03	0.04±0.00	2.79±0.03	0.12±0.01
200	4.01±0.01	5.7±0.14	5.32±0.03	3.87±0.01	0.08±0.01	0.93±0.02	0.04±0.01	2.44±0.04	0.14±0.02
250	3.81±0.01	5.3±0.12	4.88±0.03	3.78±0.03	0.06±0.02	0.83±0.01	0.04±0.01	2.11±0.02	0.12±0.01

Table shows the summary of heavy metals, Cu, Zn, Cd, Pb, Mn and nutrient, Na, P, K in *Clarias gariepinus* cultured in fertilizer solution with K having the highest value $6.1 \pm 0.14 \text{ mg.g}^{-1}$ while Cd has the lowest value $0.03 \pm 0.00 \text{ mg.g}^{-1}$.

Table 6: Water quality parameters of culture medium with detergent solution

Concentration (ml)	D ₀ control 0 ml	D ₁ 50 ml	D ₂ 100 ml	D ₃ 150 ml	D ₄ 200 ml	D ₅ 250 ml
Temperature °C	26.10 ± 0.07 ^a	26.20 ± 0.31 ^a	26.4 ± 0.12 ^a	25.6 ± 1.03 ^a	26.0 ± 0.05 ^b	24.4 ± 1.93 ^c
pH	7.1 ± 0.1 ^a	6.9 ± 0.1 ^b	6.9 ± 0.1 ^b	7.0 ± 0.1 ^b	7.0 ± 0.1 ^b	7.1 ± 0.1 ^b
Salinity %	0.2 ± 0.0 ^a	0.2 ± 0.0 ^b	0.2 ± 0.0 ^b	0.2 ± 0.0 ^b	0.2 ± 0.0 ^b	0.2 ± 0.0 ^b
(CO ₂) mg.l ⁻¹	3.0 ± 1.7 ^a	1.6 ± 0.1 ^b	1.1 ± 0.2 ^b	0.8 ± 0.4 ^b	0.8 ± 0.6 ^b	1.4 ± 0.6 ^b
(NH ₃) mg.l ⁻¹	0.10 ± 0.11 ^a	0.66 ± 0.09 ^b	0.45 ± 0.09 ^b	0.31 ± 0.12 ^b	0.45 ± 0.09 ^b	0.49 ± 0.04 ^b
(DO) mg.l ⁻¹	4.00 ± 0.4 ^a	1.8 ± 0.0 ^b	1.8 ± 0.8 ^b	0.5 ± 0.3 ^c	1.6 ± 0.6 ^b	0.4 ± 0.1 ^c

Values in the same row having the same superscript are not significantly different ($p > 0.05$).

Table 7 shows the values of water quality parameters of detergent solution, all the value of each parameters fall within the range set by FEPA except the values for DO which is lower and NH₃ which is high.

Table 7: Water quality parameters of culture medium with fertilizer solution

Concentration	F ₀ Control 0ml	F ₁ 50ml	F ₂ 100ml	F ₃ 150ml	F ₄ 200ml	F ₅ 250ml
Temperature °C	26.1 ± 1.93 ^a	25.2 ± 0.10 ^a	25.1 ± 1.13 ^a	25.1 ± 0.15 ^b	26.1 ± 0.08 ^b	27.1 ± 1.13 ^b
pH	7.0 ± 0.1 ^a	6.9 ± 0.1 ^b	7.0 ± 0.1 ^b	6.9 ± 0.5 ^b	6.9 ± 0.0 ^b	6.9 ± 0.0 ^b
Salinity %	0.2 ± 0.0 ^a	0.2 ± 0.0 ^a	0.2 ± 0.0 ^a	0.2 ± 0.0 ^b	0.2 ± 0.0 ^b	0.2 ± 0.0 ^b
(CO ₂) mg.l ⁻¹	3.0 ± 1.7 ^a	2.1 ± 1.4 ^b	3.1 ± 1.7 ^b	2.5 ± 1.3 ^b	2.7 ± 1.2 ^b	2.9 ± 1.3 ^b
(NH ₃) mg.l ⁻¹	0.10 ± 0.12 ^a	0.66 ± 0.55 ^b	0.69 ± 0.23 ^b	0.71 ± 0.06 ^b	0.67 ± 0.12 ^b	0.73 ± 0.04 ^b
(DO) mg.l ⁻¹	4.00 ± 0.10 ^a	0.70 ± 0.00 ^b	0.65 ± 0.5 ^b	2.85 ± 1.2 ^c	2.45 ± 1.8 ^c	0.90 ± 0.1 ^b

Values in the same row having the same superscript are not significantly different ($p > 0.05$).

Table 8 shows the values of water quality parameters of fertilizer solution, all the value of each parameters fall within the range set by FEPA except the values for DO which is lower and NH₃ which is high.

Table 8: Percentage survival rate of fish and toxicity behavior response in detergent solution

	Concentrations					
	D ₀ control 0ml	D ₁ 50ml	D ₂ 100ml	D ₃ 150ml	D ₄ 200ml	D ₅ 250ml
Percentage Toxicity	100%	80%	28%	16%	0%	0%
Fish Behavior Response	Normal	Slow movement	Reduce feed intake and gulping for atmospheric oxygen	Gulping, Stop feeding and erratic swimming	Stop feeding, darting and erratic swimming	Stop feeding, darting and erratic swimming

4. Discussion

As illustrated in Table 2 shows the final body weight gain of *Clarias gariepinus* treated with detergent. It indicates that the feed was utilized for body growth, but as the concentration of

the detergent increases, survival rate reduced. The final weight gain of fish treated with fertilizer in Table 3 shows that the feed were fully utilized for body growth. This is in line with the observation of [38] who reported that the

fertilization of water allows for adequate phytoplankton production which enhances a better growth and survival of the fish.

Experimental investigations revealed the presence of zinc (Zn), Lead (Pb), Copper (Cu), Cadmium (Cd), and minerals such as; Sodium (Na), Potassium (K), Magnesium (Mg), Phosphorus (P) in the samples analyzed. The values obtained are quite lower when compared to those obtained by the Federal Environmental protection agency [39]. Table 4 and 5 shows the mean value of Zn varies between 0.75 mg.g⁻¹ and 1.08 mg.g⁻¹ in both pollutants. The value was by far higher than the reported value of 0.02 mg.g⁻¹ from rivers of South Carolina [40] but lower than those of Kaduna River, Nigeria [41, 42] posited that zinc has low toxicity to man, but relatively high toxicity to fish. The appreciable levels recorded in this study would therefore be a serious cause for concern. The mean levels of Cd (0.03-0.05 mg.g⁻¹) recorded in this study was low. However, the values was higher than the reported values of <0.01 mg.g⁻¹ in rivers of South Carolina [40]. The highest Cd level recorded is an indication of anthropogenic input [43]. Cadmium has been known to have contributed to pulmonary disease, reduced glucose tolerance, severe kidney and liver damage and death in human beings [44, 45]. Also the mean levels of Pb varied between 1.94 mg.g⁻¹ and 2.50 mg.g⁻¹ in the catfish samples and these values recorded were higher than the reported mean value of 0.56 mg/g from Kaduna River, Nigeria [41]. Lead is known as a deadly and cumulative poison even when consumed in small quantities and is capable of deadening nerve receptors in man [46, 47]. The mean levels of Mn ranges from 0.11 - 0.14 mg.g⁻¹ in the catfish sampled and are below the range set by [39]. The range value of copper (0.064 - 0.081 mg.g⁻¹) recorded in sampled fish were below the value (0.11-11.9 mg.g⁻¹) observed by [48] when he tested the toxicity of this metal on some pelagic and benthic fish species. However, copper levels were lower than the other metals investigated in this study. Potassium levels in fish bred in fertilizer is high compare to fish in detergent. This simply explain that the level of potassium in the environment of greatly affect the level of potassium in the fish, this is in line with the findings of [49] who opined that fishes are susceptible to physical and chemical in the environment and may reflect in blood composition. Phosphorus level in both pollutants varies differently; the variation of phosphorus was explained by [50] who gave knowledge on role of key nutrient in pathogenesis of skeletal fish with an emphasis on mineral. There was no significant difference ($p > 0.05$) in the mean values of all the physico-chemical parameters in the treatments; they fall within the acceptable limits, except the values for ammonia and dissolved oxygen. [51] stated that the amount of oxygen in water is not as constant as in air but fluctuates markedly depending on depth. The mean dissolved oxygen values obtained in both treatments ranges from 0.4 ± 0.1 - 2.8 ± 1.2, these values are relatively lower than the range considered desirable for fish survival at 30°C [52]. The values of the pH is good for fish growth and survival as this falls within the range considered good for fish survival by [52]. The temperature and salinity values obtained in this study are characteristic of a tropical freshwater ecosystem [53]. The values of carbon dioxide are within the [39] guidelines standard for this parameter. But the ammonia was relatively higher than the limits set by [39]. The above may imply an increased toxicity with the raised values of the physico-chemical parameters. [43] stated that toxicity of pollutant of fish usually increased with temperature which may be due to an increased

uptake of toxin which added environmental stress e.g. reduced oxygen solubility [54]. The recorded values of water temperature were within tolerable range for the fish species and may not have affected the toxicity of the treatments. Ammonia (NH₃) appears to be the major source of toxicity in the experiment to which *Clarias gariepinus* was exposed. This is in line with the observations of [55] who indicated ammonia as the main cause of toxicity of pollutant from industrial plant and where death may occur in aquatic organisms.

Difference in mortality of *Clarias gariepinus* may be due to the differential toxicity of the stock solutions. As it was observed the test solution of detergent concentrate at 200ml and 250ml survival was at a quick decline as none of the fish survive, mean while the fishes increased their activity to cope with rise in level of concentration. This is similar to [56] that hyperactivity wave down with the depletion of fish energy eventually causing their death. However, the results indicate that survival time for fishes in the detergent test solution declined with increase in concentration and time of exposure. But no mortality was observed for the fertilizer test solution in this experiment because the fish were able to utilize the minerals from the fertilizer for their survivability, since fertilizers are known to boost pond phytoplankton use as natural food for fish. Meanwhile continued discharged of these effluents (i.e. detergent and fertilizer) over a long period could lead to uptake and bioaccumulation in the aquatic ecosystem.

5. Conclusion

It is concluded that 50ml of detergent and 100ml of inorganic fertilizer are the safe concentration for *Clarias gariepinus* under cultured environment. Thus it is recommended that: (i) Monitoring status of the water body for all seasons to ascertain its effect on the fisheries must be ensured by the concerned Federal and State Agencies. (ii) Environmental impact assessment and auditing should be carried out on all the industries found along the water body, and (iii) Installation of recycling facilities for aquatic waste, with stringent penalties on defaulting industries.

6. Acknowledgements

The authors are grateful to the Department of fisheries, Faculty of Science, Lagos State University, Ojo, Lagos, Nigeria. We also appreciate the contribution and efforts of Mr. Adeniyi Togunde (Farm manager) and Mr. Lamina Arubiewe (Lab technologist) at the Department of Agricultural Science, Adeniran Ogunsanya College of Education, Lagos, Nigeria.

7. References

1. Bowen A, Duffy C, Fankhauser S. 'Green Growth and The New Industrial Revolution', Policy brief, London School of Economics and Political Science. 2016, 25
2. Luodan Cao, Jialin Li, Mengyao Ye, Ruiliang Pu, Yongchao Liu, Qiangong Guo. Baixiang Feng and Xiayun Song Changes of Ecosystem Service Value in a Coastal Zone of Zhejiang Province, China, during Rapid Urbanization. Int. J Environ Res Public Health. 2018; 15(7):1301. doi: 10.3390/ijerph15071301
3. Keizire BB, Mugenyi O. Mainstreaming environment and natural resource issues in selected government sectors status, considerations and recommendations. ACODE, Kampala, Uganda, 2006.

4. Kapur R. Natural Resources and Environmental Issues. J Ecosys. Ecograph. 2016; 6:196. doi:10.4172/2157-7625.1000196G
5. Jennings S, Kaiser MJ, Reynolds JD. Marine Fisheries Ecology. Blackwell Science, Oxford, United Kingdom, 2001, 310.
6. Amosu AO, Hammed AM, Togunde KA, Joseph OO, Adekoya A. Possible Aquaculture Development in Nigeria: Evidence for Commercial Prospects. Journal of Agricultural Science and Technology B 7 2017; 194-205 doi: 10.17265/2161-6264/2017.03.007
7. FAO. The state of world fisheries and aquaculture 2008. FAO fisheries and aquaculture department. Food and Agriculture Organization of United Nations. Rome, 2009.
8. Smith MD, Roheim CA, Crowder LB, Halpern BS, Turnipseed M, Anderson JL *et al.* Sustainability and Global Seafood, Science. 2010; 327:784-786
9. FAO Fisheries Department, Fishery Information, Data and Statistics Unit. FishStatPlus. Universal Software for fishery statistical time series. Version 2.3 in 2000. 2012. Retrieved from <http://www.fao.org/docrep/013/i1820e/i1820e00>
10. World Bank. World Development Report 2008: Agriculture for Development. World Bank, Washington, DC. 2008.
11. Garcia SA, Rosenberg AA. Food security and marine capture fisheries: characteristics, trends, drivers and future perspectives. Philos Trans R Soc Lond B Biol Sci. 2010; 365(1554):2869-2880.
12. FAO. The State of World Fisheries and Aquaculture 2002 - SOFIA, Rome, Italy, 2002.
13. Clover C. End of the Line: How overfishing is changing the world and what we eat. Ebury Press, London. 2004. ISBN 0-09-189780-7
14. FAO. The State of World Fisheries and Aquaculture - Food and Agriculture Organization of the United Nations (FAO), Fisheries and Aquaculture Department. Rome. Italy, 2012.
15. Garcia SA, Rice JC. Fisheries, food security, climate change, and biodiversity: Characteristics of the sector and perspectives on emerging issues. ICES Journal of Marine Science. 2011; 68(6):1343-1353.
16. World Development Report. The Changing Nature of Work. Washington, DC: World Bank. 2019 doi:10.1596/978-1-4648-1328-3.
17. FAO. Inland Water Resources and Aquaculture Service. FAO Fishery Resources and Environment Division. FAO Fisheries Technical Paper 328. Food and Agriculture Organization of the United Nations. Rome, 1995.
18. Ipeaiyeda AR, Onianwa PC. Monitoring and assessment of sediment contamination with toxic heavy metals: case study of industrial effluent dispersion in Alaro River, Nigeria. Appl Water Sci. 2018; 8:161. <https://doi.org/10.1007/s13201-018-0815-6>
19. Edokpayi JN, Odiyo JO, Durowoju OS. Impact of Wastewater on Surface Water Quality in Developing Countries: A Case Study of South Africa, Water Quality, Hlanganani Tutu, Intech Open, 2017. DOI: 10.5772/66561. Retrieved from: <https://www.intechopen.com/books/water-quality/impact-of-wastewater-on-surface-water-quality-in-developing-countries-a-case-study-of-south-africa>
20. Alayu E, Yirgu Z. Advanced technologies for the treatment of wastewaters from agro-processing industries and cogeneration of by-products: a case of slaughterhouse, dairy and beverage industries. Int. J. Environ. Sci. Technol. 2018; 15:1581-1596. <https://doi.org/10.1007/s13762-017-1522-9>
21. Olowu RA, Jimoh AA, Onwordi CT, Onipede JO, Ogunnaike BA, Amosu AO. Potential Toxic Metals Concentrations in Different Aquatic Lives Harvested from Lagos Island and Epe Lagoons. Current World Environment. 2015; 5(2):80-89
22. Nürnberg GK. "Attempted management of cyanobacteria by Phoslock (Lanthanum-modified clay) in Canadian lakes: Water quality results and predictions". Lake and Reservoir Management. 2017; 33(2):163-170.
23. Hammed AM, Amosu AO, Fashina-Bombata HA. Effect of partial and total replacement of Soybean meal with Pigeon pea (*Cajanus cajan*) as alternative plant protein source in the diet of juveniles African Mud catfish (*Clarias gariepinus* (Burchell, 1822)). International Journal of Food Technology Photon, 2013; 105:139-145.
24. Authman MMN, Zaki MS, Khallaf EA, Abbas HH. Use of Fish as Bio-indicator of the Effects of Heavy Metals Pollution. J Aquac Res Development 2015; 6:328. doi:10.4172/2155-9546.1000328
25. Amosu AO, Robertson-Andersson DV, Kean E, Maneveldt GW, Cyster L. Biofiltering and uptake of dissolved nutrients by *Ulva armoricana* (Chlorophyta) in a land-based aquaculture system. International Journal of Agriculture and Biology 2016; 18:298-304.
26. Anifowose AJ, Oyeboode AW. Studies on heavy metals contents of Osun River at the pre-urban settlement and across Osogbo City, Nigeria, Journal of Taibah University for Science, 2019; 13(1):318-323, DOI: 10.1080/16583655.2019.1567899
27. Etori OS, Iyama WA, Amadi MC. Status of heavy metals contamination in water from the Elelenwo River, Obio-Akpor, Rivers State, Nigeria . Direct Research Journal of Chemistry and Material Science. 2019; 6(3):25-31.
28. Wagner J, Stanton TL. Formulating Rations with the Pearson Square. No. 1.618. Colorado State University Extension. Livestock Series Management, 2006; 618:1-3. Retrieved from <http://www.ext.colostate.edu/PUBS/livestk/01618.html>.
29. Royes J, Chapman FA. Preparing Your Own Fish Feeds , Department of Fisheries and Aquatic Sciences, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL 32611. Document No. Cir 97. 2012. Retrieved from <http://www.edis.ifas.ufl.edu>.
30. Hogendoorn H, koops WJ. Growth and production of African catfish *Clarias lazear*. Effects of stocking density, pond size and mix culture with tilapia under extensive field condition. Aquaculture, 1983; 34(3, 4):253-263.
31. Hammed AM, Amosu AO, Fashina-Bombata HA. Effect of partial and total replacement of Soybean meal with Pigeon pea (*Cajanus cajan*) as alternative plant protein source in the diet of juveniles African Mud catfish (*Clarias gariepinus* (Burchell, 1822)). International Journal of Food Technology Photon, 2013; 105:139-145.
32. Hammed AM. Aspects of Biology, Biochemical Characterization and Nutrient Requirements of a Cichlid species 'Wesafu' from Epe Lagoon, Lagos, Nigeria. Ph.D Thesis Dissertation, Lagos State University, Nigeria, 2012.

33. AOAC. Association of Official Analytical Chemists. Official Methods of Analysis. 16th edn, ed. p. cuniff. AOAC, International, Arlington, Virginia, U.S.A, 1990.
34. Boyd CE. Pond Aquaculture Water Quality Management. Aquaculture Series (Chapman & Hall Aquaculture Series). ed. by C.S Tucker. Springer. 1998, 700.
35. Boyd CE. Water Quality Standard: Dissolved Oxygen. Global Aquaculture Advocate. 2001; 4(6):70-72.
36. AOAC. Association official methods of analysis. Association of Official Analytical Chemists. Arlington, Virginia, 2005.
37. Duncan DB. Multiple Range and Fultple F Test. Biometric. 1955; 11:1-42.
38. Martínez Cruz P, Ibáñez AL, Monroy Herмосillo OA, Ramírez Saad HC. Use of probiotics in aquaculture. ISRN microbiology, 2012; 916845. <https://doi.org/10.5402/2012/916845>
39. Federal Environmental Protection Agency, FEPA. Guidelines and standards for environmental pollution control in Nigeria. FEPA, Lagos, Nigeria. 1991, 238.
40. Koli AK, Cauty WT, Felix KI, Reed RJ, Whitmore R. Trace metals in some fish species of South Carolina. Bull Env. Cont. Toxic. 1978; 20(3):328-331.
41. Nwaedozi JM. The determination of heavy metal pollution in some fish samples from River Kaduna. J. Chem. Soc. Nigeria, 1998; 23:21-23.
42. Alabaster JS, Lloyd R. Water quality criteria for fish, 2nd. Ed., London, Butterworths, 1980.
43. Ali H, Khan E, Ilahi I. Environmental Chemistry and Ecotoxicology of Hazardous Heavy Metals: Environmental Persistence, Toxicity, and Bioaccumulation. Journal of Chemistry. 2019, 14 |Article ID 6730305 <https://doi.org/10.1155/2019/6730305>
44. Flora SJS, Mittal M, Mehta A. Heavy metal induced oxidative stress and its reversal by chelation. Ind. J. Med. Res. 2008; 128:501-523.
45. An AT, Ali A, Haq QMR. Glutathione as an antioxidant in inorganic mercury induced nephrotoxicity. J. Postgrad. Med. 2011; 57:72-77.
46. Guedenon P, Edozh AP, Kaki C, Yehouenou APE, Gnanđi K, Montcho S, Hounkpatin A *et al.* Arsenic, Cadmium, Copper and Lead Accumulation in Water, Sediments and FishSpecies of Oueme River in Bonou . British Journal of Pharmacology and Toxicology. 2012; 3(1):13-20.
47. Agbozu IE, Ekweozor IKE, Opuene K. Survey of heavy metals in the catfish *Synodontis clarias*. Int. J. Environ. Sci. Tech., 2007; 4 (1): 93-97.
48. Oyewo O. Distribution of Cr, Pb, Cd, Zn, Fe and Mn in lake Victoria Sediments East Africa. Bull. Environ Contain. Toxicol, 1998; 42:807-813.
49. Wilson RW, Taylor EW. The physiological responses of freshwater rainbow trout, *Oncorhynchus mykiss*, during acutely lethal copper exposure. J Comp Physiol B. 1993; 163:38-47.
50. Santosh P, Lall L, Lewis – McCrea M. Role of nutrients in Skeletal Metabolism and Pathology of fish. National Research Council of Canada, 2007.
51. Aniyikaiye TE, Oluseyi TO, Odiyo JO, Edokpayi JN. Physico-Chemical Analysis of Wastewater Discharge from Selected Paint Industries in Lagos, Nigeria, Int J Environ Res Public Health. 2019; 16(7):1235. doi: 10.3390/ijerph16071235
52. Body CE. Water quality In: Warm water fish ponds. Auburn University Agr. Exp. Station Alabama 1981, 358.
53. Adeniyi IF. The ecology of the bonny estuary chemical composition of the rivers and creeks. Paper presented at man and Biosphere Workshop, Port-Harcourt 1986, 30.
54. Minnesota Pollution Control Agency. Low Dissolved Oxygen in Water Causes, Impact on Aquatic Life – An Overview. Water Quality/Impaired Waters 2009; 3(24). Retrieved from <https://www.pca.state.mn.us/sites/default/files/wq-iw3-24.pdf>
55. Savci S. An Agricultural Pollutant: Chemical Fertilizer. International Journal of Environmental Science and Development. 2012; 3(1):77-80.
56. Hammed AM, Fashina-Bombata HA, Olowu RA. Growth Response and Survival Rate of *Clarias gariepinus* Fingerlings Exposed to Varying Metal Concentrations. Pacific Journal of Science and Technology. 2003; 14(1):430-438.