

Mpabulungi, L. and Muula, A. S. (2004). Tobacco use among High School Students in Kampala, Uganda: Questionnaire Study. *Croatian Medical Journal*, 45(1), 80-83.

Murray, C. J. L. and Lopez, A. D. (1996). *The Global Burden of Disease*. Cambridge, MA: Harvard School of Public Health.

Murenzi, J. G. (2001). Habitual activity patterns among adolescent learners in Rwanda. Unpublished Master of Science thesis. Physiotherapy department, University of the Western Cape.

National Council for Population and Development (1998). *Kenya Demographic and Health Survey*. Calverton, Maryland, NDPD, CBS and Macro International Inc.

Nawi, N., Stenlund, H., Bonita, R., Hakimi, M., Wall, S. and Weinehall, L. (2006). Preventable risk factors for non-communicable diseases in Indonesia: Prevalence study using the WHO STEPS approach. *Bulletin of the World Health Organization*, 84, 305-313.

Omran, A. R. (1971). The epidemiological transition: A theory of the epidemiology of population change. *Milbank Memorial Quarterly*, 49, 51-91.



Pampel, F. C. (2005). Patterns of Tobacco Use in the Early Epidemic Stages: Malawi and Zambia, 2000-2002. *American Journal of Public Health*, 95(6), 1009-1015.

Pan American Health Organization (2005).

<http://www.paho.org/common/Display.asp?Lang=S&RECID=7157>, accessed 14, July 2009.

Pan, Z. and Hu, D. (2008). Multi-level analysis of individual and community predictors of smoking prevalence and frequency in China: 1991-2004. *Journal of Public Health Policy*, 29(1), 72-84.

Perel, P., Casas, J. P. Ortiz, Z. and Miranda, J. J. (2006). Non-communicable diseases and injuries in Latin America and the Caribbean: Time for Action. *Plos Medicine*, 3(9), 1448-1451.

- Phillips, J (2001). Recreational activities of high school learners in the Strand. Unpublished Masters Thesis. University of the Western Cape.
- Popkin, B. M. (1998). The nutrition transition and its health implications in lower-income countries. *Public Health Nutrition*, 1, 5-21.
- Reddy, K. S., Prabhakaran, D., Shah, P. and Shah, B. (2002). Differences in body mass index and waist: hip ratios in north Indian rural and urban populations. *Obesity Reviews*, 3, 197-202.
- Redfern, S. and Norman, I. (1994). Validity through triangulation. *Nursing Researcher* 2: 41-56.
- Sachs, J. D. (2005). The end of poverty. New York, NY, Penguin Press.
- Sallis, J. F., Calfas, K. J., Nichols, J. F., Sarkin, A. J., Marilyn, F. J., Caparosa, S., Thompson, S. and Alcaraz, E. J. (1999). Evaluation of a University Course to Promote Physical Activity: Project GRAD, *Research Quarterly for Exercise and Sport*, 70, 1-10.
- Sankaranarayanan, R. (2002). Effective screening programs for cervical cancer in low- and middle-income developing countries. *Bulletin of the World Health Organization*, 79, 954-962.
- Sen, A. (1998). Non-communicable diseases and achieving the Millennium Development Goals. Available. www.unescap.org/esid/hds.
- Shobhana, R., Rama, R. P., Lavanya, A., Williams, R., Vijay, V. and Ramachandran, A. (2000). Expenditure on health care incurred by diabetic subjects in a developing country - a study from southern India. *Diabetes Research and Clinical Practice*, 48, 37-42.
- Silverman, D. (2000). Doing Qualitative Research: A practical handbook. London Sage Publications.

Steyn, K., Jooste, P. L. and Bourne, L. (1991). Risk factors for coronary heart diseases in the black population of the Cape Peninsula. The BRISK study. *South African Medical Journal*, 79, 480-485.

The World Health Report (1999). Making a difference. Geneva: World Health Organization.

The World Health Report (2002) – Reducing risks, promoting healthy life. Geneva, World Health Organization.

Tumuusime, D. K. and Frantz, J. M. (2006). Influence of previous participation in physical activity on its perceptions among tertiary institution students. *African Journal for Physical, Health Education, Recreation and Dance (AJPHERD)*, 12(3), 287-297.

Unal, B., Critchley, J. L. and Capewell, S. (2005). Modelling the decline in coronary heart disease in England and Wales, 1982-2000: Comparing contribution from primary prevention and secondary prevention. *British Medical Journal*, doi: 10.1136/bmj.38561.633345.8F.

UNESCO/UNICEF/WHO/World Bank (2000). Focusing resources on effective school health: A fresh start to enhancing the equality and equity of education. World Education Forum. Final Report. (<http://portal.unesco.org/>, accessed, 14, July 2009).

Unwin, N., Setel, P., Rashid, S., Mugusi, F., Mbanya, J., Kitange, H., Hayes, L., Edwards, R., Aspray, T. and Alberti, N. (2001). Non-communicable diseases in sub-saharan Africa: Where do they feature in the health research agenda? *Bulletin of the World Health Organization*, 79, 947-953.

Van Walbeek, C. (2002). Recent trends in smoking prevalence in South Africa – Some evidence from AMPS data. *South African Medical Journal*, 92(6), 468 – 472.

Wang, W., Lee, E. T., Fabsitz, R. R., Devereux, R., Best, L., Welty, T. K. and Howard, B. V. (2006). A Longitudinal Study of Hypertension Risk Factors and their Relation to Cardiovascular Disease: The Strong Heart Study. *Hypertension*, 47, 403-409.

WHO (2005a). Preventing Chronic Diseases: A Vital Investment. *World Health Organization Global Report*.

WHO (2005b). WHO STEPS Instrument. Department of Chronic Diseases and Health Promotion. *World Health Organization*. Geneva, Switzerland.

World Health Organization (2001a). Physical Activity. Department of Non-communicable Disease Prevention and Health Promotion. *World Health Organization*: Geneva.

World Health Organization (2002). Reducing Risks: Promoting Healthy Life. *World Health Organization*. Geneva.

World Health Organization (2003). Physical inactivity: Direct and indirect health benefits. Non-communicable diseases and prevention and health promotion. *World Health Organization*, Geneva.

World Health Organization (2004). World Health Survey, Kenya.

World Health Organization (2005c). World Health Survey, Kenya.

World Health Organization (2008). Preventing Chronic Diseases: A vital investment. Available: <http://www.who.int/chp/chronic=disease=report/en/>. Accessed 30 June 2009.

World Health Organization (200b1). Summary. Surveillance of risk factors for noncommunicable diseases. *The WHO STEPwise approach*. Noncommunicable diseases and Mental Health.

World Health Report (1998). Obesity: Preventing and Managing the Global Epidemic. Geneva, Switzerland: *World Health Organization*. WHO/NUT/NCD/98.1.

World Health Report (2004). Building blocks for tobacco control, A handbook. Geneva.

Wu, X. and Stamler, Z. (1996). Changes in average blood pressure and incidence of high blood pressure 1983-1984 to 1987-1988 in four population cohorts in the people's Republic of China. The PRC-USA Cardiovascular and Cardiopulmonary Epidemiology Research Group. *Journal of Hypertension*, 14(11), 1267-1274.

Yamane, T. (1967). *Statistics, An Introductory Analysis*, 2nd Edition, New York: Harper and Row.

Yang, G., Fan, L., Tan, J., Qi, G., Zhang, Y. and Samet, J. M. (1999). Smoking in China: Findings of the 1996 National Prevalence Survey. *Journal of American Medical Association*, 33(4), 478-480.

Yusuf, S., Reddy, S., Ounpuu, S. and Anand, S. (2001). Global burden of cardiovascular diseases. Part 1: General considerations, the epidemiologic transition, risk factors and impact of urbanization. *Clinical cardiology: New Frontiers*, 104, 2746-2753.

Retrieved March 26, 2008, from <http://www.circulationaha.org>.

