

























































## Chapter 2

### Literature Review

---

#### 2.1 Hypertension as a major risk factor for CVD

Hypertension is a major driver of CVD in Africa especially stroke and hypertensive heart disease. It is commonly defined as systolic blood pressure (SBP) above 140 mmHg and/or a diastolic blood pressure (DBP) above 90 mmHg. This level of blood pressure can induce heart disease, congestive heart failure, impaired vision and diabetic nephropathy. It is generally described as a silent killer due to its asymptomatic nature and can therefore go undetected for years, affecting organ systems and shortening life expectancy by 10 – 20 years (Xie and Zhang, 2012). Epidemiologic data demonstrates a strong correlation between blood pressure (BP) and cardiovascular (CV) morbidity and mortality (**Kannel *et al.*, 2003**).

#### 2.2. Treatment and management of Hypertension

Blood pressure is controlled by neural mechanisms carried out by the autonomic nervous system and humoral mechanisms involving substances such as nitric oxide (NO) and endothelin-1 (ET-1) that are released by different cell types (**Larson, Symons, & Jalili, 2012**). Decreased vasodilation, increased vasoconstriction, and greater vascular peripheral resistance characterize hypertension. Treatment for hypertension is a life-long commitment that requires drug therapy in combination with lifestyle changes such as weight reduction if overweight, limitation of alcohol intake and a reduction in salt and fat intake. Treating hypertensive patients with antihypertensive drug therapy provides significant clinical benefits. Epidemiological studies have shown that normalization of blood pressure is associated with a significant reduction in major cardiovascular events. Evidence from large-scaled placebo controlled clinical trials have shown that the increased risks of CV events and death associated with

elevated BP are reduced substantially by antihypertensive therapy (Ettehad *et al.*, 2016). CVD and primarily hypertension can be treated with one or a combination of many therapeutic agents namely angiotensin converting enzyme inhibitors, angiotensin receptor blockers, diuretics, beta blockers, calcium channel blockers, vasodilators and centrally acting agents (Aronow *et al.*, 2012). ACE inhibitors are established as one of the main therapeutic agents for treatment of hypertension. It has favourable metabolic, renal, cardiovascular, and quality-of-life effects as compared with other regimens (Niskanen *et al.*, 2001). ACE as a component of the renin angiotensin aldosterone system plays a key role in the homeostatic mechanism of mammals, by contributing to the maintenance of normal blood arterial pressure, cell function and water-electrolyte balance, (Chen *et al.*, 2009).

### **2.3 Role of Renin Angiotensin Aldosterone system and Angiotensin converting enzyme II in Hypertension**

The renin angiotensin aldosterone system plays an integral role in the control of blood pressure and preservation of hemodynamic stability through the regulation of extracellular fluid volume, sodium balance, cardiac and vascular trophic effects. One of the key components of RAAS is angiotensin II, a main effector molecule and a potent vasoconstrictor that acts as a systemic hormone (endocrine) or as a locally generated factor (paracrine, autocrine). The over activity of RAAS has been associated with the development of atherosclerosis, hypertension, left ventricular hypertrophy and cardiovascular events such as myocardial infarction, stroke, congestive heart failure, and nephrosclerosis (Atlas, 2007).

For the past 20 years, efforts have been focused on pharmacotherapy that attenuates the activity of the renin-angiotensin-aldosterone system (RAAS). Three major classes of drugs that fulfil this purpose are the angiotensin-converting enzyme (ACE) inhibitors,  $\beta$ -blockers, and angiotensin receptor blockers. Each of these therapeutic classes exerts its inhibitory effect in different ways.  $\beta$ -Blockers primarily inhibit renin production, thus dampening the systemic

RAAS. ACE inhibitors inhibit the angiotensin converting enzyme responsible for the cleavage of peptide fragments off angiotensin I to create the active moiety angiotensin II. Angiotensin receptor blockers rather than inhibiting the formation of angiotensin II occupies its high-affinity type 1 (AT<sub>1</sub>) receptor, thus inhibiting its ability to exert biological activity (Weir and Dzau,1999).




Figure 2. 1 The Renin Angiotensin Aldosterone System (Almeida & Coimbra, 2019)

Angiotensin II, the principal effector peptide of the RAAS, has far reaching effects on vascular structure, growth and fibrosis, and is a key regulator of vascular remodelling and inflammation. Besides its vasoconstrictive effects, angiotensin II through a variety of different mechanisms, whether direct or indirect, can substantially affect the structure, function, and atherosclerotic risk of blood vessels. It stimulates growth, inhibits apoptosis, and promotes smooth muscle cell

growth and migration through a variety of different mechanisms, predominantly by causing oxidative stress and altering the redox potential of the blood vessel. Oxidative stress due to angiotensin II and free radical generation has been implicated in hypertension. ACE inhibitors such as captopril are believed to possess antioxidant activity against these factors. **(Malgorzataí *et al.*, 1997)**. The generation of Angiotensin II by ACE promotes platelet aggregation, thrombosis and endothelial dysfunction. All these factors are known to increase vascular tone, promote remodelling and restructuring and augment atherosclerotic risk **(Weir and Dzau, 1999)**.

ACE is a dipeptidyl carboxypeptidase with a zinc atom. It consists of a single polypeptide chain containing two domains: N and C. Each domain consists of two catalytic sites. The lung capillaries boast the highest concentration of ACE, organs such as renal proximal tubules, gastrointestinal tract, brain tissues and cardiac tissue contain ACE which exist as a membrane bound enzyme as well as a circulatory or globular enzyme **(Ortiz-Salmeróna, Baron, and García-Fuentes, 1998)**. The pivotal role of ACE and subsequently Angiotensin II in the sequence of events constituting the cardiovascular continuum make it a logical target in therapeutic strategies aimed at reducing the overall cardiovascular risk factor profile of an individual. According to Heart Outcomes Prevention Evaluation (HOPE) study, ACE inhibition with Ramipril reduces the risk of a primary cardiovascular event (cardiovascular death, stroke, or acute myocardial infarction) in high risk patients by 22% **(Sleight, 2000)**. ACE activity is inhibited by ACE inhibitors which has three distinct chemical structures namely sulfhydryl containing agents (Captopril), the non-sulfhydryl or carboxylic acid containing agents (Enalapril) and the phosphoric acid derivatives (Fosinopril). These inhibitors bind to the zinc ion in the ACE molecule through their sulfhydryl and phosphoric acid moieties whereas other inhibitory agents bind to the zinc through carboxyl residues **(Duncan, 1998)**. ACE inhibition has become a promising approach for drug targeting in the treatment of

cardiovascular diseases such as hypertension, heart failure, and diabetic nephropathy. Despite the benefits associated with ACE inhibitors, they exhibit adverse effects such as bronchospasm, rashes, cough (Wood, 1995) as well as contraindication in pregnancy.(Lip *et al.*, 1997; Steffensen *et al.*,1998). These side effects coupled with other factors such as high cost of conventional medicine and cultural practices point to the increasing interest in medicinal plants and the need to discover leads for novel drug discovery.




Figure 2. 2 Structures of synthetic ACE Inhibitors (Jallapally *et al.*, 2015).

#### 2.4. Traditional Use of Medicinal plants

Traditional medicine encompasses knowledge systems that have been established over generations based on the concepts, beliefs and practices native to diverse cultures prior to the modern medicine era. Since the beginning of human civilization, medicinal plants have been the backbone of traditional systems of healing throughout the globe and an integral part of history and culture. Plants have formed the basis of sophisticated traditional medicine systems that have been in existence for thousands of years and continue to provide mankind with new remedies. The earliest records dating from around 2600 BC, documents the use of some of the plant derived substances in Mesopotamia. These include oils of the *Cedrus species* (Cedar), *Glycyrrhiza glabra* (Licorice), and *Papaver somniferum* (Poppy juice), all of which are still in use today for the treatment of ailments ranging from coughs and colds to parasitic infections and inflammation. (Gurib-Fakim, 2006)

The use of herbal medicines and phytonutrients or nutraceuticals continues to expand rapidly across the world with many people now resorting to these products for the treatment of various health challenges in different national health care settings. This past decade has obviously witnessed a tremendous surge in acceptance and public interest in natural therapies both in developing and developed countries. It is estimated that more than 60% of the world's population utilize plants as their primary source of medicinal agents and moreover traditional medicine is still the only health resource available to about 60% of the world population. WHO estimates that between 60% and 90% of Africa's population rely on medicinal plants to totally or partially meet their healthcare needs (Taylor, 1999). This is valid considering 60% of South

Africa's population consult one of an estimated 200000 traditional healers. (Van wyk *et al*, 1997). Over 5000 plants are known to be used for medicinal purposes in Africa but only a few have been studied and scientifically validated.


The extensive use of traditional medicine in Africa, which composed mainly of medicinal plants, has been argued to be linked to cultural and economic reasons, this is why WHO encourages African member states to promote and integrate traditional medicinal practices into their health systems. WHO traditional medicine strategy 2014 – 2023 was developed and launched specifically for this purpose. As the use of medicinal plants grows, public health concerns surrounding their safety is increasingly recognized by regulatory authorities and further research is needed to ascertain the efficacy and safety of several of the practices and medicinal plants used by traditional medicine systems (WHO, 2003).

## **2.5. Plant Compounds and their biological activity**

Plants consist of a wide array of biologically active compounds which can act singly or synergistically together to confer therapeutic benefit. In medicinal plant study, it is of utmost importance to understand the various classes of chemical components present in a plant. Below are some of these plants compound and their biological activity.

### **2.5.1. Alkaloids**

Alkaloids are a large class of nitrogen containing secondary metabolites of plants, microbes or mammals. They exist as primary, secondary, tertiary or quaternary amines and are famous for their manifold pharmacological activities. *Papavier somniferum* from opium poppy is one of first alkaloids discovered. Alkaloids possess various remarkable biological activities such as antibacterial (berberine), antihypertensive (reserpine), vasodilation (vincamine) and anticancer (vincristine) properties.



**Figure 2. 3 Structure of Alkaloids (López *et al.*, 2018).**

### 2.5.2. Terpenes or Terpenoids

Terpenoids make up the largest class of secondary metabolites and are united by their common biosynthetic origin from acetyl-CoA or glycolytic intermediates. Terpenoids compounds are made up of five carbon units often called isoprene units that assemble in a regular pattern, usually head to tail in terpenes up to 25 carbons. They are classified on the basis of their isoprene units into monoterpenes (C<sub>10</sub>), sesquiterpenes (C<sub>15</sub>), diterpenes (C<sub>20</sub>), triterpenes (C<sub>30</sub>) and tetraterpenes (C<sub>40</sub>). They play different roles in plants such as in defence, thermotolerance, wound healing and pollination of seed crops.

**Table 2. 1 Different Classes of Terpenoids and their Examples.**

Class of Terpenoids	No of Carbons	No. of isoprene units	Example
Monoterpenoids	10	2	Thymol, Menthol
Sesquiterpenoids	15	3	Ginger, mints, Farnesol
Diterpenoids	20	4	Taxol
Sesterterpenoids	25	5	
Triterpenoids and steroids	30	6	Oleanolic acid
Tetraterpenoids	40	8	B-Carotene

#### ***2.5.2.1 Triterpenoids and steroids***

The biological activities of triterpenoids have attracted much interest. They are known as cancer chemo-preventive, antiulcer, antidiabetic agents, and inhibitors of angiogenesis. They are the biological active components of several famous herbal medicines, such as ginseng, licorice and bupleurum.

Steroids are triterpene derivatives that are based on the cyclopentane perhydrophenanthrene ring system. They are well known for their biological role as the hormones - testosterone and progesterone. Recent chemical investigation and pharmacological studies reveal various biological roles such as anti-inflammation, anti-cancer, anti-cardiovascular, hypoglycaemic and antifungal activities for steroidal compounds in plants (Liu, 2011).





Figure 2. 4 Structures of some Triterpenoids (Ruszkowski & Bobkiewicz-kozlowska, 2014)

### 2.5.3 Phenols

Phenols are a group of secondary metabolites characterized by the presence of one or several hydroxyl (OH) groups attached to an aromatic ring. Structures of phenolic compounds are very diverse and occur either as simple compounds with one aromatic ring or as complex polymers (polyphenols) with different functional groups attached. (Mann J *et al*, 1994).



**Figure 2. 5 Structure of Phenol**

**Table 2. 2 Different classes of Phenolic compounds**

Class of Phenols	Basic carbon skeleton	No. of carbons
Simple phenols and benzoquinones	C6	6
Naphthaquinones	C6-C4	10
Anthraquinones	C6-C2-C6	14
Flavonoids	C6-C3-C6	15
Tannins	(C6-C3-C6) <sub>n</sub>	N

### 2.5.3.1: Flavonoids

Flavonoids are the largest and most diversified group of polyphenolic compounds structurally based on a C<sub>15</sub> skeleton with a chromane ring bearing a second aromatic ring B in position 2, 3 or 4. Flavonoids constitute one of the most characteristic classes of compounds in higher plants, and are responsible for many of the plants colours that dazzle us with their brilliant shades of yellow, orange, or red. Classification of flavonoids is commonly based on the oxidation or saturation of the intermediate C ring. Major groups include the flavonols, flavones, isoflavonoids, and cyanidins.




Figure 2. 6 Basic structure of Flavonoid classes (Higdon, 2005)

Flavonoids are most commonly known for their antioxidant activity (**Balasuriya and Rupasinghe, 2011**) The capacity of flavonoids acting as antioxidants depends upon their molecular structure. The position of hydroxyl groups and other features in the chemical structure of flavonoids are important for their antioxidant and free radical scavenging activities. It has been discovered that flavonoids also provide other important biological activities such as antibacterial, antiviral, anti-allergic, antihypertensive, anti-allergic, antiplatelet, anti-inflammatory and anti-tumour activities (**Kumar and Pandey, 2013**).

## **2.6. Plants as Sources of ACE Inhibitors**

Plants produce and contain a vast and diverse assortment of organic compounds, the great majority of which do not appear to participate directly in growth and development. These substances or phytochemicals are divided into primary and secondary metabolites. The primary metabolites such as carbohydrates, lipids, and proteins are found in all plants and perform metabolic roles that are essential whereas secondary metabolites, which often are differentially distributed among limited taxonomic groups within the plant kingdom have an ecological role in regulating the interactions between plants and their environment. A number of these secondary metabolites in plants play defensive roles and serve as chemical messengers (**Liu, 2011**).

Secondary metabolites are the main source of pharmaceutical drugs and serve as templates for many medicinal derivatives currently in use. There are three broad categories of plant secondary metabolites; (i) Terpenes and terpenoids, (ii) alkaloids, (iii) phenolic compounds.

A number of compounds from plants have been identified to possess *in vitro* ACE inhibitory activity. These include alkaloids (**Oh et al., 2004**), phenylpropanes, proanthocyanin (**Anderson et al., 2004**), flavonoids (Oh et al., 2004; Loizzo et al., 2006), triterpenes and

terpenoids(Morigwa *et al.*, 1986; Hansen *et al.*, 1995), xanthones (C. H. Chen, Lin, Lin, and Hsu, 1992), fatty acids and peptides/amino acids (D. H. Lee, Kim, Park, Choi, and Lee, 2004). Recent studies indicate that phenolic rich foods and plants have the ability to inhibit ACE activity, both *in vitro* and *in vivo*. This was confirmed by Actis-Goretta *et al* (2006) whose study confirmed that flavanol and procyanidins exhibit ACE inhibitory activity. In that study, significant inhibition was observed for epigallocatechin which would support the idea that the extent of ACE inhibition could be associated with the number of hydroxyl groups in the flavanol/procyanidin available to establish hydrogen bonds with the ACE protein. This study will investigate *Centella asiatica* for ACE inhibition.

## 2.7 *Centella asiatica*

*Centella asiatica*, is a tropical medicinal plant from the Apiaceae family native to South Africa, Madagascar, and the Southeast Asian countries such as India, Sri Lanka, China, Indonesia and Malaysia. It is commonly known as gotu kola, asiatic pennywort, Indian pennywort, wild violet, and tiger herb. *Centella asiatica* is described as a hungry feeder with a penchant for a lot of water. It is found in abundance on moist, sandy or clayey soils, often in large clumps forming dense green carpets. In South Africa, *Centella asiatica* is found along the moist eastern parts, widely distributed from the Cape Peninsula northwards. The species has been identified to vary considerably in different parts of the world and is sometimes treated as several distinct species. (Van Wyk *et al*, 2002).

### 2.7.1. Botanical description

The plant is propagated from seeds or stolon's. The stems of the plant are green to reddish green in colour and have long stalked, green leaves with rounded apices. The edible leaves are kidney shaped, borne on pericladial petioles, smooth surfaced, thin and soft, with palmate

nerves and can measure from 1 – 5 cm in width. Nodes found on the stem give rise to long petioles (5 -15cm) which hold the leaves. The fresh or dried aerial parts consisting of leaves and stems are used for medicinal purposes (Bandara, Lee, and Thomas, 2011).



**Figure 2. 7 *Centella asiatica* (L.) Urban (Apiaceae).**

**Table 2. 3 Taxonomy of the Gotu Kola Plant**

Classification	Name
Kingdom	Plantae
Division	Magonoliophyta
Class	Magnolipsida
Order	Apiales
Family	Apiaceae
Genus	<i>Centella</i>
Species	<i>Centella asiatica</i> (L.) Urban

Ref: (Bandara *et al.*, 2011).

### 2.7.2. Therapeutic Uses of *Centella asiatica*

*C. asiatica* has been used for medicinal purposes since prehistoric times as an adaptogenic for enhancing cognitive function by revitalizing the brain and nerve cells. The plant possesses a wide range of health benefits that includes cardio-protective, anti-inflammatory, antiepileptic, sedative, antiviral, antibacterial, anti-ulcer, anti-psoriatic, anti-cytotoxic and anti-tumour properties ( **Orhan, 2012; Gohil et al, 2010**). *C. asiatica* is currently known as Gotu Kola in South Africa (Marketed by Flora force®). It is sold as a herbal product for treatment of variety of conditions such as wound healing, anxiety, depression and venous insufficiency. *C. asiatica* has been reported to act on the connective tissues of vascular walls, being effective in hypertensive micro-angiopathy and venous insufficiency and decreasing capillary filtration rate by improving microcirculatory parameters (**Cesaron et al, 1992**). Studies by **Sanctis et al (2001)** showed that the total triterpene fraction of *Centella asiatica* at 180mg/day improved venous hypertension together with capillary filtration rate and ankle oedema.

According to the **WHO (1999)**, the medicinal uses of *Centella asiatica* supported by clinical data include the treatment of wounds, burns, ulcerous skin diseases and the prevention of hypertrophic scars. Extracts of the plant have been used to treat second and third degree burns. Topical application of asiaticoside, a major constituent of *Centella asiatica*, significantly enhanced the rate of wound healing in normal and diabetic animals as assessed by an increase in collagen synthesis and tensile strength of the wound tissues (Shukla et al., 1999). Oral administration of extracts is used to treat stress induced stomach and duodenal ulcers (**Sairam, Rao, and Goel, 2001**). Uses reported in folk medicine but not supported by experimental or clinical data, include treatment for hypertension, anaemia, asthma, albinism, measles, cholera, bronchitis, cellulite, constipation, nephritis, nervous disorders, epilepsy, hepatitis, dysuria, haematemesis, jaundice, toothache and as an antipyretic, analgesic, anti-inflammatory and brain tonic. (**WHO 1999**)

### 2.7.3 Phytochemical Composition of *C. asiatica*

*Centella asiatica* have been reported to contain a lot of constituents belonging to various chemical classes. The chemical constituents identified include terpenes, phenols, alkaloids, polyacetylenes, carbohydrates, amino acids. Triterpene saponosides such as asiatic acid, madecassic acid, asiaticoside, madecassoside, betulunic acid, and thankunic acid are the major chemical classes contained in *C. asiatica*. The plant contains other triterpenes such as brahmic acid, centellin, centellicin, asiaticin, terminolic acid, centellasaponins A-D, ursolic acid, and pomolic acid (**Inamdar et al., 1996**).

Several flavonoid derivatives such as quercetin, kaempferol, patuletin, rutin, apigenin, castilliferol, castillicetin and myricetin has been reported to be present in *Centella asiatica* (Subban et al., 2008). Other classes of compounds present in *Centella asiatica* include polyacetylenes (e.g. cadinol, centellinol, centellin, centellicin, and asiaticin), Sterols (e.g., 11-oxoheneicosanil-cyclohexane, dotriacont-8-en-1-oic acid, sitosterol 3-O- $\beta$ -glucoside, stigmasterol 3-O- $\beta$ -glucoside, and castasterone), and phenolic acids (e.g., rosmarinic acid, 3,5-di-O-caffeoil quinic acid, 1, 5-di-O-caffeoil quinic acid, 3,4-di-O-caffeoil quinic acid, 4, 5-di-O-caffeoil quinic acid, ettacrynic acid, chlorogenic acid, and isochlorogenic acid (**Govindan et al., 2007; Siddiqui et al., 2007; Sondhi et al., 2010**)).

## 2.8 Drug discovery from Medicinal Plants

Plants have been used as a source of therapy for thousands of years. Medicinal use initially took the form of crude drugs such as tinctures, teas, poultices, powders and other herbal formulations. In more recent history, the use of plants as medicines has involved the isolation of active compounds that began with the isolation of morphine from opium in the early 19<sup>th</sup>









preparation of tinctures and fluid extracts. Fresh solvent is continuously added on top of the plant material until the recovered residue of the extraction solution turns out to be too little. The percolator is a cylindrical or conical container made from glass or metal with a tap at the bottom. The percolation is usually done at moderate rate (e.g 6 drops/min) until the extraction is completed before evaporation to get a concentrated extract.

### ***2.9.1.3 Extraction under Reflux***

Reflux involves the condensation of vapours and the return of the condensate to the system from which it originated. This method is suitable for heat stable compounds. It can be used for both small and large extraction for herbal study. During the extraction process, the material is immersed in an appropriate solvent in a round-bottomed flask, which is vertically connected to a condenser. The solvent is heated by water bath and electric heater to boiling. The evaporated solvent goes up into the condenser, and then is condensed into liquid form and drops down to the flask. Increasing temperature in the process generates heat that help the solvent penetrate into the plant cells thus increasing the solubility of compounds in the solvent. The extraction efficiency is higher and is one of the most popular method employed for herbal study.

### ***2.9.1.4 Sonication assisted solvent extraction***

Sonication assisted extraction is a modified maceration in which ultrasound is utilized to improve the extraction efficiency. The plant material is placed in a closed container like in maceration. The container is then placed in an ultrasonic bath. In such a condition, ultrasound transfers the mechanical power onto the plant cells, leading to breakdown of cell walls and increased solubilization of extracts in the solvent. Factors affecting extraction yields are frequency, length, and temperature. The benefit of SAE is mainly due to reduction in extraction

time and solvent consumption, therefore it is an easy and efficient method commonly used in the lab (Liu, 2011). However, use of ultrasound energy more than 20kHz may have an effect on the active phytochemicals through the formation of free radicals (Handa, Khanuja, Longo, and Rakesh, 2008)

#### ***2.9.1.5 Soxhlet extraction***

Soxhlet extraction is a convenient method for the extraction of herbal materials of small to medium volumes. The commercially available soxhlet instrument is composed of an extraction chamber with reflux condenser and collecting flask. The chamber is placed between the collection flask and refluxing condenser. The plant powder or pieces are kept in a cellulose thimble in the extraction chamber. A suitable solvent is added to the flask and heated under refluxing. The solvent will first be evaporated up into the condenser, then liquefied into the chamber. When the condensed solvent in the chamber reaches a certain height, it is siphoned into the flask, and the next extraction is initiated. Usually, 50 -60 times of recycling are necessary for an extraction. Because of the repeated extraction, this method is usually more efficient than refluxing and produces a higher yield of extract with less volume of solvent.

#### **2.9.2 Cardiovascular bioassay models for screening of medicinal plant**

In modern medicine, drug discovery has become more focused on ligands, substrates and inhibitors of specific target (enzyme and receptors) that play an important role in disease regulation. CVD models such as *in vivo* (laboratory animal model), *in vitro*, and *in silico*/computational models are used to assess the efficacy and safety of new drugs earlier in the drug development pipeline. These models are also utilized during pre- clinical evaluation to assess the risk of cardiac adverse effects of new drugs in humans. They remain a key tool for investigating the action of drugs on biological systems.

### ***2.9.2.1 in vivo models for cardiovascular activity***

*In vivo* assays assesses drug effects on the heart in its natural physiological environment. They are suited for observing the overall effect of an administered substance on a living subject, and measure endpoints such as electrocardiogram (ECG), blood pressure (BP) and heart rate (HR) via implantable telemetry. (Hanton, 2007). Some of the *in vivo* cardiovascular models commonly used include BP measurement in pithed rats, tail cuff method, direct measurement of BP in conscious rats with indwelling catheter, telemetric monitoring of CVD parameters in rats, hemodynamic evaluation in anesthetized dogs, the salt-induced hypertensive rat model, the salt-loaded hypertensive model, and the anaesthetized normotensive rat model.

Most of these *in vivo* cardiovascular models in experimental animals involve blood pressure measurement and are classified under direct and indirect methods of blood pressure measurement. The tail cuff method is an example of indirect method of measurement, it is considered a non-invasive method, technically non demanding and suitable for chronic studies considering that serious risk to animal health is minimal. However, one of the disadvantages is the need to restrain the animal and apply heat to achieve tail blood flow, these applications could negatively influence blood pressure reading. The use of an indwelling catheter is a direct method considered the most accurate method of measurement. It eliminates the need for restraining and heating the animal, but requires expertise to use (Kamadyaapa, 2008).

### ***2.9.2.2 in vitro models for cardiovascular activity***

*in vitro* assays refer to the technique of performing a given procedure in a controlled environment outside a living organism. It involves the use of isolated organs or parts of organs in a simulated physiological environment. *in vitro* models using isolated cells, cardiac tissues or whole perfused heart, provide simple, rapid and sensitive methods with a moderate to high

throughput, for screening drug candidates for biological activity. Some of the *in vitro* models used in CVD include the rabbit lung derived ACE inhibition assay, the Langendorff perfused heart model, working heart perfused heart model, isolated cardiac myocyte models. (Hanton, 2007).

The application of the assay method for screening for ACE inhibitory activity in natural product research is a very helpful method owing to its rapidity, accuracy, and simplicity. The ACE inhibitory activity has become an effective screening method in the search for new antihypertensive agents from herbal plants. Some of the *in vitro* assay methods used to examine the activity of ACE inhibitors are based on substrate usage, with the various methods distinguished by their substrates and measurement methods for their enzymatic reaction and product separation. These include Cushman and Cheung's (1971) method using hippuryl-histidyl-leucine (HHL) as a substrate, Holmquist, Bünning, and Riordan's (1979) method using furanacryloyl-tripeptide as a substrate, Elbl and Wagner's (1991) method using tripeptide dansyltriglycine as a substrate, Amos and Arieh's (1978) method using o-aminobenzoylglycyl-p-nitrophenylalanilproline as a substrate, and Lam, Himamura, Anabe, Shiyama, and Keda's (2008) method using 3-hydroxybutyrylglycyl-glycyl-glycine as a substrate. For the various methods, measurement of the enzymatic activity is carried out using either spectrophotometry, fluorometry, high-performance liquid chromatography, electrophoresis, or radiochemistry.

For this study, we utilized a fluorimetric ACE assay using HHL as the substrate as this assay is rapid, extremely sensitive and uncomplicated in terms of procedure, and has the advantage of greater efficiency with respect to both time and reagents over others (Schwager *et al*, 2006).

In this assay, the terminus of the ACE cleaved substrate binds to OPA which allows for enzyme activity to be measured based on a decrease or increase in fluorescence intensity. The presence of an inhibitor in the enzyme-substrate mix results in a decrease in fluorescence intensity. The choice of fluoroscopy as a mode of detection was due to its simplicity, sensitivity and rapidity,

and is particularly applicable for the screening of large numbers of ACE inhibitors and biological samples. The substrate and all the reagents are commercially available and relatively inexpensive. In addition, assay procedure required no solvent extraction, and has the advantage of greater efficiency with respect to both time and reagent. (Schwager *et al*, 2006).

## **2.10 Isolation and Purification techniques**

Selection of methods in the isolation and purification of herbal extract depends on the properties of the compounds and purposes of the study. Isolation and Purification techniques serves to remove interferences from the analyte as well as fractionate extracts into smaller fractions. In general, crude extracts are initially separated into several fractions by isolation methods such as liquid-liquid extraction.

Isolation and purification of compounds from complex herbal extracts depend on successful application of several isolation methods. Column chromatography and thin layer chromatography (TLC) are usually necessary, and mostly used due to their convenience, economy, and availability in various stationary phases. Many bioactive molecules have been isolated and purified by using thin layer and column chromatographic methods.

### **2.10.1 Thin layer chromatography**


TLC is an easy, economic, and quick method that can be used for qualitative and quantitative analysis as well as purification of natural products. Practically, TLC is used in identifying unknown compounds and detecting the presence of secondary metabolites in herbs. As an auxiliary tool, TLC is used to help optimize conditions of column chromatography, including selecting the mobile and solid phases, identifying compounds isolated from column as well as confirming purity of the isolates (Liu, 2011).

When applying TLC for isolation, the solution of herbal extract on a suitable solvent (ethanol or methanol) is spotted as a single round spot or thin line on a TLC plate near the edge of one side of the plate. The TLC plate is then obliquely placed into a well-closed tank containing sufficient volume of solvent but maintaining the sample spot or line above the surface of the solvent. The developing solvent will slowly migrate up the plate due adsorption of the solvent on the plate.

An important qualitative parameter in TLC of a particular sorbent and solvent system is the  $R_f$  value, which is defined as:

$$R_f = \text{Distance of analyte} / \text{Distance of solvent front}$$

In this study, TLC was used in the preliminary screening of crude plant extracts, monitoring columns during isolation and purification. TLC coupled with UV detection and spray reagents provide valuable information on type of compound in an extract. It is a simple, rapid and cost effective technique allowing for analysis of several samples.



### **2.10.2 Preparative Thin Layer Chromatography (PTLC)**

PTLC is generally applied for purification of compounds in a relatively simple fraction that have been separated by several chromatographic columns. Preparative thin layer chromatography (PTLC) is used to separate and isolate amounts of material larger than normal for analytical TLC. The quantities processed range from 10 mg to greater than 1 gram. In preparative TLC, materials to be separated are often applied as long streaks, rather than spots, in the sample application zone. The main difference from analytical TLC is in the thickness of the plates. It uses preparative plates of about 1 – 2 mm in thickness allowing the application of larger volume of sample. After development the plates are visualized under UV, and the band

with analyte of interest is scraped off together with the adsorbent using a spatula. The analyte is separated from the adsorbent by filtration and concentrated. (Sherma *et al* 1987)

### 2.10.3 Adsorption or Open Column Chromatography

The mechanism of this chromatography lies on the physical and chemical interaction between the stationary phase and solute molecules. The separation depends on the difference between the adsorption strength of the material in stationary phase to solute molecules in mobile phase. The solvent of the mobile phase competes with the solute molecules for adsorption sites. Normal silica gel is the commonly used in terms of stationary phase and is regarded as a typical polar sorbent. Due to differences in chemistry, analytes separate as they move down the column. These are continuously monitored by TLC and similar fractions are combined and concentrated. The column can be eluted in two ways, either using a single solvent or gradient elution - a series of solvents with increasing polarity/ elution strength are used to elute the column. This method is however considered very tedious and labour intensive.

### 2.10.4 High Performance Liquid Chromatography

HPLC is a popular method for chemical analysis and separation that is applied in many fields such as the food, pharmaceutical, and agricultural industries. HPLC achieves separation by utilizing the fact that different compounds have different migration rates given a particular stationary and mobile phase (Liu, 2011). HPLC fulfils different objectives; HPLC uses various separation mechanisms such as adsorption chromatography, partition chromatography (normal and reverse phase partition), ion exchange chromatography, ion pairing chromatography, and size exclusion chromatography. There is Analytical and Preparative HPLC. Analytical HPLC is used to determine the existence and possibly the concentration of analyte in a sample. It is often utilized in quality control of medicinal products. Preparative HPLC refers to the process

of isolation and purification of small quantitative compounds using HPLC. It has been applied to the isolation of most classes of compounds in herbal medicine, especially minor bioactive compounds. The HPLC instrument is an automated closed system consisting of a pump system to generate pressure, narrow columns packed with small particle size adsorbents and a detector. In this study, HPLC was used for further purification and separation of the fractions of the plant extracts. The detector emits a response to the eluting sample compound and subsequently signals a peak on the chromatogram. This is interpreted to guide the next step in the purification and identification process.

## **2.11 Compound Identification and Elucidation techniques**

Compound identification or structural elucidation involves the determination of factors such as atom number, number of bonds, configuration, and conformation of pure compounds. The development and improvement of analytical instrument to determine structure has been one of the biggest advancements in organic chemistry during the past 60 years. Today almost any structure can be determined with these instruments. Several techniques are available for elucidating and identifying the structure of a pure compound, these include NMR spectroscopy, Mass spectrometry, UV-vis Spectroscopy, and IR Spectroscopy.

### **2.11.1. Nuclear Magnetic Resonance (NMR)**

Structure elucidation and identification of novel compounds from natural and/or synthetic sources depends heavily on nuclear magnetic resonance (NMR) spectroscopy and mass spectrometry (MS) as complementary tools. Techniques such NMR spectroscopy are one of the important characterization tools in chemistry, often utilized to provide insight into the

chemical components or profile of samples such as crude extracts and their fractions, they also serve as a guide in the purification process of crude samples. NMR essentially provides simultaneous access to both qualitative (chemical structure) and quantitative information of samples and has the advantage of accuracy, reproducibility, and flexibility with respect to the nature of the analyte (Dayrit and Dios, 2017). In practice, the acquisition of a  $^1\text{H}$  NMR spectrum is the first step in NMR-based structure elucidation and metabolomics analysis. Especially for sample-limited natural products, 1D  $^1\text{H}$  NMR and its 2D counterparts are typically the first-line structural tools employed for identification and de-replication purposes. Three key parameters are important in the interpretation of NMR spectra, these include chemical shift, spin-spin coupling (splitting) and peak intensity. The chemical shift is the resonant frequency of a nucleus relative to a standard in a magnetic field. One factor that affects chemical shift is the changing of electron density from around a nucleus, such as a bond to an electronegative group. Hydrogen bonding also changes the electron density in  $^1\text{H}$  NMR, causing a larger shift. Chemical shifts can be used to identify structural properties in a molecule based on our understanding of different chemical environments. Another useful parameter that allows NMR spectra to give structural information is spin-spin coupling. Different spin states interact through chemical bonds in a molecule to give rise to this coupling. In NMR spectra, this effect is shown through peak splitting that can give direct information concerning the connectivity of atoms in a molecule. Nuclei which share the same chemical shift do not form splitting peaks in an NMR spectra. Peak intensity otherwise known as size of the peaks, the size of the peaks in the NMR spectra can give information concerning the number of nuclei that gave rise to that peak. This is done by measuring the peak's area using integration. Yet even without using integration the size of different peaks can still give relative information about the number of nuclei.

2-D NMR in itself is an effective tool for identification and validation of structure, especially for understanding the stereochemistry of the molecule. The commonly used 2D-NMR techniques are COSY (used to determine the chemical shift, coupling relationship and connecting sequence of protons), NOESY (provides important information about stereochemistry and solution conformation by analysing the connectivity through the space of atoms in a molecule), TOCSY (allows for easy recognition of complete families of coupled spins), HMQC and HSQC (allows for determination of carbon to hydrogen connectivity), and HMBC (used to identify proton nuclei that are separated by more than one bond) (Liu, 2011).

### 2.11.2 IR Spectroscopy

Infrared spectroscopy is useful in the identification of functional groups present in a compound. This technique is based on the absorption of electromagnetic radiation at wavelengths ranging between 4000 and 400  $\text{cm}^{-1}$ . At this range of wavelengths specific functional groups give characteristic vibration, bending and stretching. This is recorded by the IR instrument and gives basic information on the structure (Liu, 2011).

### 2.11.3 Mass Spectroscopy

Mass spectroscopy measures the molecular masses of individual compounds and atoms precisely, by converting them into charged ion. It is widely used for structural elucidation because of its unique ability to accurately measure molecular mass and to provide structural related information. The evolution of MS is mainly characterized by the development of an ion source and a mass analyser (Liu, 2011).

In mass spectroscopy, a compound is ionized, the ions are separated on the basis of their mass/charge ratio, and the number of ions representing each mass/charge unit is recorded as a spectrum. The recorded spectrum is analysed to reveal the molecular weight of the compound

and the fragmentation patterns of the parent compound. For unknown compounds, the molecular ion, the fragmentation pattern, and complementary forms of spectrometry such as IR and NMR put together give a molecular skeleton of the compound (Silverstein *et al*, 2005).

#### 2.11.4 UV Spectroscopy

Ultra violet spectroscopy is mainly concerned with electron transitions in conjugated systems, it uses UV-VIS light to measure the transmittance of the light that passes through a sample. the positions and intensities of the absorption band maxima depend to a large extent on the particular system under construction. The electron transitions are very sensitive to structural changes and reflect the strains imposed on the system by steric and electronic interactions. The absorption wavelength is in the range of 200 – 800 nm; it is usually possible to correlate changes in spectra with changes in structure with a fair degree of success. One of the strengths of UV spectroscopy is the relative simplicity and inexpensiveness of the instrumentation coupled with great sensitivity. The ease with which quantitative measurement can be made is also a great advantage (Scheinmann, 2013).

In this project, *Centella asiatica* will be subjected to bioassay guided fractionation following the drug discovery processes described above. The subsequent chapters present a further discussion of these processes and their outcomes.

## Chapter 3

### Materials and Method

---

#### Introduction

Bioassay guided fractionation is an important strategy in the scientific validation of the claimed traditional uses of medicinal plants as well as the discovery of novel compounds. The successful isolation of bioactive compounds from indigenous medicinal plants often validates indigenous knowledge, adds value to plants and supports the knowledge of their medicinal use. It may also contribute to the research and development of new pharmaceutical drugs for the treatment of various diseases. In this study, bioassay guided fractionation was used to isolate compounds exhibiting ACE inhibition from extracts of *Centella asiatica*.

The process starts with the extraction of the powdered plant parts, followed by sequential fractionation of the active crude extract using gravity column chromatography and solvents of increasing polarity. Fractionation and isolation was guided by ACE activity assay of the various fractions. Active fractions were further purified using high performance liquid chromatography (HPLC), and the isolated compounds characterized by NMR, HRMS, IR and UV spectral data and by comparison to literature data.

This chapter reports on the materials and equipment used in the isolation of active compounds and the bioassay used to guide the isolation of active compounds. The chapter also describes the methods applied and the protocols followed for the extraction, fractionation, isolation, purification and identification of the active compounds from *Centella asiatica*, as well as the ACE inhibition assay used to determine bioactivity of the extracts/fractions/compounds.

### 3.1 Equipment and Materials

The equipment and materials used in the study are as follows:

**Table 3.1 List of chemicals**

<b>Chemical</b>	<b>Supplier</b>
<b>Methanol (HPLC grade)</b>	Sigma Aldrich Co.
<b>Deuterated methanol</b>	“
<b>Chloroform</b>	“
<b>Hydrochloric acid (10M)</b>	“
<b>2,2-diphenyl-1-picrylhydrazyl (DPPH)</b>	“
<b>Sodium Hydroxide Pellets</b>	“
<b>Sodium Chloride</b>	“
<b>Ascorbic acid</b>	“
<b>Boric acid powder</b>	“
<b>o-phthaldialdehyde</b>	“
<b>N-Hippuryl-His- Leucine (HHL)</b>	“
<b>Rabbit lung derived Angiotensin- converting enzyme</b>	“
<b>Acetic acid (HPLC grade)</b>	“
<b>Hexane (Analytical reagent Grade)</b>	Lab Chem.
<b>Ethyl acetate (Analytical reagent Grade)</b>	Lab Chem.
<b>Ethanol (Analytical reagent Grade)</b>	Lab Chem.

**Table 3. 2 List of equipment**

<b>Equipment</b>	<b>Model and Manufacturer</b>
<b>-86°C ultralow freezer</b>	<i>(Nuair, Model NU 9668E, Nuair, Japan)</i>
<b>Oven</b>	<i>(Labotech, Model LDO-080F, DaihanLabotech Co Ltd, Korea)</i>
<b>Analytical balance</b>	<i>(Ohaus, Model PA413, Ohaus Corporation, USA),</i>
<b>Vortex mixer</b>	<i>(KK, Model VM-300, Germany Industrial Corp, Taiwan),</i>
<b>Vacuum pump</b>	<i>(Buchi, Model V-500, BuchiLarbotechnik AG, Switzerland)</i>
<b>Rota vapor</b>	<i>(Buchi, Model R-11, BuchiLarbotechnik AG, Switzerland)</i>
<b>Freeze drier</b>	<i>(Virtis TM mobile freeze-dryer, model 125L),</i>
<b>Buchner funnel</b>	
<b>Whatman No. 1 paper filter, nylon syringe filters</b> <i>(25 mm diameter, 0.45 µm pore size)</i>	
<b>Micropipettes, HPLC filter unit</b>	<i>(Millipore Cameo 25 AS, DDA 02025So MSI: Micro separation Inc., USA),</i>
<b>Membrane filters</b> <i>Durapore, 0.45 µm HV, Millipore</i>	
<b>FTIR instrument fitted with UATR and controlled with Spectrum® software version 6.3.5.0176</b>	<i>(Perkin-Elmer 100)</i>

### **3.1.1 Equipment for the isolation, purification, and identification of bioactive compounds**

#### **3.1.1.1 Column Chromatography**

Waters Sep. Pak<sup>®</sup> column (2.5 x 10cm) with silica gel 60 (0.063-0.0200 mm) (Merck, Germany), was used for column chromatography.

#### **3.1.1.2 Thin Layer Chromatography**

Thin layer chromatography was carried out on pre-coated silica gel F<sub>254</sub> plates (Merck) with a 0.2 mm layer thickness. Visualization of the TLC spots was carried out under UV light at 254nm and/or 366nm, and further detection of compounds was achieved by spraying with DPPH solution.

#### **3.1.1.3 High-Performance Liquid Chromatography**

Sample purification was carried out using an Agilent Technologies 1260 Infinity (Made in Germany), a PC with Chemstation<sup>®</sup> software (Agilent, OpenLAB CDS) and a normal phase column (Whatmann 10uM silica prep column 50cm).

#### **3.1.1.4 Nuclear Magnetic Resonance (NMR) Spectroscopy**

NMR spectra were recorded on an NMR Spectrometer (Bruker Avance 400 MHz Rheinstetten, Germany). All spectra were referenced to residual solvent signals ( $\delta_H$  of 7.2600 ppm and  $\delta_C$  of 77.00 ppm) of deuterated chloroform ( $d_{CDCl_3}$ ) and ( $\delta_H$  of 4.87, 3.31 ppm and  $\delta_C$  of 49.10 ppm) Methanol. Chemical shifts were measured in parts per million (ppm) and coupling constants ( $J$ ) in Hz.

#### **3.1.1.5 Fourier Transform – Infra-Red Spectroscopy**

Fourier transform infrared spectroscopy (FTIR) measurements were carried out using an IR spectrometer (Perkin Elmer spectrum 400 FT-IR/FT-NIR spectrometer, Waltham, USA) controlled with Spectrum<sup>®</sup> software version 6.3.5.0176 Spectrum 100 (PerkinElmer).

### **3.1.1.6 UV Visible Spectroscopy**

UV spectrum of the isolated compounds were obtained using a UV-Visible spectrophotometer (GBC CINTRA 202 New Hampshire, USA). All sample for analysis were dissolved in methanol.

### **3.1.1.7 Mass Spectroscopy**

High-resolution mass spectroscopy (HRMS) analysis was conducted using a Waters ultra-pressure liquid chromatograph (UPLC) (Waters, Midford, USA) using electrospray ionization (ESI) interface working in positive mode.

## **3.2 Plant Material**

Dried *Centella asiatica* leaf powder was purchased from Warren Chem. Specialties LTD (Cape Town, South Africa). The powdered plant came in sealed, opaque, plastic bags and was stored in a dark place away from light in a temperature controlled (20°C) laboratory room. The Certificate of Analysis of purchased plant powder is included in Appendix XXXIV of the thesis.

## **3.3. Experimental methods**

### **3.3.1. Preparation of crude extracts of *C. asiatica***

The dried leaf powder of *C. asiatica* (500 g) was macerated in 2.5 L of methanol, ethanol or distilled water for 24 hrs, with occasional shaking. The resulting methanol or ethanol extract was filtered under vacuum using a Buchner funnel with Whatman No 1 filter paper, and the residual plant material subjected to further maceration with fresh solvent up to three times to ensure exhaustive extraction. Individual methanol or ethanol filtrate were combined and

concentrated under reduced pressure at 40 °C using the rotary evaporator (Buchi Rota vapour), the obtained extract was transferred to a pre-weighed beaker and stored at -8 °C. Aqueous extracts were filtered using the Buchi vacuum pump, the filtrate was transferred to 250 ml round bottom flasks, flash frozen with liquid nitrogen and thereafter freeze dried under vacuum for 96 hours. The resultant freeze-dried extract was stored in airtight amber bottles and put in a desiccator. The percentage yield of the organic and aqueous extracts were then determined and recorded.

### **3.3.2. ACE Inhibition assay**

#### **3.3.2.1 Preparation of sodium borate buffer solution**

To prepare sodium borate buffer solution (1 L) with a final concentration of 0.4 M Boric acid, 0.3 M NaCl and a pH of 8.3, NaCl (17.532 g) and Boric acid (24.732 g) were weighed into a 1 L bottle, and 900 ml of distilled water added. The mixture was stirred using a magnetic stirrer for 25 minutes to ensure complete dissolution, and the pH determined, and adjusted to 8.3 using 0.1 M NaOH solution. The final volume of buffer was made up to 1 L and the pH was again measured and confirmed as 8.3.

#### **3.3.2.2 Preparation of Hip-His-Leu (HHL) solution**

HHL stock solution (116.4 mM) was prepared by dissolving 50 mg of HHL in 1 ml acetic acid in a 10 ml glass vial and stored at -5 °C. To prepare the 3.5 mM HHL required for the ACE assay, 300 µL of the HHL stock solution (116.4 mM) was pipetted into a glass vial, followed by 570 µL of 10M NaOH. Previously prepared sodium borate buffer solution (section 3.3.2.1) was then added up to the 8 ml mark on the glass vial, and the pH of the solution determined and adjusted to 8.3 using 0.1M NaOH. The solution was made up to a final volume of 10 ml using sodium borate buffer solution and stored at -5 °C.

### 3.3.2.3 Preparation of ACE solution

ACE solution was prepared by adding 200  $\mu\text{L}$  of NaB buffer solution to the ACE vial containing 0.25 Units of Purified rabbit lung Enzyme. The ACE solution was stored at a temperature of  $-20\text{ }^{\circ}\text{C}$ .

### 3.3.2.4 Preparation of plant extracts for ACEI assay

Crude organic extracts of *C. asiatica* were dissolved in DMSO in NaB buffer (10% DMSO for the ethanol extract and 5% DMSO for the methanol extract). A preliminary ACE assay using DMSO in NaB buffer (5% and 10%) as enzyme inhibitors did not produce any inhibition of the angiotensin converting enzyme activity. The aqueous extracts were freely soluble in the buffer solution.

### 3.3.2.5 ACE inhibitory assay protocol

ACE inhibition activity of the crude extracts/fractions/compounds of the plant was performed according to the method of Balasuriya and Rupasinghe (2012) with some modifications. In this assay, ACE cleaves the substrate HHL to expose a free N-terminus which is then labelled with o-phthalaldehyde (OPA), a fluorescence agent.

Preliminary ACEI assay of the crude extracts was carried out at concentrations of 1 mg/ml, 6 mg/ml, and 10 mg/ml of the methanol, ethanol and aqueous extracts. ACE enzyme (5.6  $\mu\text{L}$ ) was pre-incubated with 5  $\mu\text{L}$  of either the various concentrations of the plant extracts (enzyme inhibitor), deuterated water (negative control) or captopril (positive control) for 20 minutes in sodium borate buffer making a final volume of 25  $\mu\text{L}$ . After incubation, 10  $\mu\text{L}$  of HHL (3.5 mM) was added to the solution to give the reaction mixture a final volume of 35  $\mu\text{L}$ . The reaction mixture was incubated at  $37^{\circ}\text{C}$  for 60 minutes and the reaction terminated by adding 150  $\mu\text{L}$  of NaOH (0.34M). OPA (20  $\mu\text{L}$  of a 20 mg/ml) was added, the mixture incubated for 10 minutes at room temperature, and the reaction terminated by adding 50  $\mu\text{L}$  of 3M HCl. This

final assay solution was pipetted in 150  $\mu$ L aliquots into a 96 well microtitre plate and fluorescence read on a fluorescent plate reader (Synermix, Biotek) at 355nm excitation and 535nm emission wavelengths. All experiments were carried out in triplicates.

Percentage inhibition of ACE activity was then calculated from fluorescence intensity according to the following equation:

$$\text{ACE Inhibition (\%)} = (F_{\text{control}} - F_{\text{sample}}) / (F_{\text{control}} - F_{\text{blank}}) \times 100$$

Where:

$F_{\text{control}}$  = Fluorescence intensity of the control (Inhibitor replaced with D<sub>2</sub>O)

$F_{\text{sample}}$  = Fluorescence intensity of the enzyme inhibitor (captopril, crude extract, or fractions).

$F_{\text{blank}}$  = Fluorescence intensity of blank (enzyme and inhibitor replaced with distilled water)

The inhibitory concentration 50 % (IC<sub>50</sub>) was calculated from a dose-response curve obtained by plotting the percentage inhibition versus the concentrations of sample with the use of Microsoft excel and Graph Pad Prism 6 Software.

The ACE assay of fractions of the crude extracts and compounds isolated during the fractionation and purification process was carried out according the ACE assay protocol above.

### **3.3.2.6 IC<sub>50</sub> determination of captopril**

The half maximal inhibitory concentration (IC<sub>50</sub>) value of the reference inhibitor captopril was determined from a concentration - response curve using Graph Pad Prism software version 5 and Microsoft excel. The IC<sub>50</sub> value obtained was then compared with the value reported in

literature. A Stock solution of 1 mg/1ml captopril in buffer solution was initially prepared, and dilutions done to obtain various assay concentrations of 200, 100, 50, 25, 12.5 and 6.25 ng/ml. The Percentage inhibition of each of the captopril working solution was determined using the ACE assay protocol from section 3.3.2.5.

### **3.3.3. Bioassay-guided fractionation of methanol crude extracts**

Gravity column chromatography, thin layer chromatography and high performance column chromatography were utilized to isolate the bioactive compounds. Silica gel (70 – 230 mesh) was used as the stationary phase and ethyl acetate/hexane in varying proportions as mobile phase for the column chromatography. Thin layer chromatography was used to determine and monitor the purity of fractions collected.

#### **3.3.3.1 Preparation of methanol sample for fractionation and isolation**

Methanol extract (1g) was dissolved in methanol, and the solution adsorbed unto 1.5 g of silica gel (35 -70 Mesh) in a round bottom flask. Excess methanol was evaporated on a rotary evaporator at 40°C leaving behind dried granules.

#### **3.3.3.2 Fractionation of Methanol extract by Column Chromatography**


Fractionation was carried out using the Waters Sep Pak<sup>®</sup> 35cc Vac Cartridges (2.5 x 10cm), of which the tip was plugged with a small cotton wool. Silica gel 60 -230 Mesh (10g) in hexane solvent was loaded unto the column. The adsorbed sample of the methanol crude extract was added on top of the loaded silica gel column followed by a cotton wool plug to prevent the impact of the eluting solvent disturbing the sample. The column was eluted with a stepwise gradient consisting of 50 ml volumes of hexane (100%), hexane: ethyl acetate [(80:20), (60:40), (40:60), and (20:80)], ethyl acetate (100%), methanol: ethyl acetate (50:50) and methanol (100%). Each of the fractions collected were dried and transferred to a pre-weighed glass vial.

The weight and % yield of each fraction was determined and subsequently screened for ACE inhibition activity using the protocol described in section 3.3.2.5.

The decision to fractionate 1g at a time (up to 15 times), was to avoid the loss of important components that might emanate from upscaling to a larger glass column. A total of 15g of crude extract was fractionated using 750 ml of solvent in the various ratios as described above.

**Table 3. 3 Solvent System used for Fractionation of *Centella asiatica* methanol Crude Extract.**

Solvent (% v/v)	Volume (mL)	Fraction	Yield(mg)	% Percentage Yield
Hexane 100	750	M <sub>1</sub>	-	-
Hex: EtoAc 80:20	750	M <sub>2</sub>	35	0.23
Hex: EtoAc 60:40	750	M <sub>3</sub>	240	1.6
Hex: EtoAc 40:60	750	M <sub>4</sub>	150	1.0
Hex: EtoAc 20:80	750	M <sub>5</sub>	118.5	0.79
Ethyl acetate 100	750	M <sub>6</sub>	57	0.38
EtoAc: Methanol 50:50	750	M <sub>7</sub>	8005.5	53.37
Methanol 100	750	M <sub>8</sub>	4455	29.79



**Figure 3. 1 Schematic Diagram for Bioassay guided fractionation of Methanol Extract**

### 3.3.3.3 ACE activity screening of fractions

Column fractions of the methanol extracts of *C. asiatica* were screened for ACE activity using the ACE assay method described in section 3.3.2.5. Fractions M<sub>4</sub> and M<sub>5</sub> were determined to be the most active inhibitors of ACE, and were analysed by means of a thin layer chromatography and <sup>1</sup>H NMR spectroscopy to establish their purity.

### 3.3.3.4 HPLC purification of active fractions

Fractions M<sub>4</sub> and M<sub>5</sub> were purified on Agilent HPLC equipped with RI detector, and Whatman 10um Silica Prep column 50cm as stationary phase. Samples were run at a flow rate of 3ml/min, UV wavelength; 254nm, at Temperature of 30°C with a runtime of 30 - 50 minutes.

#### 3.3.3.4.1 Purification of Fraction M<sub>4</sub>

Fraction M<sub>4</sub> 150 mg (H4E6) was purified using normal phase HPLC [Whatman 10um Silica Prep column (50cm), isocratic elution using 1000 ml Hexane (40%): Ethyl acetate (60%) over 30 min, 3ml/min, detection at 254nm]. The sample was dissolved in 2.5ml of Hexane: Ethyl acetate (40:60), and 100 uL was first injected into the HPLC and eluted for 50 minutes, to establish the runtime. Subsequent injections of 200 uL each were ran for 30 minutes, with a total of 10 injections. All distinct peaks, ethyl acetate wash and fractions with no distinct peaks were collected.

#### 3.3.3.4.2 Purification of fraction M<sub>5</sub>

Fraction M<sub>5</sub> 118.5 mg (H2E8) was purified using normal phase HPLC [Whatman 10um Silica Prep column (50 cm), isocratic elution using 1000 ml Hexane (20%): Ethyl acetate (80%) over 30 min, 3ml/min, detection at 254nm]. Prior to injection onto the column, fraction M<sub>5</sub> (118.5 mg) was dissolved in 2.0 ml hexane: Ethyl acetate (20:80), filtered, and 100 µl injected into the HPLC and eluted for 30 minutes. A total of 8 injections were carried out. Similar peaks were collected for each injection and combined together. All prominent peaks, ethyl acetate wash and fractions without distinct peaks were collected. Collected sample were dried by rotary evaporation, and sent for NMR analysis.

### 3.3.4 Bioassay-guided fractionation of ethanol crude extract

The ethanol extract of *C. asiatica* was chemically profiled by NMR spectroscopy, which revealed the presence of fatty acids. The ethanol extract was therefore defatted by dissolving the ethanol extract (2.3g) in methanol (100 ml) and transferred to a 500 ml separating funnel,


100 ml hexane was added followed by 20 ml distilled water. The separating funnel was stoppered, shaken and left to stand for 20 minutes. Fatty acid, hexane and aqueous methanol layers were separately collected, dried and assayed for ACE activity.

#### **3.3.4.1** *Fractionation of the hexane fraction of the ethanol extract.*

The hexane fraction was chosen for fractionation following ACE activity assay screening. The hexane fraction (300 mg) was dissolved in 1ml of hexane: ethyl acetate (80:20) and was loaded onto the waters sepak column (2.5x 10cm) containing 10g silica gel. The column was eluted using a total of 400 ml of hexane: ethyl acetate (80:20). Fractions were collected in small volumes using glass vials to a total of 37 fractions. Collected fractions were immediately analysed on silica gel TLC plates to identify and pool similar fractions together. F2, F3 and F4 were pooled together as fraction A; F5, F6, F7, F8, and F9 as fraction B; F10, F11, F12, F13, F14, F15, F16, F17, F18 as fraction C; F19.F20, F21 as fraction D; F22, F23, F24, F25, F26, F27, F28 as fraction E; and F29, F30, F31, F32, F33, F34, F35, F36, F37 as fraction F. All pooled fractions were assayed for ACEI activity using the assay protocol in section 3.3.2.5.

##### **3.3.4.1.1** *HPLC Purification of Fraction A*

Fraction A was chosen for further purification by virtue of the NMR chemical profile and ACE inhibition activity. Fraction A (80mg) was purified using normal phase HPLC [(Whatman 10um Silica Prep column (50cm); isocratic elution using 1000ml hexane (80%): ethyl acetate (20%) over 45 min; 3ml/min; detection at 254nm]. The sample was dissolved in 1ml of hexane (80%): ethyl acetate (20%), followed by an initial 100 µl injection to establish the separation time, then subsequently four 200 µl injections were carried out. A total run time of 30 minutes was established for the purification. Pronounced peaks were collected, dried in a glass vial and assayed for ACE activity. A total of 5 peaks and ethyl acetate wash were collected.



**Figure 3. 2 Schematic Diagram for Bioassay guided fractionation of Hexane extract from Defatted Ethanol extract.**

#### 3.3.4.1.2 NMR Spectroscopy of fraction A

NMR spectral profile of sub-fraction A<sub>3</sub> (fraction of interest) revealed the presence of a coumaric acid structure. However, due to the difficulty in obtaining a clear elucidation of the full structure of the compound of interest as well as the insufficient quantity obtained from the purification process, methyl, ethyl and butyl esters of the synthetic coumaric acid were synthesized and compared to the proposed structure of the compound from fraction A<sub>3</sub>. This was to facilitate the confirmation of the proposed structure.

Synthesis of the compounds were done by reflux, to synthesize the methyl ester of coumaric acid, 1.0014g of coumaric acid was added to a three neck round bottom flask, followed by 10ml Methanol and 0.5ml Conc. H<sub>2</sub>SO<sub>4</sub>. The sample was spotted on a TLC at intervals during the reflux process to monitor the progress of the reaction. After 2 hours, the sample was pipetted into a 250ml round bottom flask and dried on a rotary evaporator. Once drying was completed, 10ml CHCl<sub>3</sub> (ethyl acetate) was added to the flask, swirled and transferred to a separating

funnel. Distilled water was added leading to formation of layers, the aqueous layer was removed and saturated aqueous NaHCO<sub>3</sub> added. The aqueous layer was again removed, and the remaining sample washed with brine and allowed to dry in a desiccator.

For ethyl and propyl ester of coumaric acid, the process above was repeated but replacing Methanol with ethanol and propan-1-ol respectively.

### **3.3.5 NMR Spectroscopy of extracts/fractions/compounds**

In this study, NMR was easily available and cost-effective, therefore was utilized for profiling of extracts/fractions/compounds and to monitor their purity at various steps. The preliminary analysis of the crude extracts and fractions were based on their 1D <sup>1</sup>H and 2D <sup>13</sup>C NMR and only the final pure compounds required a full data set (<sup>1</sup>H, <sup>13</sup>C, DEPT, COSY, HMBC, HSQC). The methanol and ethanol crude extracts were initially profiled to get an overview of their chemical profile. As the fractionation process progressed, the resultant fractions at each stage were also profiled to determine the effect of solvent elution on the resolution of various components of the sample as well as to establish the purity of the fractions.

For the NMR analysis, Crude extracts were dissolved in 0.5ml CDCl<sub>3</sub>, whereas some fractions and compounds were dissolved in 0.5ml CD<sub>3</sub>OD due to poor solubility in CDCl<sub>3</sub>. Samples were subsequently transferred into a 5mm NMR tube, which were sealed, appropriately labelled and sent for analysis. The generated spectra were analysed using Topspin® Software 4.0.6. Analysed samples were retrieved by evaporation

### **3.3.6 Spectroscopic identification of bioactive compound**

#### **3.3.6.1 Infrared (IR) spectroscopy**

Infrared spectra of the isolated compounds was measured using Perkin Elmer Spectrum 400 Fourier transform infrared (FTIR) spectrometer (Perkin Elmer Scientific, Massachusetts, United States). For measurement, a small dried sample of the isolated compounds were transferred to the crystal of the IR equipment. Some Pressure (50 - 60%) was gradually applied using the pressure gauge. Measurement was done at a scanning range of 500 to 4000  $\text{cm}^{-1}$ . Generated IR spectra was identified and analysed by OMNIC Spectra software provided in the spectrometer system.

#### **3.3.6.2 UV Spectroscopy**

UV spectrum of the compounds were measured using GBC CINTRA 202 UV-VIS Spectrometer. Samples were dissolved in methanol. The UV spectra was analysed using the CINTRA Software.

#### **3.3.6.3 NMR spectrometry**

NMR experiments were performed using Bruker AV400 NMR spectrometer in the NMR laboratory of University of the Western Cape. Samples were dissolved in deuterated methanol ( $\text{CD}_3\text{OD}$ ) and  $\text{CDCl}_3$ , filtered through a cotton wool plugged Pasteur pipette and placed in 5mm NMR tube (Norell, United States). All chemical shifts were referenced relative to corresponding residual solvent signals. The NMR experiments were analysed using “Topspin” software. The 1D  $^1\text{H}$  and  $^{13}\text{C}$  NMR experiments were carried out. In addition, homo and Heteronuclear 2D experiments, Correlated Spectroscopy with gradients (g-COSY) and Heteronuclear Multiple-Quantum Correlation (HMQC) spectra were also acquired.

#### **3.3.6.4 Mass spectrometry**

Mass Spectroscopy analysis of compounds was carried on Waters Synapt G2 quadrupole time-of-flight (QTOF) mass spectrometer at Stellenbosch University Central Analytical Facility. 1 mg of each sample was dissolved in 1 ml methanol (Romil), followed by a further 10-fold dilution into methanol. 2  $\mu$ L of sample was injected into a stream of methanol flowing at 0.3 ml/min, using a Waters ultra-pressure liquid chromatograph (UPLC) (Waters, Midford, USA) which conveyed the sample to a Waters Synapt G2 quadrupole time-of-flight (QTOF) mass spectrometer used for high-resolution accurate mass analysis.

Data was acquired in scan mode, the mass spectrometer was optimized for best sensitivity, a cone voltage of 15 V, desolvation gas was nitrogen at 650 L/hr and desolvation temperature 275 °C. The instrument was operated with an electrospray ionization probe in the positive mode. Sodium formate was used for calibration and leucine encephalin was infused in the background as lock mass for accurate mass determinations

#### **3.3.7 Antioxidant assay of compounds**

##### **3.3.7.1 DPPH radical scavenging assay**

Radical scavenging activity of isolated active compounds against stable DPPH (2,2-diphenyl-1-picrylhydrazyl hydrate, Sigma-Aldrich, Germany) was determined using spectrophotometry. When DPPH (a stable free radical) reacts with an antioxidant compound, which can donate hydrogen, it is reduced. The change in form of absorbance were measured at 517 nm on a UV/visible light spectrophotometer (GBC CINTRA 202 New Hampshire, USA).

The radical scavenging activity of the extracts were estimated according to the procedure described by (Shen *et al.*, 2010) with some modification. Briefly, a 0.1 mM solution of DPPH

radical solution in methanol was prepared and 1 ml of this solution was mixed vigorously with 3ml of different concentrations (31.25, 62.5, 125, 250, 500 µg/ml in methanol) of each extract. After 30 min incubation in the dark and at room temperature, absorbance (*A*) was measured at 517 nm using a UV/VIS spectrometer. The percentage of the radical scavenging activity (RSA) was calculated based on the following equation:

$$\text{DPPH Scavenged (\%)} = (A_{\text{Control}} - A_{\text{Sample}}) / A_{\text{Control}} \times 100$$

*A*<sub>control</sub> and *A*<sub>sample</sub> are the absorbance values (at 517 nm) for the control and sample, respectively.

Sample = 1ml DPPH solution + 3ml Sample Solution

Negative Control = 1ml DPPH solution + 3ml Distilled water

Positive Control = 1ml DPPH + 3ml various concentration of Ascorbic acid in methanol

EC<sub>50</sub> value, defined as the concentration of the sample leading to 50% reduction of the initial DPPH concentration, was obtained from the linear regression of plots of mean percentage of the DPPH radical scavenging activity against the concentration of the test compounds (µg/ml) obtained from three replicate assays.

### 3.4 Statistical Analysis

All experiments/ analyses in the present study were performed in triplicates. The results were expressed as mean. EC<sub>50</sub> and IC<sub>50</sub> calculation were determined from non-linear regression plot using Microsoft Excel.

The next chapter will report and discuss the results of the methods, protocols and procedures followed above.

## Chapter 4

### Results

---

#### Introduction

ACE Inhibitors remain crucial in the therapeutic strategy for treatment of cardiovascular disease and its risk factor hypertension. Increasing use and interest in plants highlights the need for increased research towards their quality control to ensure safe use, validate their claimed therapeutic efficacy and to identify bioactive/novel compounds that could replenish already depleting therapeutic armory and provide alternatives to ineffective and problematic drugs. The aim of the study was to isolate a bioactive ACE inhibitory compound, thereby validating the claimed use of *Centella asiatica* in the treatment of hypertension.

This chapter presents the results of ACE inhibition activity screening of crude extracts, fractions and compounds derived from the bioassay guided fractionation of *Centella asiatica*, and the identification, elucidation and antioxidant activity of the isolated bioactive compounds.

#### 4.1 Extraction and preliminary bioassay of plant extracts

The extraction of 500 g of *Centella asiatica* using methanol, ethanol and water as solvents resulted in methanol, ethanol and freeze dried aqueous extracts with percentage yields of 15% w/w (75 g), 11.2% w/w (56 g) and 4.6% w/w (23 g) respectively. The greater yields with methanol and ethanol as compared to water suggests that the plant's constituent compounds were more likely to exhibit greater solubility in polar than non-polar solvents.

The ACE inhibition assay of the crude organic and aqueous extracts was conducted according to modified method of Balasuriya and Rupasinghe (2012) as detailed in section 3.3.2.5. Concentrations of 1 mg/ml, 6 mg/ml and 10 mg/ml of the methanol, ethanol and aqueous extracts of *Centella asiatica* inhibited ACE activity in a dose dependent manner (table 4.1).

Inhibition of ACE was greatest with the methanol extract (85% at 10 mg/ml concentration), while the aqueous extract exhibited the least inhibition of enzyme activity (36.2% at 10 mg/ml concentration). The methanol and ethanol extracts exhibited inhibition significantly greater than that obtained with similar concentrations of the aqueous extract for all concentrations assessed. Although both methanol and ethanol extracts exhibited greater than 50% enzyme inhibition at the 1mg/ml concentration, inhibition was significantly lower than that observed with the same concentration of the standard ACEI captopril (98.5%) (table 4.1). Of the three extracts tested, the methanol and ethanol extracts demonstrated inhibition of ACE above 50% at all concentrations, and these extracts were subsequently subjected to the bioassay-guided fractionation process as described in section 3.3.2.5 of **Chapter 3**.

**Table 4. 1 Percentage inhibition of ACE activity by the methanol, ethanol and aqueous crude extracts of *Centella asiatica*.**

Plant Extract	% Inhibition		
	( 1mg/ml)	(6mg/ml)	(10mg/ml)
<b>Methanol Extract</b>	51.36	63.0	85.0
<b>Ethanol Extract</b>	55.2	66.4	78.6
<b>Aqueous Extract</b>	18.2	25.7	36.2
<b>Captopril</b>	98.5	-	-

#### **4.2 Bioassay guided fractionation of the methanol crude extract**

The fractionation of the methanol extract using silica gel column chromatography, yielded 8 fractions (M<sub>1</sub>– M<sub>8</sub>), which were assayed for ACE inhibition activity using the assay protocol described in section 3.3.2.5. Fractions M<sub>1</sub> and M<sub>2</sub> produced no inhibition of enzyme activity when tested for inhibition of ACE activity at the 1 mg/ml concentration, while fractions M<sub>3</sub>, M<sub>6</sub>, M<sub>7</sub> and M<sub>8</sub> all produced inhibition of enzyme activity below 50% (table 4.2). Fraction M<sub>4</sub>

was the most active (60.6 % inhibition), followed by fraction M<sub>5</sub> (53.3 % inhibition) at the 1 mg/ml concentration (table 4.2). There was an observed increase in the percentage inhibition of ACE by both fractions (M<sub>4</sub> and M<sub>5</sub>) compared to that obtained with the crude methanol extract (51.36% Inhibition) at a similar concentration. This increase could be attributed to concentration of active components as inactive diluents are fractionated out. Due to the greater inhibitory activity observed with fractions M<sub>4</sub> and M<sub>5</sub>, they were chosen for NMR analysis and further HPLC purification to isolate active compound.

**Table 4. 2 Percentage inhibition of ACE activity by fractions of the methanol extract of *Centella asiatica*.**

Column Fractions	% ACE Inhibition (1 mg/ml)	Percentage Yield
M <sub>1</sub>	-	-
M <sub>2</sub>	-	0.23
M <sub>3</sub>	0.5	1.6
M <sub>4</sub>	60.6*	1.0
M <sub>5</sub>	53.3*	0.79
M <sub>6</sub>	20.1	0.38
M <sub>7</sub>	28.3	53.37
M <sub>8</sub>	30.4	29.79

\* Active fractions subjected for further purification.

#### 4.2.1. HPLC purification of active fractions (M<sub>4</sub> and M<sub>5</sub>)

Fractions M<sub>4</sub> and M<sub>5</sub> were purified using a normal phase preparative HPLC according to the method described in sections 3.3.3.4.1 and 3.3.3.4.2.

HPLC fractionation of fraction M<sub>4</sub> (150 mg) resulted in 5 fractions (M<sub>4A</sub>, M<sub>4B</sub>, M<sub>4C</sub>, M<sub>4D</sub> and, M<sub>4E</sub>) (see appendix XXXII for chromatogram), which were subjected to the ACEI bioassay at a concentration of 1mg/ml. Fraction M<sub>4B</sub> exhibited the greatest ACE inhibition of 65% followed by M<sub>4C</sub> at 20.5%. The other fractions showed ACE inhibition less than 15% (table 4.3). Apart from the ethyl acetate wash, M<sub>4B</sub> and M<sub>4C</sub> showed the highest yield of 12.67% and 12.4% respectively in relation to the 150mg starting material (M<sub>4</sub>). Further bioassays and NMR spectroscopy focused on fraction M<sub>4B</sub> as the fraction with the greatest ACE inhibition (see appendix II for NMR spectra).

**Table 4. 3 Percentage inhibition of ACE activity by sub-fractions of fraction M<sub>4</sub> of the methanol extract of *Centella asiatica*.**

Fractions	Retention time	% Yield	% ACE inhibition (1mg/ml)
M <sub>4A</sub>	11.383	8.53	3.2
M <sub>4B</sub> *	13.386	12.67	65
M <sub>4C</sub>	15.241	6.6	20.5
M <sub>4D</sub>	17.182	5.4	10
(M <sub>4E</sub> ) Ethyl acetate wash	-	22.23	15
Fractions with no peak	-	37.47	12.36

\*Most Active Fraction identified as pure compound

HPLC fractionation of fraction M<sub>5</sub> (115mg) yielded 5 fractions (M<sub>5A</sub>, M<sub>5B</sub>, M<sub>5C</sub>, M<sub>5D</sub> and M<sub>5E</sub>) (see appendix XXXIII for chromatogram). ACEI bioassay at a concentration of 1mg/ml revealed fraction M<sub>5B</sub> to exhibit the highest ACE inhibition of 70.5% followed by M<sub>5C</sub> (23.2%). Fraction M<sub>5A</sub> exhibited no inhibition of enzyme activity, while all other fractions showed ACE

inhibition less than 15% (table 4.4). Further bioassays and NMR spectroscopy focused on M<sub>5B</sub> as the fraction with the greatest ACE inhibition.

**Table 4. 4 Percentage inhibition of ACE activity by sub-fractions of fraction M<sub>5</sub> of the methanol extract of *Centella asiatica*.**

Purified Fractions	Retention time	% Yield	% ACE inhibition (1mg/ml)
M <sub>5A</sub>	9.150	5.04	-
M <sub>5B</sub> *	10.804	11.02	70.5
M <sub>5C</sub>	11.453	4.2	23.2
M <sub>5D</sub>	13.713	8.17	10.4
M <sub>5E</sub> (Ethyl acetate Wash)	-	20.57	7
Fractions with no peak	-	40.26	5

\* Most active Fraction identified as pure compound

### 4.3. Bioassay guided fractionation of the ethanol extract

NMR profiling of the ethanol extract of *Centella asiatica* revealed the presence of fatty acids. Prior to the fractionation process, the extract was defatted according to the method described in section 3.3.3.5.1.

Defatting yielded methanol (550.6mg) and hexane (848.4mg) extracts which were subjected to ACE inhibition activity assay. The hexane extract exhibited a greater inhibition of ACE activity (51%) compared to the methanol extract (45 %), and so was subjected to further fractionation to isolate active compound/s.

Fractionation of the hexane extract yielded 5 fractions (F<sub>A</sub> - F<sub>E</sub>), with Fraction F<sub>A</sub> exhibiting the highest ACE inhibition (49 %) (Table 4.5). Fraction A was subjected to further purification to isolate active compound(s)

**Table 4. 5 Percentage Inhibition of ACE activity by Hexane extract fractions**

<b>Fractions</b>	<b>% ACE Inhibition</b>
<b>F<sub>A</sub></b>	49
<b>F<sub>B</sub></b>	12.5
<b>F<sub>C</sub></b>	21
<b>F<sub>D</sub></b>	8.1
<b>F<sub>E</sub></b>	5.2

#### 4.3.1. Purification of fraction F<sub>A</sub>

Fraction F<sub>A</sub> was purified using normal phase HPLC (see section 3.3.3.5.2). Four peaks eluting at 9.456, 11.076, 14.856, and 18.440 minutes, and the ethyl acetate wash were collected, yielding sub-fractions F<sub>A1</sub> - F<sub>A5</sub>. ACE inhibition assay at a concentration of 1mg/ml produced the greatest inhibition (39%) with F<sub>A3</sub> (see table 4.6). Fraction F<sub>A1</sub> and F<sub>A5</sub> exhibited no ACE inhibition activity while F<sub>A2</sub> and F<sub>A4</sub> showed less than 20% inhibition.

**Table 4. 6 Percentage Inhibition of ACE activity by HPLC fractions of Hexane extract.**

<b>HPLC Fractions</b>	<b>Retention Time</b>	<b>% Yield</b>	<b>% ACE inhibition</b>
<b>F<sub>A1</sub></b>	<b>9.456</b>	<b>5.2</b>	-
<b>F<sub>A2</sub></b>	<b>11.076</b>	<b>8.5</b>	<b>4.2</b>
<b>F<sub>A3</sub></b>	14.856	15.6	39
<b>F<sub>A4</sub></b>	18.440	18.1	15.8
<b>F<sub>A5</sub> (ethyl acetate wash)</b>	-	43.8	-

NMR analysis of F<sub>A3</sub> indicated that it consisted of one compound. However, some difficulty was experienced in elucidating the complete structure. The compound was suspected to be an

ester of coumaric acid, hence the decision to synthesize the esters of a synthetic coumaric acid to enable the identification of the purified compound through the comparison of its NMR spectra to that of the synthesized compounds.

#### 4.3.1.1 *Synthesis of methyl, ethyl and propyl ester of coumaric acid*

Three esters of coumaric acid were successfully synthesized with yields of 95.75% (methyl hydroxycinnamate), 85.2% (ethyl hydroxycinnamate) and 51.67% (propyl hydroxycinnamate) (see appendix I for synthesis pathway). The NMR profile of the three compounds were compared to isolated compound F<sub>A3</sub> to fully elucidate the structure (see appendix XXVIII – XXX for NMR data), and the isolated compound identified as ethyl hydroxycinnamic acid.


ACE activity was conducted on the synthesized ethyl hydroxycinnamic acid due to insufficient quantities of the isolated compound. The synthesized ethyl hydroxycinnamate (1 mg/ml) exhibited low ACE inhibitory activity of 15.2%, significantly lower than the inhibition produced by the compounds isolated from the methanol extract. The synthesized compound was not pursued further owing to its low ACE inhibition and as well as insufficient quantities of the isolated compound.

### **4.4 Structural elucidation and characterization of the isolated compounds**

Column chromatographic separation of the methanol extract yielded fractions M<sub>4</sub> and M<sub>5</sub> with significant ACE inhibition activity. The fractions were further purified using HPLC to produce M<sub>4B</sub>, from fraction M<sub>4</sub>, and two compounds M<sub>5B</sub> and M<sub>5C</sub> from fraction M<sub>5</sub>. NMR, MS, IR and UV Spectroscopy were used as tools for elucidation and characterization of the isolated compounds, comparable literature data of the isolated compounds were thoroughly searched using Scifinder, Molbase and PubChem. The results are presented below.

#### **4.4.1. Structural elucidation and characterization of compound M<sub>4B</sub> (Kaempferol)**

M<sub>4B</sub> (19 mg, 12.7% w/w yield) was a yellow powder with good solubility in ethanol, methanol, dimethyl sulfoxide, and poor solubility in water, and a melting point of 277 °C. The polar solubility of the isolated compound could be related to the initial solvent of extraction (methanol). The UV spectrum showed  $\lambda_{\text{max}}$  at 265 and 366 nm suggestive of the presence of flavonol bond (see appendix VIII). The <sup>1</sup>H-NMR (400MHz, CD<sub>3</sub>OD,) showed 2 peaks at  $\delta$  6.11 (<sup>1</sup>H, d,  $J = 1.87$  Hz) and 6.30 ppm (<sup>1</sup>H, d,  $J = 2.34$  Hz) consistent with the meta protons H-6 and H-8 on A-ring and an AA'BB' system at 7.99 (2H, d,  $J = 8.9$  Hz, H-2', 6') and 6.79 (2H, d,  $J = 8.87$  Hz, H-3', 5') corresponding to the protons on B-ring. <sup>13</sup>C NMR (100 MHz, CD<sub>3</sub>OD,) ppm 146.47 (s, C-2), 135.1 (s, C-3), 176.37 (s, C-4), 161.26 (s, C-5), 97.86 (d, C-6), 164.17 (s, C-7), 93.1 (d, C-8), 156.86 (s, C-9), 122.33 (s, C-1'), 129.28 (d, C-2'), 115.48 (d, C-3'), 159.16 (C-4'), 121.9 (d, C-5'), 134.7(d, C-6'), 109.9 (d, C-1''), 83.5 (d, C-2''), 78.8 (d, C-3''), 88.6 (d, C-4''), 62.4 (t, C-5'') (see appendix III and IV). The HRMS yielded a quasi-molecular ion peak [M-H]<sup>-</sup> at  $m/z$  285 and [M+H]<sup>+</sup> at  $m/z$  287 and corresponded with the molecular formulae C<sub>15</sub>H<sub>10</sub>O<sub>6</sub>, and a molecular mass of 286.04 g/mol (See appendix X). The compound was elucidated as **3,4',5,7-Tetrahydroxyflavone** (kaempferol) (see figure 4.1) by analysis of the various spectral data (MS, <sup>1</sup>H-, <sup>13</sup>C-NMR), and by comparison to literature values (Aisyah *et al.*, 2017)(**table 4.7**).




**Figure 4. 1 Structure of 3,4',5,7-Tetrahydroxyflavone** (kaempferol)

**Table 4.7 NMR Spectroscopic data of 3,4',5,7-Tetrahydroxyflavone** (kaempferol)

Carbon Position	$\delta_H$ (ppm)	Mult.	$J$ (Hz)	$\delta_C$ (ppm)	Type	HMBC	Literature Values ( $\delta_H$ ) (Aisyah <i>et al.</i> , 2017)	Literature Values ( $\delta_C$ ) (Aisyah <i>et al.</i> , 2017)
2				146.64	C			146.8
3					C			
4				176.37	C			176.6
5				161.26	C			162.3
6	6.11	d	1.87	97.86	CH	C8	6.28	99.2
7				164.17	C			164.9
8	6.30	d	2.34	93.1	CH	C6	6.52	94.4
9				156.86	C			157.7
10				103.15	C			104.1
1'				122.33	C			123.3
2'	7.99	d	8.90	129.28	CH	C6'	8.04	125.9
3'	6.79	d	8.87	115.48	C		6.95	116.3
4'	-----			159.16	C			160.1
5'	6.79	d	8.87	115.48	C	C1'	6.95	116.3
6'	7.99	d	8.90	129.28	CH	C2', C4', C2	8.04	125.9

#### 4.4.2 Structural elucidation and characterization of compound M<sub>5B</sub> (quercetin)

M<sub>5B</sub> (5.5mg 4.8% w/w Yield) was a yellow powder with good solubility in ethanol, methanol, dimethyl sulfoxide, and poor solubility in water, and with a melting point of 313 °C. The UV spectrum showed  $\lambda$  max at 256 and 370 nm, suggesting the presence of a flavonol bond (see appendix XVII). The IR spectrum, recorded in the 4000 to 600 cm<sup>-1</sup> range, was typical of a flavonoid structure. The spectra showed absorption bands with maxima at the following frequencies ( $V_{\max}$ ) 3400.79, 2925, 1607.46, 1520.49, 1362.89. The peaks at 3400.79 and 2925 are characteristic of the aromatic ring structure and absorption maxima at 1607 and 1520 indicating the presence of quinoid structure and –C=C- bonds, respectively. The absorption maximum at 1362 cm<sup>-1</sup> is due to phenolic OH groups (see appendix XIX). The <sup>1</sup>H-NMR spectrum showed 2 characteristic peaks at  $\delta$ 6.20 (1H, d,  $J$  = 2.0 Hz) and 6.41 ppm (1H, d,  $J$  = 2.0 Hz) consistent with the meta protons H-6 and H-8 on A-ring and implying the presence of a 5,7-dihydroflavonol in an A ring, which connects to no glycosides. Peaks at 7.55 (1H, d,  $J$  = 2.0 Hz, 8.4 Hz, H-6'), 7.68(1H, d,  $J$  = 2.2 Hz, H-2'), and 6.88 (1H, d,  $J$  =8.6 Hz, H-5') corresponding to the catechol protons on B ring. The <sup>13</sup>C-NMR spectrum revealed <sup>13</sup>CNMR (100MHz CD<sub>3</sub>OD) 8 92.99 (C-8), 97.82 (C-6), 103.11 (C-10), 115.48 (C-3', C-5'), 122.73 (C-1'), 129.28 (C-2', C-6'), 135.38 (C-3), 147.37 (C-2), 156.83(C-9), 147.7 (C-4'), 161.12 (C-5), 164.17 (C-7), 175.93 (C-4) (see appendix XII- XIII). The HRMS yielded a molecular peak [M-H]<sup>-</sup> at  $m/z$  301 and [M+H]<sup>+</sup> at  $m/z$  303 and corresponded to the molecular formulae C<sub>15</sub>H<sub>10</sub>O<sub>7</sub>, and a molecular mass 302.05g/mol (see appendix XVIII).The compound was elucidated as **3,5,7,3',4'- pentahydroxyflavone** (quercetin) by analysis of the various spectral data (MS, <sup>1</sup>H-, <sup>13</sup>C-NMR), and by comparing to literature values(Aisyah *et al.*, 2017; Cao, Wan, Yu, Zhou, and Tian, 2011) (figure 4.8).



**Figure 4. 2 Structure of 3,5,7,3',4'- pentahydroxyflavone (quercetin)**




**Table 4. 8 NMR Spectroscopic data of 3,5,7,3',4'- pentahydroxyflavone**

Carbon Position	$\delta_H$ (ppm)	Mult	$J$ (Hz)	$\delta_C$ (ppm)	Type	HMBC	Literature Values( $\delta_H$ ) (Aisyah <i>et al.</i> , 2017)	Literature Values( $\delta_C$ ) (Aisyah <i>et al.</i> , 2017)
2				147.37	C			147.7
3				135.83	C			135.7
4				175.93	C			176.8
5				161.12	C			160.7
6	6.20	d	2.0	97.82	CH	C8	6.20	94.5
7				164.17	C			163.9
8	6.41	d	2.0	92.99	CH	C6	6.40	94.5
9				156.83	C			156.1
10				103.11	C			103.0
1'				122.73	C			121.9
2'	7.68	d	2.2	114.60	CH	C6'	7.65	115.0
3'				145.20	C			145.0
4'				147.7	C			145.8
5'	6.88	d	8.6	116.05	C	C1', C3'	6.88	115.6
6'	7.55	d	8.4	129.68	CH	C2',C4', C2	7.50	124.5

#### 4.4.3 Structural elucidation and characterization of compound M<sub>5C</sub> (4 Hydroxybenzoic acid)

M<sub>5C</sub> was a white powder soluble in ethanol and methanol, and slightly soluble in water, with a melting point of 214.5°C. The UV spectrum showed  $\lambda$  max at 256 and 369 (see appendix XXVI). The NMR spectrum shows <sup>1</sup>H-NMR (MeOH-d, 400 MHz)  $\delta$  7.78 (2H, dd, J = 8.75,1.96 Hz, H-2,6), 6.72 (2H, dd, J = 8.78,1.9.2 Hz, H-3,5). <sup>13</sup>C-NMR (CD<sub>3</sub>OD, 100 MHz)  $\delta$ C: 168.18 (COOH), 163.35 (C-4), 131.59 (C-2,6), 121.29 (C-1), 114.91(C-3,5) (see appendix XXI - XXII). The HRMS yielded a molecular peak [M-H]<sup>-</sup> at  $m/z$  137 and [M+H]<sup>+</sup> at  $m/z$  139 and corresponded to the molecular formulae C<sub>7</sub>H<sub>6</sub>O<sub>3</sub>, and a molecular mass 138.02 g/mol (see appendix XXVII). The compound was elucidated as 4-Hydroxybenzoic acid (figure 4.13) by analysis of the spectral data and comparison to literature data(Lee *et al.*, 2011)



**Figure 4. 3 Structure of 4-Hydroxybenzoic acid**

**Table 4. 9 NMR Spectroscopic data of 4-Hydroxybenzoic acid**

Carbon atom	$\delta_H$ (ppm)	Mult	$J$ (Hz)	Type	$\delta_C$ (ppm)	Literature Values( $\delta_H$ ) (Lee <i>et al.</i> , 2011)	Literature Values( $\delta_C$ ) (Lee <i>et al.</i> , 2011)
1				C 1	121.29		122.9
2	7.78	dd	8.75	C 2	131.59	7.88	133.2
3	6.72	dd	8.78	C 3	114.91	6.82	116.2
4				C 4	163.35		163.5
5	6.72	dd	8.78	C 5	114.91	6.82	116.2
6	7.78	dd	8.75	C 6	131.59	7.88	133.2
7				- COOH	168.18		170.2

#### 4.5 Determining the IC<sub>50</sub> of captopril

The IC<sub>50</sub> value of an inhibitor is dependent upon inhibition mechanisms and assay conditions used and may vary from one laboratory to another, but is useful for comparing activities within a set of determinations (**Bush, 1983**).


IC<sub>50</sub> of captopril was determined and compared to the literature value of 23nM (**Ondetti, 1988**).

Table 4.5 shows the percentage ACE inhibition by captopril at various concentrations.

Captopril in this study exhibited a higher IC<sub>50</sub> of 85.76nM in comparison to the literature value of 23nM, but was still significant in relation the isolated compounds.

**Table 4.10 IC<sub>50</sub> of Reference standard (Captopril)**

Concentration of Captopril in assay (ng/ml)	% ACE Inhibition
6,25	27
12,5	45
25	60
50	65,8
100	78
200	91,3



**Figure 4. 4 IC<sub>50</sub> and percentage inhibition of ACE by captopril**


#### 4.6 ACE inhibition and IC<sub>50</sub> of the isolated compounds

The ACE activity of the compounds were assayed based on the fluorimetric determination of the amount of histidyl-leucine (His-Leu) cleaved from the enzyme substrate, hippuryl-histidyl-leucine (Hip-His-Leu). Detection of fluorescence intensity was aided by a fluorescence compound o-phthaldialdehyde (OPA) that binds to His-Leu. The ACE bioassay was conducted on the isolated compounds (3,4',5,7-Tetrahydroxyflavone and 3,5,7,3',4'-Pentahydroxyflavone) that exhibited significant inhibitory effect on ACE activity.

Both compounds exhibited a concentration dependent inhibition of ACE activity greater than 50% at 500 µg/ml concentration. 3,5,7,3',4'- pentahydroxyflavone expressed a greater ACE inhibition at all concentrations compared to 3,4',5,7-Tetrahydroxyflavone. 3,5,7,3',4'-pentahydroxyflavone at the highest dose of 1000 µg/ml, showed ACE inhibition of 73% compared to 3,4',5,7-Tetrahydroxyflavone (59.9%), and the 50% inhibitory concentration of 3,5,7,3',4'- pentahydroxyflavone (302.72 µg/ml) was 1.8 fold lower than that obtained with 3,4',5,7-Tetrahydroxyflavone (542.39 µg/ml), indicating that 3,5,7,3',4'- pentahydroxyflavone was a better ACE inhibitor than 3,4',5,7-Tetrahydroxyflavone (table 4.11 and figure 4.15). The IC<sub>50</sub> of the reference inhibitor (captopril) as determined in this study was 18.63 ng/ml

**Table 4. 11 Percentage inhibition of ACE activity by kaempferol and quercetin isolated from *C. asiatica*.**

% ACE Inhibition activity of isolated compounds		
Concentration (µg/ml)	3,4',5,7-tetrahydroxyflavone (M <sub>4B</sub> ) Kaempferol	3,5,7,3',4'- pentahydroxyflavone (M <sub>5B</sub> ) Quercetin
62.5	2	11.4
125	14.0	21.5
250	34.6	47.7
500	53.0	68.3
1000	59.9	73.0
IC 50	542.39 µg/ml	302.72 µg/ml




**Figure 4. 5 ACE inhibitory activity of 3,4',5,7-tetrahydroxyflavone (M<sub>4B</sub>) *Kaempferol* and 3,5,7,3',4'- pentahydroxyflavone (M<sub>5B</sub>) *Quercetin***

#### 4.7 DPPH radical scavenging assay

The DPPH assay method is a rapid, simple and inexpensive method to measure antioxidant capacity using DPPH (a stable free radical). The DPPH radical scavenging activity indicates hydrogen donating ability of compounds. The degree of reduction in absorbance measurement is indicative of scavenging potential of compounds. In this study, the DPPH radical scavenging capacity of the active compounds (3,4',5,7-Tetrahydroxyflavone and 3,5,7,3',4'-Pentahydroxyflavone) was evaluated.

All the test compounds exhibited dose dependent scavenging of DPPH radical. Compounds (3,4',5,7-Tetrahydroxyflavone and 3,5,7,3',4'-Pentahydroxyflavone) showed comparable antioxidant activity to the positive control (ascorbic acid). At a concentration of 125 µg/ml, 3,4',5,7-Tetrahydroxyflavone, 3,5,7,3',4'-Pentahydroxyflavone and ascorbic acid scavenged more than 50% of the free radicals. At double the concentration i.e. 250 µg/ml, 3,5,7,3',4'-Pentahydroxyflavone and ascorbic acid showed greater than 85% scavenging capacity compared to 3,4',5,7-Tetrahydroxyflavone at 75.4%. The highest scavenging activity of 91.2% was exhibited by 3,5,7,3',4'-Pentahydroxyflavone followed by 3,4',5,7-Tetrahydroxyflavone (81%) at the highest concentration of 500 µg/ml. The EC<sub>50</sub> value for 3,5,7,3',4'-Pentahydroxyflavone was determined as 84.83 µg/ml, which was 1.3 fold more potent than 3,4',5,7-Tetrahydroxyflavone (109.99 µg/ml) and 2.2 fold less potent than ascorbic acid (38.77 µg/ml). Figure 4.16 below depicts the scavenging activity of the isolated compounds and ascorbic acid.



**Figure 4. 6 DPPH radical scavenging activity of the isolated active compounds.**

**Table 4. 12 EC<sub>50</sub> of compounds**

Compound	EC <sub>50</sub> (µg/ml)	Regression Equation	Correlation coefficient (R <sup>2</sup> )
<b>3,4',5,7-Tetrahydroxyflavone(M<sub>4B</sub>)</b>	109.99	$y = 22,65\ln(x) - 56,463$	0,9746
<b>3,5,7,3',4'-Pentahydroxyflavone(M<sub>5B</sub>)</b>	84.83	$y = 25,031\ln(x) - 61,156$	0,9393
<b>Ascorbic acid</b>	38.77	$y = 19,65\ln(x) - 21,874$	0,9425

## Chapter 5

### Discussion and Conclusion

---

#### 5.1 Discussion

As part of our search for an alternative therapeutic approach for treatment of high blood pressure, as well as scientific validation of claimed therapeutic use of medicinal plants in hypertension. The methanol, ethanol and aqueous extracts of *Centella asiatica* were screened for their inhibitory effects on ACE. Amongst the extracts screened, methanol and ethanol extracts exhibited significant ACE inhibitory activity of 51.36% and 55.2% at 1mg/ml concentration respectively and 85% and 76.8% at 10mg/ml concentration respectively.

The extraction yield of the methanol and ethanol extracts in this study were 15% and 11.7% respectively compared to a much lower yield of 4.6% for the aqueous extract. This reflected an order of decreasing solvent polarity from water (lowest yield) to methanol (highest yield). Similar observation was made with regards to the ACE inhibitory effects of the crude extracts of methanol and ethanol which was greater compared to the aqueous extract. These differences in ACE inhibition by various extracts could be attributed to the solvent polarity and its ability to extract specific types of compound. In this study, less polar solvents such as methanol and ethanol seemed to extract ACE inhibitory compounds compared to the more polar aqueous solvent. A study by **Truong *et al* (2019)** showed methanol solvent extracted the highest level of phenolics, flavonoids, alkaloids and terpenoids compared to aqueous and ethanol solvent, aqueous solvent was said to show high efficiency in extraction of phenolic but not flavonoids, alkaloids and terpenoids. For this study, the aqueous extract exhibited a low yield and poor ACEI activity possibly due to poor solubility of the ACE inhibitory compounds in aqueous solvent.

The bioassay guided fractionation of the methanol and ethanol extracts afforded three pure compounds. These compounds were identified as 3,4',5,7-tetrahydroxyflavone (Kaempferol), 3,5,7,3',4'- pentahydroxyflavone (Quercetin), and the inactive 4 Hydroxybenzoic acid. The structures of the compounds were determined on the basis of their spectral data and compared to literature data as previously described. The two active compounds, 3,4',5,7-tetrahydroxyflavone, and 3,5,7,3',4'- pentahydroxyflavone, isolated from the methanol extract of *Centella asiatica* exhibited a dose dependent inhibition of ACE Activity, the 50% inhibitory concentration were determined as 542.39 µg/ml and 302.72 µg/ml respectively. This is the first report of ACE inhibitory effect of these compounds in *Centella asiatica*.

The isolated active compounds 3,4',5,7-tetrahydroxyflavone (Kaempferol), and 3,5,7,3',4'-pentahydroxyflavone (Quercetin) both consist of a flavonoid structure which is comprised of two phenyl rings (A and C rings) joined to a catechol group (B ring). Both compounds have a primary flavanone structure but 3,5,7,3',4'- pentahydroxyflavone (Quercetin) has an additional OH group at position 3' of the catechol group and therefore exhibited a 1.8-fold decrease in IC<sub>50</sub> relative to 3,4',5,7-tetrahydroxyflavone (Kaempferol). **Quin *et al* (2012)** in a structure activity relationship study of the ability of different flavonoids to inhibit ACE activity confirmed that the presence of the catechol group in the B-ring (3',4'- dihydroxy) appears to be fundamental to achieving an increased ACE inhibitory activity, the study showed that the absence of the 3' hydroxyl group in apigenin caused a 57% reduction of ACEI activity at 100 mM relative to the luteolin. This presence of additional OH group on C3' possibly explain the lower IC<sub>50</sub> exhibited by 3,5,7,3',4'- pentahydroxyflavone. In addition, the study identified that the double bond between C2 and C3 at C-ring and keto group at the C4 carbon on the C=ring to be essential for inhibiting ACE activity. The two active compounds isolated in this study comprised of these key functional groups. The importance of these functional groups were further confirmed in a study by **Tsutsumi *et al* (1998)**, where docking results of both quercetin

and epicatechin showed that the absence of the keto group and the C2–C3 double bond shifts the zinc binding site to the 7-OH moiety resulting in a reduced ACEI activity. Both 3,4',5,7-tetrahydroxyflavone (Kaempferol), and 3,5,7,3',4'- pentahydroxyflavone (Quercetin) isolated in this study contained these key functional group and double bond. **Shukor *et al.*, (2013)** further confirmed the role of the catechol B ring group , and C2=C3 double bond in ACEI activity . In contrast to the active compounds, the inactive 4 Hydroxybenzoic acid lacked these basic functional structures/groups which could possibly explain the reason for its lack of ACE inhibitory effect. The 4 Hydroxybenzoic acid comprises a benzoic acid structure carrying a hydroxyl substituent at the C-4 of the benzene ring, and is considered a phenolic derivative of the benzoic acid. Another important component of structure – ACE activity relationship is ACE, a zinc-containing peptidyl dipeptide hydrolase (**Becker and Scholkens, 1987**). Binding to ACE sub site is considered essential for ACEI activity. The active site of ACE consists of three parts; a carboxylate binding functionality such as the guanidium group of Arg, a pocket that accommodates the side chain of c-terminal amino acid residues, and a zinc ion. The zinc ion coordinates to the carbonyl of the penultimate peptide bond of the substrate, whereby the carbonyl group becomes polarized and is subjected to a nucleophilic attack(Oh *et al.*, 2004). Some flavonoids are suggested to show *in vitro* activity via the generation of chelate complexes within the active centre of ACE. Hydroxyl groups are suggested to be structural moieties that chelates zinc ions thus inactivating ACE activity(**Chen and Lin, 1992**). The hydroxyl group on B ring catechol group of 3,4',5,7-tetrahydroxyflavone (Kaempferol), and 3,5,7,3',4'-pentahydroxyflavone (Quercetin) can make charge-charge interactions with the active site  $Zn^{2+}$  of ACE thus exhibiting ACE inhibitory effect. Interestingly, **Quin *et al.*, 2012** showed that charge-charge interaction with 3' hydroxyl is stronger than that of with the 4' interaction. Which could explain why kaempferol showed a lower ACE inhibition effect relative to

quercetin. Therefore, these structural features and hydroxyl groups are considered a putative zinc-chelating sites.

The renin angiotensin system has been implicated in inducing oxidative stress in vascular and endothelial smooth muscle cells and may be an important mechanism also participating in the pathophysiology of hypertension. Studies have suggested that free radicals might mediate the pathogenesis of vascular hyper permeability and cellular damage associated with oxidative stress (**Dhalla et al., 2000**). Oxidative stress has also been implicated as a risk factor for cardiovascular disease through its triggering of the atherogenic processes that hardens and narrows arteries thus impeding the flow of blood around the body. In a study by **Wilson** (1990), rats made acutely hypertensive by administering angiotensin II showed severe and extensive endothelial and smooth muscle lesions that were inhibited upon antioxidant treatment. It could be said that an important synergistic role exists between antioxidant and ACE inhibition in the treatment of hypertension and cardiovascular disease. Flavonols act as prominent antioxidants in biological systems. The antioxidant capacity of the isolated compounds from *Centella asiatica* was determined to be comparable to the reference compound (Ascorbic acid). 3,4',5,7-tetrahydroxyflavone (Kaempferol), and 3,5,7,3',4'- pentahydroxyflavone (Quercetin) scavenged 50% of the DPPH radical at 84.83 and 109.99 µg/ml compared to the reference standard (38.77 µg/ml). A study by **Okawa et al** (2001) showed quercetin (8 µM) to exhibit stronger DPPH radical scavenging activity than kaempferol (41.2 µM). The structural properties once more play a role in the biological activity of these compounds, the presence of a dihydroxy group at C-3' and 4' (catechol) position in the B-ring of quercetin flavanol skeleton is suggested to enhance the DPPH radical scavenging activity.

In conclusion, the structure– function relationships can be useful for designing new ACE inhibitors based on phenolic compounds and further strategies such as glycosylation of C3 and C7 could confer greater ACE inhibitory activity. Although the ACEI activity of these

compounds do not reach the potency of drugs commonly used in the treatment of hypertension, the synergistic role of antioxidant and ACE inhibition properties can play a role in mild to moderate treatment of hypertension and a plant like *Centella asiatica*, serve as a source of dietary supplementation or considered as a naturally functional food.

## 5.2 Conclusion

The use of ACE inhibitors to treat hypertension, which is considered a major risk factor for cardiovascular disease remains a very important therapeutic strategy, but the manifestation of side effects has stirred up interest in alternative source of therapy hence the increasing interest in medicinal plant as source of safe and effective treatment. The aim of this study was to investigate the antihypertensive effect of *Centella asiatica* using an *in vitro* bioassay. The objectives were to evaluate the inhibition of ACE activity by crude extracts of *Centella asiatica* and to isolate and identify the ACE inhibitory compounds using bioassay guided fractionation. It was therefore hypothesized that *Centella asiatica* mediates its anti-hypertensive activity by means of ACE inhibition.

In this study, methanol, ethanol and aqueous extract were assayed for ACE activity. The active methanol and ethanol extract were subjected to a bioassay guided fractionation. Active fractions were further purified on a high performance liquid chromatography to yield the final pure compounds which were identified by means of NMR, MS, UV and IR spectroscopy. ACE Inhibition and antioxidant activity were conducted on the active compounds.

From the results obtained methanol extract yielded three compounds, 3,4',5,7-tetrahydroxyflavone (kaempferol), 3,5,7,3',4'- pentahydroxyflavone (quercetin), and 4-Hydroxybenzoic acid. Quercetin and kaempferol were determined to exhibit significant ACE inhibition as well as antioxidant activity.

These compounds have been previously reported to be present in *Centella asiatica*, but this is the first report of their ACE inhibitory effect in this plant. Though the compounds show less inhibition than the reference standard captopril, the synergistic effect of ACE inhibition and antioxidant activity is believed to play an important role in treatment of hypertension and cardiovascular disease. Furthermore, these compounds could be utilized as leads by synthesising glycosylated conjugates which has been shown to exhibit significant ACE inhibition activity. It is noteworthy that quercetin and kaempferol are commercialized products, therefore the results of this study highlights the need to be mindful of drug-drug interactions that might arise in hypertensive patients who are also on antihypertensive drugs. In other words, there is a possibility of an increased antihypertensive effect in patients concomitantly taking quercetin and kaempferol containing herbal products together with an antihypertensive drug

### **5.3 Limitations**

An *in vitro* assay was the only method used in this study, due to lack of access to animals, *in vivo* assay was not conducted. It is worthy of note that *in vitro* conditions/results does not necessarily reflect or translate to *in vivo* outcome.

### **5.4 Recommendation**

Future studies are required to investigate the *in vivo* effect of the isolated compounds and to synthesize conjugates of the compounds which can be examined for activity using *in vitro* and *in vivo* experiments.

## References

- Actis-Goretta, L., Ottaviani, J. I., & Fraga, C. G. (2006). Inhibition of angiotensin converting enzyme activity by flavanol-rich foods. *Journal of Agricultural and Food Chemistry*, 54(1), 229–234. <https://doi.org/10.1021/jf052263o>
- Aiko Morigwa, Katsuaki Kitabatake, Yoshinori Fujimoto, N. I. (1986). Angiotensin Converting Enzyme Inhibitory triterpenes from *Ganoderma lucidum*. *Chemical & Pharmaceutical Bulletin*, 34(7), 3025–3028. <https://doi.org/10.1248/cpb.37.3229>
- Aisyah, L. S., Yun, Y. F., Herlina, T., Julaeha, E., Zainuddin, A., & Nurfarida, I. (2017). Flavonoid Compounds from the Leaves of *Kalanchoe prolifera* and Their Cytotoxic Activity against P-388 Murine Leukimia Cells. *Natural Product Sciences*, 23(2), 139–145.
- Almeida, L. F. De, & Coimbra, T. M. (2019). When Less or More Isn't Enough: Renal Maldevelopment Arising From Disequilibrium in the Renin-Angiotensin System. *Frontiers in Paediatrics*, 7(July), 1–8. <https://doi.org/10.3389/fped.2019.00296>
- Alwan, A., Armstrong, T., Bettcher, D., & Branca, F. (2010). Global status report on noncommunicable diseases. Retrieved April 30, 2017, from [https://apps.who.int/iris/bitstream/handle/10665/44579/9789240686458\\_eng.pdf?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/44579/9789240686458_eng.pdf?sequence=1)
- Amos, C., & Ariei, Y. (1978). An Intramolecularly Quenched Fluorescent Tripeptide as a Fluorogenic Substrate of Angiotensin-I-Converting Enzyme and of Bacterial Dipeptidyl

Carboxypeptidase. *European Journal of Biochemistry*, 87, 265–273.

Anderson, G. H., & Moore, S. E. (2004). The Emerging Role of Dairy Proteins and Bioactive Peptides in Nutrition and Health Dietary Proteins in the Regulation of Food Intake and Body Weight in Humans. *Journal of Nutrition*, 134(4), 974–979.  
<https://doi.org/10.1128/JB.188.5.1691>

Atlas, S. A. (2007). The Renin-Angiotensin Aldosterone System: Pathophysiological Role and Pharmacologic Inhibition. *Journal of Managed Care Pharmacy*, 13(8 Supp B), 9–20. <https://doi.org/10.18553/jmcp.2007.13.s8-b.9>

Balasuriya, B. W. N., & Rupasinghe, H. P. V. (2011). Plant flavonoids as angiotensin converting enzyme inhibitors in regulation of hypertension. *Functional Foods in Health Disease*, 5(5), 172–188. Retrieved from  
<http://www.functionalfoodscenter.net/files/49461525.pdf>

Balasuriya, N., & Rupasinghe, H. P. V. (2012). Antihypertensive properties of flavonoid-rich apple peel extract. *Food Chemistry*, 135(4), 2320–2325.  
<https://doi.org/10.1016/j.foodchem.2012.07.023>

Bandara, M. S., Lee, E. L., & Thomas, J. E. (2011). Gotu Kola ( *Centella asiatica* L .): An Under-utilized Herb. *American Journal of Plant Science and Biotechnology*, 1–12.

Becker, R. H. ., & Scholkens, B. (1987). Ramipril : Review of Pharmacology. *Journal of Cardiology*, 59(10), D3–D11.

Bhat, R. (2014). Medicinal plants and traditional practices of Xhosa people in the Transkei region of Eastern Cape , South Africa. *Indian Journal of Traditional Knowledge*, 13(2), 292–298.

Bush, K. (1983). Screening and Characterization of Enzyme Inhibitors as Drug Candidates \*.

*The Squibb Institute for Medical Research*, 14(4), 689–708.

Cao, S., Wan, C., Yu, Y., Zhou, S., & Tian, S. (2011). Isolation and identification of phenolic compounds from *Gynura divaricata* leaves. *Pharmacognosy Magazine*, 7(26), 101.

<https://doi.org/10.4103/0973-1296.80666>

Cappuccio, F. P., & Miller, M. A. (2016). Cardiovascular disease and hypertension in sub-Saharan Africa : burden , risk and interventions. *Internal and Emergency Medicine*,

11(3), 299–305. <https://doi.org/10.1007/s11739-016-1423-9>

Chandrakant Katiyar, Arun Gupta, Satyajyoti Kanjilal, S. K. (2012). Drug discovery from plant sources: An integrated approach. *An International Quarterly Journal of Review in Ayurveda*, 33(10–19).

Chen, C. H., & Lin, J. Y. (1992). Inhibition of Angiotensin-I-Converting Enzyme by Tetrahydroxyxanthenes Isolated from *Tripterospermum lanceolatum*. *Natural Products*, 55(5), 691–695.

Chen, C. H., Lin, J. Y., Lin, C. N., & Hsu, S. Y. (1992). Inhibition of angiotensin-i-converting enzyme by tetrahydroxyxanthenes isolated from *tripterospermum lanceolatum*. *Journal of Natural Products*, 55(5), 691–695.

<https://doi.org/10.1021/np50083a025>

Chen, Z. Y., Peng, C., Jiao, R., Wong, Y. M., Yang, N., & Huang, Y. (2009). Anti-hypertensive nutraceuticals and functional foods. *Journal of Agricultural and Food Chemistry*, 57(11), 4485–4499. <https://doi.org/10.1021/jf900803r>

Cushman, D. W., & Cheung, H. S. (1971). Spectrophotometric assay and properties of the angiotensin-converting enzyme of rabbit lung. *Biochemical Pharmacology*, 20(7), 1637–1648. [https://doi.org/10.1016/0006-2952\(71\)90292-9](https://doi.org/10.1016/0006-2952(71)90292-9)

- Dhalla, N. S., Temsah, R. M., & Netticadan, T. (2000). Role of oxidative stress in cardiovascular diseases. *Journal of Hypertension*, 18(6), 655–673.
- Duncan, A. C. (1998). *An Investigation of Plants used in South Africa for the Treatment of Hypertension*. University of Natal.
- Dyubeni, L., & Buwa, L. V. (2012). An ethnobotanical study of plants used for the treatment of ear , nose and throat ( ENT ) infections in Nkonkobe Municipality , South Africa. *Journal of Medicinal Plant Research*, 6(14), 2721–2726.  
<https://doi.org/10.5897/JMPR10.869>
- Ekor, M. (2014). The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in Neurology*, 4 JAN(January), 1–10.  
<https://doi.org/10.3389/fphar.2013.00177>
- Elbl, G., & Wagner, H. (1991). A New Method for the in vitro Screening of Inhibitors of Angiotensin-Converting Enzyme (ACE), Using the Chromophore- and Fluorophore-Labelled Substrate, Dansyltriglycine. *Planta Medica*, 57(2), 137–141.
- Ettehad, D., Emdin, C. A., Kiran, A., Anderson, S. G., Callender, T., Emberson, J., ... Rahimi, K. (2016). Blood pressure lowering for prevention of cardiovascular disease and death: A systematic review and meta-analysis. *The Lancet*, 387(10022), 957–967.  
[https://doi.org/10.1016/S0140-6736\(15\)01225-8](https://doi.org/10.1016/S0140-6736(15)01225-8)
- Fatema, K., Zwar, N. A., Milton, A. H., & Ali, L. (2016). Prevalence of Risk Factors for Cardiovascular Diseases in Bangladesh : A Systematic Review and Meta-Analysis. *PLoS ONE*, 11(8), 1–14. <https://doi.org/10.1371/journal.pone.0160180>
- Fennell, C. W., Lindsey, K. L., McGaw, L. J., Sparg, S. G., Stafford, G. I., Elgorashi, E. E., ... Van Staden, J. (2004). Assessing African medicinal plants for efficacy and safety:

- Pharmacological screening and toxicology. *Journal of Ethnopharmacology*, 94(2–3), 205–217. <https://doi.org/10.1016/j.jep.2004.05.012>
- Gersh, B. J., Sliwa, K., Mayosi, B. M., & Yusuf, S. (2010). Novel therapeutic concepts: The epidemic of cardiovascular disease in the developing world: Global implications. *European Heart Journal*, 31(6), 642–648. <https://doi.org/10.1093/eurheartj/ehq030>
- Govindan, G., Sambandan, T. G., Govindan, M., Sinskey, A., Vanessendelft, J., Adenan, I., & Rha, C. K. (2007). A bioactive polyacetylene compound isolated from *Centella asiatica*. *Planta Medica*, 73(6), 597–599. <https://doi.org/10.1055/s-2007-981521>
- Gurib-Fakim, A. (2006). Medicinal plants: Traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine*, 27(1), 1–93. <https://doi.org/10.1016/j.mam.2005.07.008>
- Handa, S., Khanuja, S., Longo, G., & Rakesh, D. (2008). Extraction Technologies for Medicinal and Aromatic Plants. *United Nations Industrial Development Organization and International Centre for Science and High Technology*, 66.
- Hansen, K., Nyman, U., Smitt, U. W., Adersen, A., Gudiksen, L., Rajasekharan, S., & Pushpangadan, P. (1995). In vitro screening of traditional medicines for anti-hypertensive effect based on inhibition of the angiotensin converting enzyme (ACE). *Journal of Ethnopharmacology*, 48(1), 43–51. [https://doi.org/10.1016/0378-8741\(95\)01286-M](https://doi.org/10.1016/0378-8741(95)01286-M)
- Hanton, G. (2007). Preclinical Cardiac Safety Assessment of Drugs. *Drugs in R & D*, 8(4), 213–228.
- Higdon, J. (2005). Flavonoids [online] Linus Pauline institute, micronutrient information centre. Available at:

<https://lpi.oregonstate.edu/mic/dietary-factors/phytochemicals/flavonoids> [Accessed 30th April. 2017].

Holmquist, B., Bünning, P., & Riordan, J. F. (1979). A continuous spectrophotometric assay for angiotensin converting enzyme. *Analytical Biochemistry*, 95(2), 540–548.

[https://doi.org/10.1016/0003-2697\(79\)90769-3](https://doi.org/10.1016/0003-2697(79)90769-3)

Horton, R. (2013). NCDs : a challenge to sustainable human development. *The Lancet*, 381(9866), 510–511. [https://doi.org/10.1016/S0140-6736\(13\)60058-6](https://doi.org/10.1016/S0140-6736(13)60058-6)

Inamdar, P. K., Yeole, R. D., Ghogare, A. B., & Souza, N. J. De. (1996). Determination of biologically active constituents in *Centella asiatica*. *Journal of Chromatography*, 742(1–2), 127–130.

Israili, Z. H., & Hall, W. D. (1992). Cough and Angioneurotic Edema Associated with Angiotensin-converting Enzyme Inhibitor Therapy A Review of the Literature and Pathophysiology. *Annals of Internal Medicine*, 3(117), 234–242.

Jallapally, A., Addla, D., Bagul, P., Sridhar, B., Banerjee, S. K., & Kantevari, S. (2015). Design , synthesis and evaluation of novel 2-butyl-4-chloroimidazole derived peptidomimetics as Angiotensin Converting Enzyme ( ACE ) inhibitors. *Bioorganic and Medicinal Chemistry*, 23(13), 3526–3533. <https://doi.org/10.1016/j.bmc.2015.04.024>

Kamadyaapa, D. R. J. (2008). *Mechanisms of the cardiovascular effects of some medicinal plants: an experimental study*. PhD. University of Kwazulu Natal. South Africa.

Kannel, W. B., Vasan, R. S., & Levy, D. (2003). Is the Relation of Systolic Blood Pressure to Risk of Cardiovascular Disease Continuous and Graded , or Are There Critical Values ? *Hypertension*, 42(4), 453–456. <https://doi.org/10.1161/01.HYP.0000093382.69464.C4>

Kashmira, G., Jagruti, P., & Gajjar, A. K. (2010). Pharmacological Review on *Centella*

- asiatica : A Potential Herbal Cure-all. *Indian Journal of Pharmaceutical Sciences*, 72(5)(October), 546–556.
- Kearney, P. M., Whelton, M., Reynolds, K., Muntner, P., Whelton, P. K., & He, J. (2005). Global burden of hypertension : analysis of worldwide data. *The Lancet*, 365(9455), 2147–2223.
- Kinghorn, A. D. (2001). Pharmacognosy in the 21 st century\*. *Journal of Pharmacy and Pharmacology*, 53, 135–148. <https://doi.org/10.1211/0022357011775334>
- Kumar, S., & Pandey, A. K. (2013). Chemistry and biological activities of flavonoids: An overview. *The Scientific World Journal*, 1–16. <https://doi.org/10.1155/2013/162750>
- Lam, L. H. L., Himamura, T. S., Anabe, S. M., Shiyama, M. I., & Keda, H. U. (2008). Assay of Angiotensin I-converting Enzyme-inhibiting Activity Based on the Detection of 3-Hydroxybutyrate with Water-soluble Tetrazolium Salt. *Analytical Sciences*, 24(8), 1057–1060. <https://doi.org/10.2116/analsci.24.1057>
- Larson, A. J., Symons, J. D., & Jalili, T. (2012). Therapeutic Potential of Quercetin to Decrease Blood Pressure : Review of Ef fi cacy. *An International Review Journal*, 3, 39–46. <https://doi.org/10.3945/an.111.001271.studies>
- Lee, D. H., Kim, J. H., Park, J. S., Choi, Y. J., & Lee, J. S. (2004). Isolation and characterization of a novel angiotensin I-converting enzyme inhibitory peptide derived from the edible mushroom *Tricholoma giganteum*. *Peptides*, 25(4), 621–627. <https://doi.org/10.1016/j.peptides.2004.01.015>
- Lee, H., Cho, J., Moon, J., & Park, K. (2011). Isolation and Identification of Antioxidative Phenolic Acids and Flavonoid Glycosides from *Camellia japonica* Flowers. *Horticulture, Environment, and Biotechnology*, 52(3), 270–277.

<https://doi.org/10.1007/s13580-011-0157-x>

Light, M. E., Sparg, S. G., Stafford, G. I., & Staden, J. Van. (2005). Riding the wave : South Africa ' s contribution to ethnopharmacological research over the last 25 years. *Journal of Ethnopharmacology*, 100(1–2), 127–130. <https://doi.org/10.1016/j.jep.2005.05.028>

Lip, G. Y. H., Churchill, D., Beevers, M., Auckett, A., & Beevers, D. G. (1997). Angiotensin-converting-enzyme inhibitors in early pregnancy *Borrelia burgdorferi* infection in patients with suspected acute myocardial infarction. *Lancet (London, England)*, 350, 1446–1447.

Liu, W. J. H. (2011). *Traditional herbal medicine research methods : identification, analysis, bioassay, and pharmaceutical and clinical studies*. Hoboken, N.J. : John Wiley & Sons.

Loizzo, M. R., Said, A., Tundis, R., Rashed, K., Antonio, G., Hufner, A., & Menichini, F. (2006). Inhibition of Angiotensin Converting Enzyme ( ACE ) by Flavonoids isolated from *Ailanthus excelsa* ( Roxb ) ( Simaroubaceae ). *Phytotherapy Research*, 21(1), 32–36. <https://doi.org/10.1002/ptr>

López, P., Fauw, D. P. K. H. P., Mulder, P. P. J., Spanjer, M., Stoppelaar, J. De, Mol, H. G. J., & Nijs, M. De. (2018). Straightforward analytical method to determine opium alkaloids in poppy seeds and bakery products. *Food Chemistry*, 242, 443–450. <https://doi.org/10.1016/j.foodchem.2017.08.045>

Malgorzata, B., Jozef, K., & Grzegorz, B. (1997). Antioxidant and Prooxidant Properties of Captopril. *Free Radical Biology and Medicine*, 23(5), 729–735.

Niskanen, L., Hedner, T., Hansson, L., Lanke, J., & Niklason, A. (2001). Reduced cardiovascular morbidity and mortality in hypertensive diabetic patients on first-line therapy with an ACE inhibitor compared with a diuretic/ $\beta$ -blocker-based treatment

- regimen: A subanalysis of the captopril prevention project. *Diabetes Care*, 24(12), 2091–2096. <https://doi.org/10.2337/diacare.24.12.2091>
- Obikeze, K. C. (2009). *Cardiovascular effects of (13S)-9<sub>α</sub>, 13<sub>α</sub>-epoxylabda-6<sub>α</sub>(19), 15(14)diol dilactone, a diterpenoid isolated from the organic extract of Leonotis leonurus leaves, in anaesthetized normotensive rats*. PhD. University of The Western Cape, South Africa.
- Oh, H., Kang, D.-G., Kwon, J.-W., Kwon, T.-O., Lee, S.-Y., Lee, D.-B., & Lee, H.-S. (2004). Isolation of angiotensin converting enzyme (ACE) inhibitory flavonoids from *Sedum sarmentosum*. *Biological & Pharmaceutical Bulletin*, 27(12), 2035–2037. <https://doi.org/10.1248/bpb.27.2035>
- Oh, H., Kang, D.-G., Lee, S., Lee, Y., & Lee, H.-S. (2003). Angiotensin converting enzyme (ACE) inhibitory alkaloids from *Fritillaria ussuriensis*. *Planta Medica*, 69(6), 564–565. <https://doi.org/10.1055/s-2003-40659>
- Oh, H., Kang, D. G., Lee, S., & Lee, H. S. (2002). Angiotensin converting enzyme inhibitors from *Cuscuta japonica* Choisy. *Journal of Ethnopharmacology*, 83(1–2), 105–108. [https://doi.org/10.1016/S0378-8741\(02\)00216-7](https://doi.org/10.1016/S0378-8741(02)00216-7)
- Ojeda, D., Jiménez-ferrer, E., Zamilpa, A., Herrera-arellano, A., Tortoriello, J., & Alvarez, L. (2010). Inhibition of angiotensin convertin enzyme ( ACE ) activity by the anthocyanins delphinidin- and cyanidin-3- O -sambubiosides from *Hibiscus sabdariffa*. *Journal of Ethnopharmacology*, 127(1), 7–10. <https://doi.org/10.1016/j.jep.2009.09.059>
- Okawa, M., Kinjo, J., Nohara, T., & Ono, M. (2001). DPPH ( 1, 1-Diphenyl-2-Picrylhydrazyl ) Radical Scavenging Activity of Flavonoids Obtained from Some Medicinal Plants. *Biological & Pharmaceutical Bulletin*, 24(10), 1202–1205.

- Orhan, I. E. (2012). Centella asiatica ( L .) Urban : From Traditional Medicine to Modern Medicine with Neuroprotective Potential. *Evidence-Based Complementary and Alternative Medicine*, 1–8. <https://doi.org/10.1155/2012/946259>
- Ortiz-Salmeróna, E., Barón, C., & García-Fuentes, L. (1998). Enthalpy of captopril-angiotensin I-converting enzyme binding. *Febs Letters*, 435(2–3), 219–224.
- Pan, S., Zhou, S., Gao, S., Yu, Z., Zhang, S., Tang, M., ... Ko, K. (2013). New Perspectives on How to Discover Drugs from Herbal Medicines : CAM ' s Outstanding Contribution to Modern Therapeutics. *Evidence-Based Complementary and Alternative Medicine*, 1–25.
- Quin, M., Garcia-vallve, S., & Pujadas, G. (2012). Inhibition of Angiotensin-Converting Enzyme Activity by Flavonoids : Structure-Activity Relationship Studies. *PLoS ONE*, 7(11), 1–11. <https://doi.org/10.1371/journal.pone.0049493>
- Ruszkowski, P., & Bobkiewicz-kozlowska, T. (2014). Natural Triterpenoids and their Derivatives with Pharmacological Activity Against Neurodegenerative Disorders. *Mini-Reviews in Organic Chemistry*, 11(3), 307–315.
- Sairam, K., Rao, C. V., & Goel, R. K. (2001). Effect of Centella asiatica Linn on physical and chemical factors induced gastric ulceration and secretion in rats. *Indian Journal of Experimental Biology*, 39(2), 137–142. Retrieved from <http://www.scopus.com/inward/record.url?eid=2-s2.0-0035112251&partnerID=40&md5=9bcea7feadf99e6e052f367b4af4a048>
- Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K. M., & Yoga Latha, L. (2011). Extraction, isolation and characterization of bioactive compounds from plants' extracts. *African Journal of Traditional, Complementary and Alternative Medicines*, 8(1), 1–10. <https://doi.org/10.4314/ajtcam.v8i1.60483>

- Schwager, S. L., Carmona, A. K., & Sturrock, E. D. (2006). A high-throughput fluorimetric assay for angiotensin I-converting enzyme. *Nature Protocols*, *1*(4), 1961–1964.  
<https://doi.org/10.1038/nprot.2006.305>
- Shen, Q., Zhang, B., Xu, R., Wang, Y., Ding, X., & Li, P. (2010). Anaerobe Antioxidant activity in vitro of the selenium-contained protein from the Se-enriched *Bifidobacterium animalis* 01. *Anaerobe*, *16*(4), 380–386.  
<https://doi.org/10.1016/j.anaerobe.2010.06.006>
- Shukla, a, Rasik, a M., Jain, G. K., Shankar, R., Kulshrestha, D. K., & Dhawan, B. N. (1999). In vitro and in vivo wound healing activity of asiaticoside isolated from. *Journal of Ethnopharmacology*, *65*(1), 1–11.
- Shukor, N. Al, Camp, J. Van, Gonzales, G. B., Staljanssens, D., Struijs, K., Zotti, M. J., ... Smagghe, G. (2013). Angiotensin-Converting Enzyme Inhibitory Effects by Plant Phenolic Compounds: A Study of Structure Activity Relationships. *Journal of Agricultural and Food Chemistry*, *61*(48), 11832–11839.
- Sica, D. A. (2007). Angiotensin-Converting Enzyme Inhibitor Use in the Year 2005. *The Journal of Clinical Hypertension*, *7*(s8), 8–11. <https://doi.org/10.1111/j.1524-6175.2005.04598.x>
- Siddiqui, B. S., Aslam, H., Ali, S. T., Khan, S., & Begum, S. (2007). Chemical constituents of *Centella asiatica*. *Journal of Asian Natural Products Research*, *9*(4), 407–414.  
<https://doi.org/10.1080/10286020600782454>
- Sleight, P. (2000). The HOPE Study ( Heart Outcomes Prevention Evaluation ). *Journal of Renin-Angiotensin-Aldosterone System*, *1*(1), 18–20.
- Somova, L. O., Nadar, A., Rammanan, P., & Shode, F. O. (2003). Cardiovascular ,

antihyperlipidemic and antioxidant effects of oleanolic and ursolic acids in experimental. *Phytomedicine*, *10*(2–3), 115–121.

Sondhi, N., Bhardwaj, R., Kaur, S., Chandel, M., Kumar, N., & Singh, B. (2010). Inhibition of H<sub>2</sub>O<sub>2</sub>-induced DNA damage in single cell gel electrophoresis assay (comet assay) by castasterone isolated from leaves of centella asiatica. *Health*, *02*(06), 595–602.  
<https://doi.org/10.4236/health.2010.26088>

Steffensen, F. H., Nielsen, G. L., Sørensen, H. T., Olesen, C., & Olsen, J. (1998). Pregnancy outcome with ACE-inhibitor use in early pregnancy. *Lancet (London, England)*, *351*(9102), 596. [https://doi.org/10.1016/S0140-6736\(05\)78584-6](https://doi.org/10.1016/S0140-6736(05)78584-6)

Subban, R., Veerakumar, A., Manimaran, R., Hashim, K. M., & Balachandran, I. (2008). Two new flavonoids from *Centella asiatica* (Linn.). *Journal of Natural Medicines*, *62*(3), 369–373. <https://doi.org/10.1007/s11418-008-0229-0>

Taylor, J. L.S., Rabe, T., McGaw, L. J., Jäger, A. K., & Van Staden, J. (2001). Towards the scientific validation of traditional medicinal plants. *Plant Growth Regulation*, *34*(1), 23–37. <https://doi.org/10.1023/A:1013310809275>

Taylor, Joeslyn Leanda Susan. (1999). *An Investigation into the biology and medicinal properties of Eucomis species*. Phd. University of Natal.

Truong, D., Nguyen, D. H., Thuy, N., Ta, A., Bui, A. V., & Do, T. H. (2019). Evaluation of the Use of Different Solvents for Phytochemical Constituents, Antioxidants, and In Vitro Anti-Inflammatory Activities of *Severinia buxifolia*. *Journal of Food Quality*, *2019*, 1–9.

Tsutsumi, Y., Shimada, A., Miyano, A., Nishida, T., & Mitsunaga, T. (1998). In vitro screening of angiotensin I-converting enzyme inhibitors from Japanese cedar (

Cryptomeria japonica ). *Japan Wood Science*, 44, 463–468.

W. Aronow. (2012). Treatment of systemic hypertension. *American Journal of Cardiovascular Disease*, 2(3), 160–170. [https://doi.org/10.1016/0002-9149\(87\)90535-2](https://doi.org/10.1016/0002-9149(87)90535-2)

Weir, M. R., & Dzau, V. J. (1999). The Renin-Angiotensin-Aldosterone System: A Specific Target for Hypertension Management. *American Journal of Hypertension*, 12(12), 205S-213S. [https://doi.org/10.1016/S0895-7061\(99\)00103-X](https://doi.org/10.1016/S0895-7061(99)00103-X)

WHO. (1999). *WHO monographs on selected medicinal plants*. World Health Organization (Vol. 1).

WHO. (2010). Global status report on noncommunicable diseases. *World Health Organization*. Retrieved from [https://apps.who.int/iris/bitstream/handle/10665/44579/9789240686458\\_eng.pdf?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/44579/9789240686458_eng.pdf?sequence=1)

Wilson, S. K. (1990). Role of Oxygen-Derived Free Radicals in Acute Angiotensin II-Induced Hypertensive Vascular Disease in the Rat. *Circulation Research*, 66(3), 722–734.


Wood, R. (1995). Bronchospasm and cough as adverse reactions to the ACE inhibitors captopril, enalapril and lisinopril. A controlled retrospective cohort study. *British Journal of Clinical Pharmacology*, 39(3), 265–270. <https://doi.org/10.1111/j.1365-2125.1995.tb04447.x>

Xie, Y., & Zhang, W. (2012). Antihypertensive activity of Rosa rugosa Thunb. flowers: Angiotensin i converting enzyme inhibitor. *Journal of Ethnopharmacology*, 144(3), 562–566. <https://doi.org/10.1016/j.jep.2012.09.038>


Zimmet, P. Z., & Alberti, K. G. M. M. (2006). Introduction : Globalization and the Non-

## APPENDIX


### Appendix I: Synthetic Pathway for *p*-Coumaric acid derivatives




**Appendix II :<sup>1</sup>H-NMR spectrum of 3,4',5,7-Tetrahydroxyflavone (kaempferol)  
(CD<sub>3</sub>OD,400 MHz)**




**Appendix III:  $^{13}\text{C}$ -NMR data of 3,4',5,7-Tetrahydroxyflavone (kaempferol) ( $\text{CD}_3\text{OD}$ , 400 MHz)**




**Appendix IV: C13 Dept. Spectra for 3,4',5,7-Tetrahydroxyflavone (kaempferol) (CD<sub>3</sub>OD, 400 MHz)**




**Appendix V: COSY Spectra for 3,4',5,7-Tetrahydroxyflavone (kaempferol) (CD<sub>3</sub>OD, 400 MHz)**





**Appendix VI: HSQC Spectra for 3,4',5,7-Tetrahydroxyflavone (kaempferol) (CD<sub>3</sub>OD, 400 MHz)**




**Appendix VII: HMBC Spectra for 3,4',5,7-Tetrahydroxyflavone (kaempferol) (CD<sub>3</sub>OD, 400 MHz)**




## Appendix VIII: UV Spectrum of 3,4',5,7-Tetrahydroxyflavone (kaempferol).





## Appendix IX: IR Spectra for 3,4',5,7-Tetrahydroxyflavone (kaempferol).




## Appendix X: HRMS Spectra for 3,4',5,7-Tetrahydroxyflavone (kaempferol).








**Appendix XI:  $^1\text{H}$ -NMR spectra of 3,5,7,3',4'- pentahydroxyflavone (Quercetin) ( $\text{CD}_3\text{OD}$ , 400 MHz)**




Appendix XII: <sup>13</sup>C-NMR spectra for 3,5,7,3',4'- pentahydroxyflavone (Quercetin) (CD<sub>3</sub>OD, 400 MHz)




**Appendix XIII: C13 Dept. spectra for 3,5,7,3',4'- pentahydroxyflavone (Quercetin) (CD<sub>3</sub>OD, 400 MHz)**




**Appendix XIV: COSY Spectra for 3,5,7,3',4'- pentahydroxyflavone (Quercetin) (CD<sub>3</sub>OD, 400 MHz)**





**Appendix XV: HSQC Spectra for 3,5,7,3',4'- pentahydroxyflavone (Quercetin) (CD<sub>3</sub>OD, 400 MHz)**




**Appendix XVI: HMBC Spectra for 3,5,7,3',4'- pentahydroxyflavone(Quercetin) (CD<sub>3</sub>OD, 400 MHz)**




## Appendix XVII: UV Spectra for 3,5,7,3',4'- pentahydroxyflavone (Quercetin)




## Appendix XVIII: HRMS Spectra for 3,5,7,3',4'- pentahydroxyflavone (Quercetin)







## Appendix XIX: IR Spectra for 3,5,7,3',4'- pentahydroxyflavone (Quercetin)








Appendix XXI:  $^{13}\text{C}$ -NMR data of 4-Hydroxybenzoic acid ( $\text{CD}_3\text{OD}$ , 100 MHz)




## Appendix XXII: C13 Dept. Spectra for 4 Hydroxybenzoic acid (CD<sub>3</sub>OD, 400 MHz)




### Appendix XXIII: COSY Spectra for 4 Hydroxybenzoic acid (CD<sub>3</sub>OD, 400 MHz)




## Appendix XXIV: HSQC Spectra for 4-Hydroxybenzoic acid (CD<sub>3</sub>OD, 400 MHz)



## Appendix XXV: HMBC Spectra for 4 Hydroxybenzoic acid (CD<sub>3</sub>OD, 400 MHz)





## Appendix XXVI: UV Spectra for 4 Hydroxybenzoic acid



UNIVERSITY of the  
WESTERN CAPE


## Appendix XXVII: HRMS Spectra for 4 Hydroxybenzoic acid





## Appendix XXVIII: IR Spectra for 4 Hydroxybenzoic acid





## Appendix XXIX: $^1\text{H}$ NMR of Ethyl Hydroxycinnamic acid




### Appendix XXX: <sup>1</sup>H NMR of Methyl Hydroxycinnamic acid



# Appendix XXXI: <sup>1</sup>H NMR of Propyl Hydroxycinnamic acid




## Appendix XXXII: HPLC Chromatogram of Fraction M4



UNIVERSITY of the  
WESTERN CAPE

### Appendix XXXIII: HPLC Chromatogram of Fraction M5



## Appendix XXXIV: Certificate of Analysis of *Centella asiatica* (Gotu Kola)



Shaanxi Jiabe Phytochem Co., Ltd. Tel: 0086-29-48344361 Fax: 0086-29-48325119  
A-408 Floor, No.66 Beiyue 1st Road, Xi'an, China 710077  
http://www.jiaherb.com

### CERTIFICATE OF ANALYSIS

#### Gotu Kola Extract 4:1

Batch No.: CJXC-A-822919 Manufacture Date: 20180413 Expiry Date: 20200412

General Information			
Part Used	Grass	Solvents Used	Water & Ethanol
Botanical Source	<i>Centella asiatica</i> (L.) Urb.	Country Of Origin	China
ITEMS	SPECIFICATION	METHOD	TEST RESULTS
Physical & Chemical Data			
Color	Greenish Brown	Organoleptic	Conform
Odour	Characteristic	Organoleptic	Conform
Appearance	Fine Powder	Organoleptic	Conform
Analytical Quality			
Identification	Identical to R.S. sample	HPTLC	Identical
Extract Ratio	4:1		Conform
Sieve analysis	100 % through 80 mesh	USP <786>	Conform
Loss on Drying	≤5.0 %	Eur.Ph.7.0 [2.8.17]	3.75%
Total Ash	≤10.0 %	Eur.Ph. <2.4.16>	2.95%
Bulk density	40-60 g/100mL	Eur.Ph. <2.9.34>	47 g/100mL
Tap density	60-90g/100mL	Eur.Ph. <2.9.34>	69 g/100mL
Contaminants			
Lead (Pb)	≤3.0 mg/kg	Eur.Ph. <2.2.58> ICP-MS	0.0845 mg/kg
Arsenic (As)	≤2.0 mg/kg	Eur.Ph. <2.2.58> ICP-MS	0.1789 mg/kg
Cadmium (Cd)	≤1.0 mg/kg	Eur.Ph. <2.2.58> ICP-MS	0.0784 mg/kg
Mercury (Hg)	≤0.1 mg/kg	Eur.Ph. <2.2.58> ICP-MS	0.0372 mg/kg
Solvents Residue	Meet Eur. Ph.7.0 <5.4>	Eur.Ph. <2.4.24>	Conform
Pesticides Residue	Meet USP Requirements	USP36 <561>	Conform
Microbiological			
Total Plate Count	≤10000 cfu/g	USP36 <2021>	70 cfu/g
Yeast & Mold	≤1000 cfu/g	USP36 <2021>	10 cfu/g
E.Coli.	Negative	USP36 <2022>	Conform
Salmonella	Negative	USP36 <2022>	Conform
General Status			
Non-Irradiation	≤700	EN13751.2002<PSL>	241
Packing & Storage			
	Packed in paper-drum and two plastic-bags inside. N.W.25kgs I.D.35×H51cm. Store in a well-closed container. Away from moisture, light, oxygen.		
Shelf life			
	24 months under the conditions above and in its original packaging.		
Manufacturer			
	Shaanxi Jiabe Phytochem Co., Ltd. Xi'an, P.R. China.		

Quality Assurance Officer



Analyst

75292314  
28153  
ZA10007434

Direktweg South Africa (Pty) Ltd  
11 Marcell Road  
Elsenburg Gardens, Cape Town  
7845  
Tel: 0215561900

No: STP-QCP-<2> (818)

**APPROVED**

Shaanxi Jiabe Phytochem Co., Ltd. Factory Location: NO.66 Hanguang Road, Heping Industrial Park, Xi'an, China 710006